



## A Study of Some Neutrosophic Differential Equations as a Direct Application of One-Dimensional Geometric AH-Isometry

Ahmed A. Salama<sup>1</sup>, Malath F Alaswad<sup>2</sup>, Rasha Dallah<sup>3</sup>, Rozina Ali<sup>4</sup>

Departement of Mathematics and Computer science, port said, Egypt

PHD of Mathematics, GaziAntep, Turkey

Departement of Mathematics, Albaath University, Homs, Syria

Departmrnt Of Mathematics, Cairo University, Egypt

Emails: [drsalama44@gmail.com](mailto:drsalama44@gmail.com); [Malaz.Aswad@yahoo.com](mailto:Malaz.Aswad@yahoo.com); [rasha.dallah20@gmail.com](mailto:rasha.dallah20@gmail.com)

[rozyyy123n@gmail.com](mailto:rozyyy123n@gmail.com)

### Abstract

In this paper, we use the one dimensional AH-isometry to find the structure of the solutions of many neutrosophic differential equations. These equations will be handled by the algebraic direct image of the neutrosophic AH-isometry taken in one dimension.

**Keywords:**One-Dimensional Geometric AH-Isometry; Neutrosophic linear Differential Equation; Neutrosophicreal number.

### 1.Introduction

Neutrosophic logic. Neutrosophy, Neutrosophic set, Neutrosophic probability, are recently creations of Smarandache, being characterized by having the indeterminacy as component of their framework, and a notable feature of neutrosophic logic is that can be considered a generaliazationof fuzzy logics, encompassing the classical logic as well[1]. Also.F. Smarandache, has defined the concept of continuation of a neutrosophic function in year 2015 in[1],and neutrosophicmereo-limit[1], mereo-continuity.Moreover, in 2014, he has defined the concept of a neutro-oscillator differentialin [3], and mereo-derivative. Finallyin 2013 he introduced neutrosophic integration in [2], and mereo-integral, besides the classical defintions of limit, continuity, deverative, and integral respectively. Among the recent applications there are: neutrosophic crisp set theory in image processing,neutrosophic setsmedical field [6-10], in information geographic systems and possible applications to database. Also,neutrosophic triplet group application to physics.Moreover Several researches have made multiple contributions to neutrosophic topology and algebra [14-20, 34-50], Also More researches have made multiple contributions to neutrosophic analysis[21 – 33]. Finally the neutrosophic integration may have application in calculus the areas between two neutrosophic functions.

## 2. Preliminaries

### Definition :Neutrosophic Real Number

Suppose that  $w$  is a neutrosophic number, then it takes the following standard form:  $w = a + bI$  where  $a, b$  are real coefficients, and  $I$  represents the indeterminacy, where  $0 \cdot I = 0$  and  $I^n = I$  for all positive integers  $n$ .

For example:

$$w = 1 + 2I, w = 3 = 3 + 0I.$$

### Definition :Division of neutrosophic real numbers

Suppose that  $w_1, w_2$  are two neutrosophic number, where

$$w_1 = a_1 + b_1I, w_2 = a_2 + b_2I$$

Then:

$$\frac{w_1}{w_2} = \frac{a_1 + b_1I}{a_2 + b_2I} = \frac{a_1}{a_2} + \frac{a_2b_1 - a_1b_2}{a_2(a_2 + b_2)}I$$

### Definition

Let  $R(I) = \{a + bI ; a, b \in R\}$  where  $I^2 = I$  be the neutrosophic field of reals. The one-dimensional isometry (AH-Isometry) is defined as follows:

$$\begin{aligned} T: R(I) &\rightarrow R \times R \\ T(a + bI) &= (a, a + b) \end{aligned}$$

### Remark

$T$  is an algebraic isomorphism between two rings, it has the following properties:

- 1)  $T$  is bijective.
- 2)  $T$  preserves addition and multiplication, i.e.:

$$T[(a + bI) + (c + dI)] = T(a + bI) + T(c + dI)$$

And

$$T[(a + bI) \cdot (c + dI)] = T(a + bI) \cdot T(c + dI)$$

- 3) Since  $T$  is bijective, then it is invertible by:

$$\begin{aligned} T^{-1}: R \times R &\rightarrow R(I) \\ T^{-1}(a, b) &= a + (b - a)I \end{aligned}$$

- 4)  $T$  preserves distances, i.e.:

The distance on  $R(I)$  can be defined as follows:

$$\begin{aligned} \text{Let } A = a + bI, B = c + dI \text{ be two neutrosophic real numbers, then } L = \|\overline{AB}\| &= d[(a + bI, c + dI)] = \\ |a + bI - (c + dI)| &= |(a - c) + I(b - d)| = |a - c| + I[|a + b - c - d| - |a - c|]. \end{aligned}$$

On the other hand, we have:

$$\begin{aligned} T(\|\overline{AB}\|) &= (|a - c|, |(a + b) - (c + d)|) = (d(a, c), d(a + b, c + d)) = d[(a, a + b), (c, c + d)] = d(T(a + \\ bI), T(c + dI)) \\ &= \|T(\overline{AB})\|. \end{aligned}$$

This implies that the distance is preserved up to isometry. i.e.  $\|T(AB)\| = T(\|AB\|)$

**Neutrosophic linear differential equation.**

In this section is defined a linear differential equation by Using the One-Dimensional Geometric AH-Isometry and solutions are found for this equation.

**Neutrosophic identical linear differential equation.****Definition**

Let  $Y = y_1 + y_2I$ ,  $X = x_1 + x_2I$  We define the Neutrosophic identical linear differential equation by Using the One-Dimensional Geometric AH-Isometry as form:

$$\dot{Y} + f(x)Y = 0$$

This equation can be written as follow:

$$\begin{aligned} (y_1' + I[(y_1 + y_2)' - y_1']) & (f(x_1) + I[f(x_1 + x_2) - f(x_1)])(y_1 + y_2I) = 0 \\ \Rightarrow y_1' + f(x_1)y_1 + I[(y_1 + y_2)' - f(x_1 + x_2)(y_1 + y_2) - (y_1' + f(x_1)y_1)] & = 0 \end{aligned}$$

**Method of solution.**

1. Take AH-Isometry for the differential equation, we have.

$$T(y_1' + f(x_1)y_1 + I[(y_1 + y_2)' - f(x_1 + x_2)(y_1 + y_2) - (y_1' + f(x_1)y_1)]) = T(0)$$

$$\Rightarrow (y_1' + f(x_1)y_1, (y_1 + y_2)' - f(x_1 + x_2)(y_1 + y_2)) = (0, 0)$$

Then.

$$\begin{cases} y_1' + f(x_1)y_1 = 0 \dots \dots (1) \\ (y_1 + y_2)' - f(x_1 + x_2)(y_1 + y_2) = 0 \dots \dots (2) \end{cases}$$

The equations (1) and (2) are two differential equations classical.

2. We find the solution to the equations classical (1) and (2), we have.

$y_1$  the solution to the equation (1).

$(y_1 + y_2)$  the solution to the equation (2).

3. We Take invertible AH-Isometry, then, we have the solution of a Neutrosophic identical linear differential equation.

$$Y = y_1 + y_2I = T^{-1}(y_1, y_1 + y_2) = y_1 + ((y_1 + y_2) - y_1)I$$

**Example.** Find a solution to the equation:

$$\dot{Y} + \frac{1}{X}Y = 0$$

**Solution.**

Let  $Y = y_1 + y_2I$ ,  $X = x_1 + x_2I$ . Then.

$$y_1' + \frac{1}{x_1}y_1 + I \left[ (y_1 + y_2)' - \frac{1}{(x_1 + x_2)}(y_1 + y_2) - \left( y_1' + \frac{1}{x_1}y_1 \right) \right] = (0)$$

Now, Take AH-Isometry for the differential equation, we have.

$$\begin{cases} y_1' + \frac{1}{x_1}y_1 = 0 \dots \dots (3) \\ (y_1 + y_2)' - \frac{1}{x_1 + x_2}(y_1 + y_2) = 0 \dots \dots (4) \end{cases}$$

The solution of equation (3) written as follow:

$$y_1 = e^{-\int \frac{1}{x_1} dx_1} + a = e^{-\ln x_1} + a = \frac{1}{x_1} + a, \text{ where } a \in R.$$

The solution of equation (4) written as follow:

$$(y_1 + y_2) = e^{-\int \frac{1}{x_1 + x_2} d(x_1 + x_2)} + b = e^{-\ln(x_1 + x_2)} + b = \frac{1}{x_1 + x_2} + b, \text{ where } b \in R$$

Now, Take invertibleAH-Isometry, then, we have the solution of aNeutrosophic identical linear differential equation

$$Y = y_1 + y_2I = T^{-1} \left( \frac{1}{x_1} + a, \frac{1}{x_1 + x_2} + b \right) = \frac{1}{x_1} + a + \left( \frac{1}{x_1 + x_2} + b - \frac{1}{x_1} - a \right) I$$

$$Y = y_1 + y_2I = \frac{1}{x_1} + \left( \frac{1}{x_1 + x_2} - \frac{1}{x_1} \right) I + a + (b - a)I$$

we have.

$$\frac{1}{x_1} + \left( \frac{1}{x_1 + x_2} - \frac{1}{x_1} \right) I = \frac{1}{x_1} + \left( \frac{-x_2}{x_1(x_1 + x_2)} \right) I = \frac{1}{x_1 + x_2I} = \frac{1}{X}$$

So that,

$$Y = y_1 + y_2I = \frac{1}{X} + (a + bI)$$

where  $a + bI \in R(I)$ .

**Example** Find a solution to the equation:

$$\dot{Y} + 2XY = 0$$

**Solution.**

Let  $Y = y_1 + y_2I$ ,  $X = x_1 + x_2I$ . Then.

$$(y_1' + 2x_1y_1, (y_1 + y_2)' - 2(x_1 + x_2)(y_1 + y_2)) = (0)$$

Now, Take AH-Isometry for the differential equation, we have.

$$\begin{cases} y_1' + 2x_1y_1 = 0 \dots \dots (5) \\ (y_1 + y_2)' - 2(x_1 + x_2)(y_1 + y_2) = 0 \dots \dots (6) \end{cases}$$

The solution of equation (5) written as follow:

$$y_1 = e^{-\int 2x_1 dx_1} + a = e^{-(x_1)^2} + a, \text{ where } a \in R.$$

The solution of equation (6) written as follow:

$$(y_1 + y_2) = e^{-\int 2(x_1+x_2) d(x_1+x_2)} + b = e^{-(x_1+x_2)^2} + b, \text{ where } b \in R$$

Now, Take invertible AH-Isometry, then, we have the solution of a Neutrosophic identical linear differential equation

$$Y = y_1 + y_2I = T^{-1}(e^{-(x_1)^2} + a, e^{-(x_1+x_2)^2} + b) = \frac{1}{x_1} + a + (e^{-(x_1+x_2)^2} + b - e^{-(x_1)^2} - a)I$$

$$Y = y_1 + y_2I = e^{-(x_1)^2} + (e^{-(x_1+x_2)^2} - e^{-(x_1)^2})I + a + (b - a)I$$

we have.

$$e^{-(x_1)^2} + (e^{-(x_1+x_2)^2} - e^{-(x_1)^2})I = e^{-X^2}$$

So that,

$$Y = y_1 + y_2I = e^{-X^2} + (a + bI)$$

where  $a + bI \in R(I)$ .

**Neutrosophic non-homogeneous linear differential equation.**

**Definition**

We define the Neutrosophic non-homogeneous linear differential equation is taken the following forms:

$$\dot{Y} + f(X)Y = g(X)$$

This equation can be written as follow:

$$\begin{aligned} (y_1' + I[(y_1 + y_2)' - y_1'])(f(x_1) + I[f(x_1 + x_2) - f(x_1)])(y_1 + y_2I) &= g(x_1) + I[g(x_1 + x_2) - g(x_1)] \\ \Rightarrow y_1' + f(x_1)y_1 + I[(y_1 + y_2)' - f(x_1 + x_2)(y_1 + y_2) - (y_1' + f(x_1)y_1)] &= g(x_1) + I[g(x_1 + x_2) - g(x_1)] \end{aligned}$$

**Method of solution.**

1. Take AH-Isometry for the differential equation, we have.

$$\begin{aligned} T(y_1' + f(x_1)y_1 + I[(y_1 + y_2)' - f(x_1 + x_2)(y_1 + y_2) - (y_1' + f(x_1)y_1)]) &= T(g(x_1) + I[g(x_1 + x_2) - g(x_1)]) \\ \Rightarrow (y_1' + f(x_1)y_1, (y_1 + y_2)' - f(x_1 + x_2)(y_1 + y_2)) &= (g(x_1), g(x_1 + x_2)) \end{aligned}$$

Then.

$$\begin{cases} y_1' + f(x_1)y_1 = g(x_1) \dots \dots (7) \\ (y_1 + y_2)' - f(x_1 + x_2)(y_1 + y_2) = g(x_1 + x_2) \dots \dots (8) \end{cases}$$

The equations (7) and (8) are two differential equations classical.

2. We find the solution to the equations classical (7) and (8), we have.

$y_1$  the solution to the equation (7).

$(y_1 + y_2)$  the solution to the equation (8).

3. We Take invertible AH-Isometry, then, we have the solution of a Neutrosophic identical linear differential equation .

$$Y = y_1 + y_2 I = T^{-1}(y_1, y_1 + y_2) = y_1 + ((y_1 + y_2) - y_1)I$$

**Example** Find the general solution for the following neutrosophic non- homogeneous linear differential equation:

$$\dot{Y} + 2XY = X$$

**Solution.**

Let  $Y = y_1 + y_2 I, X = x_1 + x_2 I$ . Then.

$$(y_1' + 2x_1 y_1, (y_1 + y_2)' - 2(x_1 + x_2)(y_1 + y_2)) = (x_1, x_1 + x_2)$$

Now, Take AH-Isometry for the differential equation, we have.

$$\begin{cases} y_1' + 2x_1 y_1 = x_1 \dots \dots (9) \\ ((y_1 + y_2)' - 2(x_1 + x_2)(y_1 + y_2)) = x_1 + x_2 \dots \dots (10) \end{cases}$$

The solution of equation (10) written as follow:

$$y_1 = \frac{1}{\mu(x_1)} (a + \int x_1 \mu(x_1) dx) = \frac{1}{e^{(x_1)^2}} (a + \int x_1 e^{(x_1)^2} dx) = \frac{1}{e^{(x_1)^2}} (a + \frac{1}{2} e^{(x_1)^2}), \text{ where } a \in R.$$

By the method same, The solution of equation (10) written as follow:

$$(y_1 + y_2) = \frac{1}{e^{(x_1+x_2)^2}} (b + \frac{1}{2} e^{(x_1+x_2)^2}), \text{ where } b \in R$$

Now, Take invertible AH-Isometry, then, we have the solution of a Neutrosophic identical linear differential equation .

$$\begin{aligned} Y = y_1 + y_2 I &= T^{-1} \left( \frac{1}{e^{(x_1)^2}} (a + \frac{1}{2} e^{(x_1)^2}), \frac{1}{e^{(x_1+x_2)^2}} (b + \frac{1}{2} e^{(x_1+x_2)^2}) \right) \\ &= T^{-1} \left( \frac{1}{e^{(x_1)^2}}, \frac{1}{e^{(x_1+x_2)^2}} \right) \cdot \left[ T^{-1}(a, b) + T^{-1} \left( \frac{1}{2} e^{(x_1)^2}, \frac{1}{2} e^{(x_1+x_2)^2} \right) \right] \end{aligned}$$

$$Y = y_1 + y_2 I = \left[ \frac{1}{e^{(x_1)^2}} + \left( \frac{1}{e^{(x_1+x_2)^2}} - \frac{1}{e^{(x_1)^2}} \right) I \right] \cdot \left[ a + (b - a)I + \frac{1}{2} e^{(x_1)^2} + \left( \frac{1}{2} e^{(x_1+x_2)^2} - \frac{1}{2} e^{(x_1)^2} \right) I \right]$$

By Definition 2.5, we have.

$$\frac{1}{e^{(x_1)^2}} + \left( \frac{1}{e^{(x_1+x_2)^2}} - \frac{1}{e^{(x_1)^2}} \right) I = \frac{1}{e^{(x_1+x_2)I^2}} = \frac{1}{e^{x^2}}$$

$$\text{And } \frac{1}{2} e^{(x_1)^2} + \left( \frac{1}{2} e^{(x_1+x_2)^2} - \frac{1}{2} e^{(x_1)^2} \right) I = \frac{1}{2} [e^{(x_1)^2} + (e^{(x_1+x_2)^2} - e^{(x_1)^2})I] = \frac{1}{2} e^{(x_1+x_2)I^2} = \frac{1}{2} e^{x^2}$$

So that,

$$Y = y_1 + y_2 I = \frac{1}{e^{x^2}} \left[ a + bI + \frac{1}{2} e^{x^2} \right]$$

where  $a + bI \in R(I)$ .

**Example** Find the general solution for the following neutrosophic non- homogeneous linear differential equation:

$$\dot{Y} + \cot g(X)Y = \sin(X)$$

**Solution.**

Let  $Y = y_1 + y_2 I$ ,  $X = x_1 + x_2 I$ . Then.

$$(y_1' + \cot g(x_1)y_1, (y_1 + y_2)') + \cot g(x_1 + x_2)(y_1 + y_2) = (\sin(x_1), \sin(x_1 + x_2))$$

Now, Take AH-Isometry for the differential equation, we have.

$$\begin{cases} y_1' + \cot g(x_1)y_1 = \sin(x_1) \dots \dots (11) \\ ((y_1 + y_2)' + \cot g(x_1 + x_2)(y_1 + y_2) = \sin(x_1 + x_2) \dots \dots (12) \end{cases}$$

The solution of equation (11) written as follow:

$$y_1 = \frac{1}{\mu(x_1)} \left( a + \int x_1 \mu(x_1) dx_1 \right) = \frac{1}{\sin(x_1)} \left( a + \int \sin(x_1) \cdot \sin(x_1) dx \right) = \frac{1}{\sin(x_1)} \left( a + \int \sin^2(x_1) dx \right)$$

$$y_1 = \frac{1}{\sin(x_1)} \left( a + \frac{1}{2} x_1 + \frac{1}{4} \sin(2x_1) \right), \text{ where } a \in R.$$

By the method same, The solution of equation (12) written as follow:

$$(y_1 + y_2) = \frac{1}{\sin(x_1 + x_2)} \left( b + \frac{1}{2} (x_1 + x_2) + \frac{1}{4} \sin(2(x_1 + x_2)) \right), \text{ where } b \in R$$

Now, Take invertible AH-Isometry, then, we have the solution of a Neutrosophic identical linear differential equation

$$Y = y_1 + y_2 I = T^{-1} \left( \frac{1}{\sin(x_1)} \left( a + \frac{1}{2} x_1 + \frac{1}{4} \sin(2x_1) \right), \frac{1}{\sin(x_1 + x_2)} \left( b + \frac{1}{2} (x_1 + x_2) + \frac{1}{4} \sin(2(x_1 + x_2)) \right) \right)$$

$$Y = y_1 + y_2 I = T^{-1} \left( \frac{1}{\sin(x_1)}, \frac{1}{\sin(x_1 + x_2)} \right) \cdot \left[ T^{-1}(a, b) + T^{-1} \left( \frac{1}{2} x_1, \frac{1}{2} (x_1 + x_2) \right) + T^{-1} \left( \frac{1}{4} \sin(2x_1), \frac{1}{4} \sin(2(x_1 + x_2)) \right) \right]$$

$$Y = y_1 + y_2 I = \left[ \frac{1}{\sin(x_1)} + \left( \frac{1}{\sin(x_1 + x_2)} - \frac{1}{\sin(x_1)} \right) I \right] \cdot \left[ a + (b - a)I + \frac{1}{2} x_1 + \left( \frac{1}{2} (x_1 + x_2) - \frac{1}{2} x_1 \right) I + \frac{1}{4} \sin(2x_1) + \left( \frac{1}{4} \sin(2(x_1 + x_2)) - \frac{1}{4} \sin(2x_1) \right) I \right]$$

On the other hand, we have.

$$\frac{1}{\sin(x_1)} + \left( \frac{1}{\sin(x_1 + x_2)} - \frac{1}{\sin(x_1)} \right) I = \frac{1}{\sin(x_1 + x_2 I)} = \frac{1}{\sin(X)}$$

$$\text{And } \frac{1}{2}x_1 + \left(\frac{1}{2}(x_1 + x_2) - \frac{1}{2}x_1\right)I = \frac{1}{2}[x_1 + ((x_1 + x_2) - x_1)I] = \frac{1}{2}(x_1 + x_2I) = \frac{1}{2}X$$

$$\text{And } \frac{1}{4}\sin(2x_1) + \left(\frac{1}{4}\sin(2(x_1 + x_2)) - \frac{1}{4}\sin(2x_1)\right)I = \frac{1}{4}\left[\sin(2x_1) + \left(\sin(2(x_1 + x_2)) - \sin(2x_1)\right)\right]I = \frac{1}{4}\sin 2(x_1 + x_2I) = \frac{1}{4}\sin 2X$$

So that,

$$Y = y_1 + y_2I = \frac{1}{\sin(X)}\left[a + bI + \frac{1}{2}X + \frac{1}{4}\sin 2X\right]$$

where  $a + bI \in R(I)$ .

## Conclusion

In this work we have used the one dimensional AH-isometry to find the solutions of many neutrosophic differential equations. Also, we have illustrated many examples to clarify the validity of our work.

In the future, we aim to study refined neutrosophic differential equations.

## References

- [1] Smarandache F. Neutrosophic Precalculus and Neutrosophic Calculus, University of New Mexico, 705 Gurley Ave. Gallup, NM 87301, USA, 2015.
- [2] Smarandache F. Neutrosophic Measure and Neutrosophic Integral, In Neutrosophic Sets and Systems, 3 – 7, Vol. 1, 2013.
- [3] A. A Salama, Smarandache F. Kroumov, Neutrosophic Closed Set and Continuous Functions, in Neutrosophic Sets and Systems, Vol.4, 4- 8, 2014.
- [4] A. A Salama; I. M Hanafy; Hewayda Elghawalby Dabash M.S, Neutrosophic Crisp Closed Region and Neutrosophic Crisp Continuous Functions, New Trends in Neutrosophic Theory and Applications.
- [5] A. A Salama; Hewayda Elghawalby; M.S, Dabash; A.M. NASR, Retrac Neutrosophic Crisp System For Gray Scale Image, Asian Journal Of Mathematics and Computer Research, Vol 24, 104-117, (2018).
- [6] F. smarandache. "Neutrosophy and Neutrosophic Logic, First International Conference on Neutrosophy, neutrosophic Logic, Set, Probability, and Statistics" University of New Mexico, Gallup, NM 87301, USA 2002.
- [7] M. Abdel-Basset; E. Mai. Mohamed; C. Francisco; H. Z. Abd EL-Nasser. "Cosine similarity measures of bipolar neutrosophic set for diagnosis of bipolar disorder diseases" Artificial Intelligence in Medicine Vol. 101, 101735, (2019).
- [8] Abobala, M., Bal, M., Aswad, M., "A Short Note On Some Novel Applications of Semi Module Homomorphisms", International journal of neutrosophic science, 2022.
- [9] Smarandache F., and Abobala, M., "n-Refined Neutrosophic Vector Spaces", International Journal of Neutrosophic Science, Vol. 7, pp. 47-54. 2020.
- [10] Abobala, M., On Refined Neutrosophic Matrices and Their Applications In Refined Neutrosophic Algebraic Equations, Journal Of Mathematics, Hindawi, 2021
- [11] Abobala, M., A Study of Maximal and Minimal Ideals of n-Refined Neutrosophic Rings, Journal of Fuzzy Extension and Applications, Vol. 2, pp. 16-22, 2021.
- [12] M. Abdel-Basset; E. Mohamed; G. Abdullah; and S. Florentin. "A novel model for evaluation Hospital medical care systems based on plithogenic sets" Artificial Intelligence in Medicine 100 (2019), 101710.
- [13] M. Abdel-Basset; G. Gunasekaran Mohamed; G. Abdullah. C. Victor, "A Novel Intelligent Medical Decision Support Model Based on soft Computing and Iot" IEEE Internet of Things Journal, Vol. 7, (2019).
- [14] M. Abdel-Basset; E. Mohamed; G. Abdullah; G. Gunasekaran; L. Hooang Viet." A novel group decision making model based on neutrosophic sets for heart disease diagnosis" Multimedia Tools and Applications, 1-26, (2019).

- [15] A. A Salama. Basic Structure of Some Classes of Neutrosophic Crisp Nearly Open Sets and Possible Application to GIS Topology. *Neutrosophic Sets and Systems*, Vol. 7, 18-22, (2015).
- [16] A. A Salama; F. Smarandache. Neutrosophic Set Theory, *Neutrosophic Sets and Systems*, Vol. 5, 1-9, (2014).
- [17] F. Smarandache, The Neutrosophic Triplet Group and its Application to physics, Seminar Universidad Nacional de Quilmes , Department of science and Technology, Beunos Aires, Argentina, 20 June 2014.
- [18] A. B.AL-Nafee; R.K. Al-Hamido; F.Smarandache. "Separation Axioms In Neutrosophic Crisp Topological Spaces", *Neutrosophic Sets and Systems*, vol. 25, 25-32, (2019).
- [19] Abobala, M., Bal, M., and Hatip, A.," A Review On Recent Advantages In Algebraic Theory Of Neutrosophic Matrices", *International Journal of Neutrosophic Science*, Vol. 17, 2021.
- [20] R.K. Al-Hamido, Q. H. Imran, K. A. Alghurabi, T. Gharibah, "On Neutrosophic Crisp Semi Alpha Closed Sets", *Neutrosophic Sets and Systems*", vol. 21, 28-35, (2018).
- [21] Sankari, H., and Abobala, M., "Neutrosophic Linear Diophantine Equations With two Variables", *Neutrosophic Sets and Systems*, Vol. 38, pp. 22-30, 2020.
- [22] Sankari, H., and Abobala, M." *n*-Refined Neutrosophic Modules", *Neutrosophic Sets and Systems*, Vol. 36, pp. 1-11. 2020.
- [23] Abobala, M., "AH-Subspaces in Neutrosophic Vector Spaces", *International Journal of Neutrosophic Science*, Vol. 6 , pp. 80-86. 2020.
- [24] Abobala, M.,. "A Study of AH-Substructures in *n*-Refined Neutrosophic Vector Spaces", *International Journal of Neutrosophic Science*", Vol. 9, pp.74-85. 2020.
- [25] Agboola, A.A.A., "On Refined Neutrosophic Algebraic Structures," *Neutrosophic Sets and Systems*,Vol.10, pp. 99-101. 2015.
- [26] Abobala, M., "Classical Homomorphisms Between Refined Neutrosophic Rings and Neutrosophic Rings", *International Journal of Neutrosophic Science*, Vol. 5, pp. 72-75. 2020.
- [27] Q. H. Imran, F. Smarandache, R.K. Al-Hamido, R. Dhavasselán, "On Neutrosophic Semi Alpha open Sets", *Neutrosophic Sets and Systems*, vol. 18, 37-42, (2017).
- [28] Al-Hamido, R. K.; "A study of multi-Topological Spaces", PhD Theses, AlBaath university , Syria, (2019).
- [29] Al-Hamido, R. K.; "Neutrosophic Crisp Supra Bi-Topological Spaces", *International Journal of Neutrosophic Science*, Vol. 1, 66-73, (2018).
- [30] R.K. Al-Hamido, "Neutrosophic Crisp Bi-Topological Spaces", *Neutrosophic Sets and Systems*, vol. 21, 66-73, (2018).
- [31] R.K. Al-Hamido, T. Gharibah, S. JafariF.Smarandache, "On Neutrosophic Crisp Topology via N-Topology", *Neutrosophic Sets and Systems*, vol. 21, 96-109, (2018).
- [32] A. Hatip, "The Special Neutrosophic Functions," *International Journal of Neutrosophic Science (IJNS)*, p. 13, 12 May 2020.\
- [33] Kandasamy, I., Kandasamy, V., and Smarandache, F., "Algebraic structure of Neutrosophic Duplets in Neutrosophic Rings", *Neutrosophic Sets and Systems*, Vol. 18, pp. 85-95. 2018.
- [34] Sankari, H., and Abobala, M.," AH-Homomorphisms In neutrosophic Rings and Refined Neutrosophic Rings", *Neutrosophic Sets and Systems*, Vol. 38, pp. 101-112, 2020.
- [35] Olgun, N., Hatip, A., Bal, M., and Abobala, M., " A Novel Approach To Necessary and Sufficient Conditions For The Diagonalization of Refined Neutrosophic Matrices", *International Journal of Neutrosophic Science*, Vol. 16, pp. 72-79, 2021.
- [36] Aswad, F, M., " A Study of Neutrosophic Complex Number and Applications", *Neutrosophic Knowledge*, Vol. 1, 2020.
- [37] Aswad, M., " A Study of The Integration Of Neutrosophic Thick Function", *International journal of neutrosophic Science*, 2020.
- [38] Abobala, M, "*n*-Cyclic Refined Neutrosophic Algebraic Systems Of Sub-Indeterminacies, An Application To Rings and Modules", *International Journal of Neutrosophic Science*, Vol. 12, pp. 81-95 . 2020.
- [39] M. Ali, F. Smarandache, M. Shabir and L. Vladareanu., "Generalization of Neutrosophic Rings and Neutrosophic Fields", *Neutrosophic Sets and Systems*, vol. 5, pp. 9-14, 2014.
- [40] Smarandache, F., and Abobala, M., *n*-Refined neutrosophic Rings, *International Journal of Neutrosophic Science*, Vol. 5 , pp. 83-90, 2020.

- [41] Abobala, M., " Semi Homomorphisms and Algebraic Relations Between Strong Refined Neutrosophic Modules and Strong Neutrosophic Modules", *Neutrosophic Sets and Systems*, Vol. 39, 2021.
- [42] Abobala, M., "On Some Neutrosophic Algebraic Equations", *Journal of New Theory*, Vol. 33, 2020.
- [43] Abobala, M., "On The Representation of Neutrosophic Matrices by Neutrosophic Linear Transformations", *Journal of Mathematics*, Hindawi, 2021.
- [44] Abobala, M., "On Some Algebraic Properties of n-Refined Neutrosophic Elements and n-Refined Neutrosophic Linear Equations", *Mathematical Problems in Engineering*, Hindawi, 2021
- [45] Kandasamy V, Smarandache F., and Kandasamy I., *Special Fuzzy Matrices for Social Scientists* . Printed in the United States of America,2007, book, 99 pages.
- [46] Abobala, M., *Partial Foundation of Neutrosophic Number Theory*, *Neutrosophic Sets and Systems*, Vol. 39 , 2021.
- [47] Giorgio, N, Mehmood, A., and Broumi, S.," Single Valued neutrosophic Filter", *International Journal of Neutrosophic Science*, Vol. 6, 2020.
- [48] Abobala, M., Hatip, A., Olgun, N., Broumi, S., Salama, A,A., and Khaled, E, H., "The algebraic creativity In The Neutrosophic Square Matrices", *Neutrosophic Sets and Systems*, Vol. 40, pp. 1-11, 2021.
- [49] Abobala, M., "Neutrosophic Real Inner Product Spaces", *Neutrosophic Sets and Systems*, Vol. 43, 2021.
- [50] Abobala, M., "On Some Special Elements In Neutrosophic Rings and Refined Neutrosophic Rings", *Journal of New Theory*, vol. 33, 2020.
- [51] Chellamani, P., and Ajay, D., "Pythagorean neutrosophic Fuzzy Graphs", *International Journal of Neutrosophic Science*, Vol. 11, 2021.
- [52] Abobala, M., "On The Characterization of Maximal and Minimal Ideals In Several Neutrosophic Rings", *Neutrosophic Sets and Systems*, Vol. 45, 2021.
- [53] Abobala, M., Hatip, A., Bal,M.," A Study Of Some Neutrosophic Clean Rings", *International journal of neutrosophic science*, 2022.