



A Study on Compact Operators in Locally K -Convex Spaces

Karla Zayood

Online Islamic University, Department Of Science and Information Technology, Doha, Qatar

Email: zayyyoood134@gmail.com

Abstract

In this paper we give an equivalent definition of continuous and compact linear operators by using orthogonal bases in non-archimedean locally K - convex spaces. We also show that if E is a d_1 space and F is a semi-Montel d_2 space, then every continuous linear operator $T: E \rightarrow F$ is compact.

Keywords: Operator; Convex space; Compact set

1. Introduction

In archimedean functional analysis V. P. Zahariuta showed in [7] that if E, F are locally convex space with absolute bases, E is d_1 space and F is a Montel d_2 space, then every continuous linear operator $T: E \rightarrow F$ is compact.

In this paper we prove this result in non-archimedean functional analysis with E, F are locally K - convex spaces with orthogonal basis.

Throughout K is a non-archimedean valued field that is complete under the metric induced by non-trivial valuation $|\cdot|: K \rightarrow [0, \infty)$.

Let E be a K - vector space. A non-archimedean seminorm on E is a map $\|\cdot\|: E \rightarrow R$ satisfying

- (i) $\|x\| \in \overline{|K|}$ Where $\overline{|K|}$ is the closure of $\{|\lambda|: \lambda \in K\}$ in R .
- (ii) $\|\lambda x\| = |\lambda| \|x\|$
- (iii) $\|x + y\| \leq \max(\|x\|, \|y\|)$

For all $x, y \in E, \lambda \in K$.

In the sequel $E, F, \dots$ are locally K - convex Hausdorff spaces over K , the topology of which is determined by a family $(\|\cdot\|_k)_k$ of non-archimedean seminorms. For the elementary notion and notations concerning locally K - vector spaces we refer to [2], [5] and [6]. By $L(E, F)$ will denote a K - vector spaces consisting of all continuous linear maps.

1.1 Definition Let E be a locally K - convex space. A sequence (x_n) in E is called a (topological) base of E if each $x \in E$ can be written uniquely as $x = \sum_{n=1}^{\infty} \lambda_n x_n$ with $\lambda_n \in K$. If the coefficient functionals $f_n: x \in E \rightarrow \lambda_n \in N, (n \in N)$ are continuous, then (x_n) is called a Schauder basis (see[4]).

1.2 Definitions (see [1])

A basis (x_i) of a locally K - convex space E is said to be an orthogonal basis if the topology of E can be determined by a sequences $(|\cdot|_p)$ of nonarchimedean seminorms satisfying the conditions:

If $x \in E, x = \sum_{i=1}^{\infty} \lambda_i x_i$, then $|x|_p = \max_i |\lambda_i x_i|_p \forall p \in N$.

Note that every orthogonal basis is a Schauder basis.

2. Compact operators:

A compactoid in a locally K - convex space E is a subset B of E such that for every zero-neighborhood U in E , there exists a finite set $A \subset E$ such that $B \subset U + co(A)$, where $co(A)$ is the absolutely convex hull of A . E is called semi-Montel if every bounded subset of E is compactoid. An operator $T \in L(E, F)$ is called compact if there exists a zero- neighborhood U in E such that $T(U)$ is compactoid in F (see[3]).

2.1 Theorem: Let E, F be two locally K - convex space and let $(|\cdot|_k), (\|\cdot\|_k)$, be two sequences of non-archimedean seminorms, defining the topologies on E and F respectively. Then

- a) $T \in L(E, F)$ if and only if $\forall m \exists k = k(m)$ and $C(m) \in K$ with $|C(m)| > 0$ such that $\forall x \in E$
 $\|Tx\|_m \leq |C(m)| |x|_k$
- b) If (e_n) is any orthogonal basis in E , then $T \in L(E, F)$ if and only if $\forall m \exists k = k(m)$ such that

$$\sup_n \frac{\|Te_n\|_m}{|e_n|_k} < \infty$$

c) If F is a semi-Montel space, then $T: E \rightarrow F$ is a compact operator if and only if $\exists k \forall m \exists D(m) \in K$ with $|D(m)| > 0$ such that $\forall x \in E$

$$\|Tx\|_m \leq |D(m)| |x|_k$$

d) If e_n is any orthogonal basis in E and F is a semi-Montel space, then $T: E \rightarrow F$ is a compact operator if and only if $\exists k \forall m$

$$\sup_n \frac{\|Te_n\|_m}{|e_n|_k} < \infty$$

Proof : a)

$T \in L(E, F)$ if and only if for the a zero-neighborhood $B_{\|\cdot\|_m}(0,1)$ in F , there exist zero-neighborhood $B_{|\cdot|_k}(0,1)$ in E and $C'(m) \in K$ with such $|C'(m)| > 0$ that $T(C'(m) \cdot B_{|\cdot|_k}(0,1)) \subseteq B_{\|\cdot\|_m}(0,1)$. Now if $x \in E$, then $\left| \frac{C'(m)}{2} \frac{x}{|x|_k} \right|_k = \left| \frac{C'(m)}{2} \right| \frac{|x|_k}{|x|_k} = \left| \frac{C'(m)}{2} \right| < |C'(m)|$, and so $\frac{C'(m)}{2} \frac{x}{|x|_k} \in C'(m) \cdot B_{|\cdot|_k}(0,1)$ and this implies that

$$T\left(\frac{C'(m)}{2} \frac{x}{|x|_k}\right) \in T(C'(m) \cdot B_{|\cdot|_k}(0,1)) \subseteq B_{\|\cdot\|_m}(0,1).$$

$$\text{Thus, } \left\| \frac{C'(m)}{2} \frac{x}{|x|_k} \right\|_m \leq 1, \text{ and so } \|Tx\|_m \leq |C(m)| |x|_k \text{ where } |C(m)| = \left| \frac{2}{C'(m)} \right|.$$

b) ($\Rightarrow$) Let $T \in L(E, F)$. Then by part (a) $\forall m \exists k = k(m)$ and $C(m) \in K$ with $|C(m)| > 0$ such that

$$\|Tx\|_m \leq |C(m)| |x|_k \quad \forall x \in E$$

Since $e_n \in E$, then $\forall m \exists k = k(m)$ and $C(m) \in K$ with $|C(m)| > 0$ such that

$$\|Te_n\|_m \leq |C(m)| |e_n|_k \quad \forall n \in N$$

Hence $\forall m \exists k = k(m)$ and $C(m) \in K$ with $|C(m)| > 0$ such that

$$\frac{\|Te_n\|_m}{|e_n|_k} \leq |C(m)| \quad \forall n \in N$$

It follows that, $\forall m \exists k = k(m)$ such that

$$\sup_n \frac{\|Te_n\|_m}{|e_n|_k} < \infty$$

($\Leftarrow$) Let $T: E \rightarrow F$ be any linear operator and (e_n) be any orthogonal basis in E . Suppose that $\forall m \exists k = k(m)$ such that

$$\sup_n \frac{\|Te_n\|_m}{|e_n|_k} < \infty$$

Then $\forall m \exists k = k(m)$ and $C(m) \in K$ with $|C(m)| > 0$ such that

$$\|Te_n\|_m \leq |C(m)| |e_n|_k \quad \forall n \in N$$

Now if $x \in E$. Then x has a unique representation $x = \sum_{n=1}^{\infty} \lambda_n e_n$. Since

$$Tx = \sum_{n=1}^{\infty} \lambda_n Te_n, \text{ then}$$

$$\|Tx\|_m = \left\| \sum_{n=1}^{\infty} \lambda_n Te_n \right\|_m \leq \max_n |\lambda_n| \|Te_n\|_m \leq \max_n |C(m)| |\lambda_n| |e_n|_k = |C(m)| |x|_k.$$

and hence $T \in L(E, F)$

c) $T: E \rightarrow F$ is compact if and only if there exists a zero-neighborhood $B_{|\cdot|_k}(0,1)$ in E , such that $T(B_{|\cdot|_k}(0,1))$ is compact in F . Since F is semi-Montel space, then $T(B_{|\cdot|_k}(0,1))$ is bounded in F . So there exist a zero-neighborhood $B_{\|\cdot\|_m}(0,1)$ in F and $D'(m) \in K$ with $|D'(m)| > 0$ such that $T(B_{|\cdot|_k}(0,1)) \subseteq D'(m) \cdot B_{\|\cdot\|_m}(0,1)$.

Now let $x \in E$, then $\left| \frac{1}{2} \frac{x}{|x|_k} \right|_k = \frac{1}{2} < 1$, and so $\frac{1}{2} \frac{x}{|x|_k} \in B_{|\cdot|_k}(0,1)$ and this implies that $T\left(\frac{1}{2} \frac{x}{|x|_k}\right) \in T(B_{|\cdot|_k}(0,1)) \subseteq D'(m) \cdot B_{\|\cdot\|_m}(0,1)$. Thus $\left\| \frac{1}{2} \frac{Tx}{|x|_k} \right\|_m \leq |D'(m)|$, and so $\|Tx\|_m \leq |D(m)| |x|_k$ where $|D(m)| = |2D'(m)|$.

d) ($\Rightarrow$) let $T: E \rightarrow F$ be any compact operator. Then by part (c) $\exists k \forall m \exists D(m) \in K$ with $|D(m)| > 0$ such that $\forall x \in E$

$$\|Tx\|_m \leq |D(m)| |x|_k$$

Since $e_n \in E$, then $\exists k \forall m \exists D(m) \in K$ with $|D(m)| > 0$ such that

$$\|Te_n\|_m \leq |D(m)| |e_n|_k \quad \forall n \in N.$$

Hence $\exists k \forall m \exists D(m) \in K$ with $|D(m)| > 0$ such that

$$\frac{\|Te_n\|_m}{|e_n|_k} \leq |D(m)| \quad \forall n \in N$$

It follows that, $\exists k \forall m$ such that

$$\sup_n \frac{\|Te_n\|_m}{|e_n|_k} < \infty$$

($\Leftarrow$) Let $T: E \rightarrow F$ be any linear operator and (e_n) be any orthogonal basis in E . Suppose that $\exists k \forall m$ such that

$$\sup_n \frac{\|Te_n\|_m}{|e_n|_k} < \infty$$

then $\exists k \forall m \exists D(m) \in K$ with $|D(m)| > 0$ such that

$$\|Te_n\|_m \leq |D(m)| |e_n|_k \quad \forall n \in N.$$

Now if $x \in E$, then x has a unique representation $x = \sum_{n=1}^{\infty} \lambda_n e_n$. Since

$$Tx = \sum_{n=1}^{\infty} \lambda_n Te_n,$$

$$\|Tx\|_m = \left\| \sum_{n=1}^{\infty} \lambda_n Te_n \right\|_m \leq \max_n |\lambda_n| \|Te_n\|_m \leq \max_n |D(m)| |\lambda_n| |e_n|_k = |D(m)| |x|_k$$

And so $T: E \rightarrow F$ is a compact operator.

2.2 Definition

We shall say that a locally K -convex space $E \in d_i, i = 1, 2$ if there exists in E an orthogonal basis x_n such that

$$\exists p \forall q \exists r \text{ such that } |x_k|_q^2 \leq |x_k|_p |x_k|_r \quad \forall k \in N \text{ for } i=1, \text{ and}$$

$$\forall q \exists p \forall r \text{ such that } |x_k|_q^2 \geq |x_k|_p |x_k|_r \quad \forall k \in N \text{ for } i=2.$$

2.3 Theorem

Let $E \in d_2, F \in d_1$ and F be a semi-Montel space. Then every continuous linear operator $T: E \rightarrow F$ is compact.

Proof: Let $T: E \rightarrow F$ be an arbitrary continuous linear operator. Since $E \in d_2, F \in d_1$, then there exist two orthogonal bases $(x_n), (y_n)$ in E, F respectively and sequences of non-archimedean seminorms $(|\cdot|_p), (\|\cdot\|_q)$ defining the topology on E, F respectively, satisfying the conditions:

$$\text{if } x \in E, x = \sum_{i=1}^{\infty} \lambda_i x_i, \text{ then } |x|_p = \max_i |\lambda_i x_i|_p \quad \forall p \in N, \text{ and}$$

$$\text{if } x \in F, x = \sum_{i=1}^{\infty} \lambda_i x_i, \text{ then } \|x\|_p = \max_i \|\lambda_i x_i\|_p \quad \forall p \in N.$$

Now let $Tx_k = \sum_{i=1}^{\infty} t_{ik} y_i$. Since $x = \sum_{k=1}^{\infty} \lambda_k x_k$ and T is continuous, then

$$Tx = \sum_{i=1}^{\infty} \eta_i y_i, \text{ where } \eta_i = \sum_{k=1}^{\infty} t_{ik} \lambda_k, \text{ that is } T \text{ can be represented by the}$$

Matrix (t_{ik}) . Now Since T is continuous, by proposition (2.1(a)) $\forall p \exists q = q(p)$ and $C(p) \in K$ with $|C(p)| > 0$ such that $\forall x \in E$

$$\|Tx\|_p \leq |C(p)| |x|_q.$$

It follows that

$$\left\| \sum_{i=1}^{\infty} \eta_i y_i \right\|_p = \left\| \sum_{i=1}^{\infty} \left(\sum_{k=1}^{\infty} t_{ik} \lambda_k \right) y_i \right\|_p \leq |C(p)| \left\| \sum_{i=1}^{\infty} \lambda_i x_i \right\|_q$$

And so,

$$\max_i \left\| \left(\sum_{k=1}^{\infty} t_{ik} \lambda_k \right) y_i \right\|_p \leq |C(p)| \max_i |\lambda_i x_i|_q$$

Hence

$$\frac{\max_i \left\| \left(\sum_{k=1}^{\infty} t_{ik} \lambda_k \right) y_i \right\|_p}{\max_i |\lambda_i x_i|_q} \leq |C(p)|$$

Now if $x = x_k$, then

$$\frac{\max_i |t_{ik}| \|y_i\|_p}{|x_k|_q} \leq |C(p)|$$

Taking the supremum over k , we have

$$\sup_k \frac{\max_i |t_{ik}| \|y_i\|_p}{|x_k|_q} \leq |C(p)| \quad (1)$$

Now since F is a semi-Montel space, to prove that T is compact, it sufficient to show that for some neighborhood $U_{q_0} = \{x \in X: |x|_{q_0} \leq 1\}$ in $E, T(U_{q_0})$ is bounded in F .i.e. for all $V_p = \{y \in F: |y|_p \leq 1\}, p \in N$

There exists $M(p) \in K$ with $|M(p)| > 0$ such that $T(U_{q_0}) = \{Tx \in F: |x|_{q_0} \leq 1\} \subseteq M(p)V_p$, that is if $x \in U_{q_0}$ then for all $p \in N$,

$$\|Tx\|_p = \max_i \left\| \left(\sum_{k=1}^{\infty} t_{ik} \lambda_k \right) y_i \right\|_p \leq |M(p)|$$

Now let $x \in E$, then $\left| \frac{x}{2|x|_{q_0}} \right|_{q_0} = \frac{1}{2} < 1$. Hence,

$$T\left(\frac{x}{2|x|_{q_0}}\right) \in T(U_{q_0}) \subseteq M(p)V_p. \text{ So } \frac{\|Tx\|_p}{|x|_{q_0}} \leq 2|M(p)|, \text{ and hence}$$

$$\frac{\max_i \left\| \left(\sum_{k=1}^{\infty} t_{ik} \lambda_k \right) y_i \right\|_p}{\max_i |\lambda_i x_i|_{q_0}} \leq 2|M(p)|. \text{ Now if we take } x = x_k, \text{ then}$$

$$\frac{\max_i |t_{ik}| \|y_i\|_p}{|x_k|_{q_0}} \leq 2|M(p)|$$

Taking the supremum over k we have

$$\sup_k \frac{\max_i |t_{ik}| \|y_i\|_p}{|x_k|_{q_0}} \leq 2|M(p)| \quad (2)$$

To prove $T(U_{q_0})$ is bounded in F we want to prove that (2) holds.

Now, since $F \in d_1$, then

$$\exists p_1 \forall p \exists p_2 = p_2(p) \text{ such that } \|y_i\|_p^2 \leq \|y_i\|_{p_1} \|y_i\|_{p_2} \quad \forall i \in N, \quad (3)$$

And since $E \in d_2$, then

$$\forall q_1 = q(p_1) \exists q = q_0 \forall q_2 = q(p_2) \text{ such that } |x_k|_{q_0}^2 \geq |x_k|_{q_1} |x_k|_{q_2} \quad \forall k \in N, \quad (4)$$

Now by (1), (3), (4), we have

$$\begin{aligned} \max_i \frac{|t_{ik}| \|y_i\|_p}{|x_k|_{q_0}} &\leq \left(\max_i \frac{|t_{ik}|^2 (\|y_i\|_p)^2}{(|x_k|_{q_0})^2} \right)^{\frac{1}{2}} \leq \left(\max_i \frac{|t_{ik}|^2 \|y_i\|_{p_1} \|y_i\|_{p_2}}{|x_k|_{q_1} |x_k|_{q_2}} \right)^{\frac{1}{2}} \\ &\leq \left(\max_i \frac{|t_{ik}| \|y_i\|_{p_1}}{|x_k|_{q_1}} \right)^{\frac{1}{2}} \left(\max_i \frac{|t_{ik}| \|y_i\|_{p_2}}{|x_k|_{q_2}} \right)^{\frac{1}{2}} \leq \left(|C(p_1)|^{\frac{1}{2}} \right) \left(|C(p_2)|^{\frac{1}{2}} \right) < \infty \quad \forall p \in N \end{aligned}$$

If we take the supremum over k , then the inequality (2) holds, and hence T is a compact operator.

References

- [1] N.De Grande-De Kimpe, On the structure of locally K-convex spaces with a Schauder basis, Indag. Math., 34 (1972), 396-406.
- [2] N.De Grande-De Kimpe, C-compactness in locally K-convex spaces, Indag. Math., 33, (1971), 176-180.
- [3] C. C. Perez-Garcia and W.H. Schikhof, Compact operators and the Orlicz-Pettis property in p-adic analysis, Report 9101, Department of Math, Catholic University, Nijmegen, the Netherlands, 1991, 1 27.
- [4] C. Perez-Garcia, On compactoidity in non-Archimedean locally convex spaces with a Schauder basis, Nederl. Akad. Wetensch. Indag., 50,(1988), 85-88.
- [5] W.H. Schikhof, Locally convex spaces over non-spherically complete valued fields, I-II, Bull. Soc. Math. Belgique, XXXVIII (ser. B),38, (1986), 187-224.
- [6] A. C. M. Van Rooij, Non-Archimedean Functional Analysis, Marcel Dekker, New York, (1978).
- [7] V.P. Zahariuta, On the isomorphism of Cartesian products of locally convex spaces, Studia Math. 46, (1973), 201-221.
- [8] Multu, F., Tercan, A., and Yasar, R., Eventually semisimple weak FI-extending modules, Mathematica BOHEMICA, Vol 148, 2023
- [9] Yasar, R., C11-modules via left exact modules preradicals, Turkish journal of mathematics, vol 45, 2021.
- [10] Yasar, R., On weak Projection invariant semisimple modules, Fundamental journal of mathematics and applications, 2021.
- [11] Yasar, R., and Tercan, A., Extending property on EC-fully submodules, Sakarya university Journal of Science, 2018.