



MBJ-Neutrosophic Positive Implicative LI-ideals, Associative LI-ideals, and Fantastic LI-ideals

Amarendra Babu V.¹, Abida Begum K.², Siva Naga Malleswari V.³

¹Department of Mathematics, Acharya Nagarjuna university, Guntur Dist., A.P, India

²Department of H & S, KRUCET, Krishna university, Machilipatnam, A.P, India

³PVP Siddhartha institute of Technology, Vijayawada, A.P, India

Emails: vamarendrababu@anu.ac.in; kruteacher211@kru.ac.in; malleswari.v@pvpsiddhartha.ac.in

Abstract

The MBJ-Neutrosophic positive implicative LI-ideals, the MBJ-Neutrosophic associative LI-ideals, and the MBJ-Neutrosophic fantastic LI-ideals are all introduced in this study. Of all these ideals, several their qualities and corresponding conditions were explored. We have demonstrated how every positive implicative MBJ-Neutrosophic LI-ideal evolved into an MBJ-Neutrosophic LI-ideal, MBJ-Neutrosophic implicative LI-ideal, and MBJ-Neutrosophic fantastic LI-ideal. Additionally, it was demonstrated that any MBJ-Neutrosophic fantastic LI-ideal is a MBJ-Neutrosophic associative LI-ideal.

Keywords: Lattice implication algebra (LIA); MBJ-Neutrosophic LI-ideal (MBJ-NLII), MBJ-Neutrosophic implicative LI-ideal (MBJ-NILII), MBJ-Neutrosophic positive implicative LI-ideal (MBJ-NPILII), MBJ-Neutrosophic associative LI-ideal (MBJ-NALII), and MBJ-Neutrosophic fantastic LI-ideal (MBJ-NFLII) .

1. Introduction:

The concept of Neutrosophic set (Nset), which Smarandache proposed in 1998 [15], includes the interval-valued fuzzy (intuitionistic) set and the standard set (fuzzy). According to [11], "Jun, his colleagues examined, properties and relations of BCK/BCI-algebras using the idea of Neutrosophic set theory" (Nset) was defined after the Neutrosophic elements T, I, and F were innovated. It indicates the truth-membership, indeterminacy, and false-membership values individually. Then, according to [15], "the interval Neutrosophic set was inaugurated. Additionally, "Interval Neutrosophic ideals were interpreted, and several features examined" in [11]. They showed a variety of interval Neutrosophic ideals in [10], examined some of their traits, and found a connection between them.

In 1993, Xu [16] put up the idea of a logical algebra-lattice implication algebra. In lattice implication algebra, the lattice is defined to explain uncertainty, notably beyond comparison, and is intended to capture human reasoning. The use of Neutrosophic sets in lattice implication algebra offers a fresh method for identifying uncertainty while avoiding many of the drawbacks of earlier theoretical methods. Applications of neutrosophic set theory include information theory, probability theory, control theory, decision-making, measurement theory, and others. Since then, a considerable body of literature on the theory of Neutrosophic sets has been published as a result of the work of several researchers in this area. Smarandache introduces the idea of the MBJ-Neutrosophic set as another generalisation of the Neutrosophic set.

In [9], Jun also suggested the notion of LI-ideals and examined a number of their characteristics as well as their interrelationships. The Neutrosophic set theory idea was also used to LIA. The MBJ Neutrosophic set abstraction is a novel generalisation of the N set.

We examined the properties and corresponding circumstances of the MBJ-Neutrosophic LI-Ideals in Lattice Implication Algebras in [1] and introduced them.

The MBJ-Neutrosophic Implicative LI-Ideals in Lattice Implication Algebras are explored in [2], where their features are provided and related relations are analysed.

The MBJ-Neutrosophic positive implicative LI ideal was introduced in section 3.1 of this work, and it was demonstrated that any MBJ-Neutrosophic positive implicative LI ideal is both an MBJ-Neutrosophic LI ideal and an MBJ-Neutrosophic implicative LI-ideal. We introduced the MBJ-Neutrosophic associative LI-ideal in section 3.2. We study the MBJ-Neutrosophic fantastic LI-ideal of a LIA in section 3.3 and look at a number of its features. Additionally, it demonstrated how these ideals linked to one another.

2. Related Work:

Definition-2.1. ([18]) : LIA is described as a Lattice $(L, \vee, \wedge, 0, 1)$ with bounds $(0, 1)$, order-reversing involution “’” comforting propositions given below under a binary operation “ $\rightarrow$ ” are

$$(I1) \quad \zeta \rightarrow (q \rightarrow \varphi) = q \rightarrow (\zeta \rightarrow \varphi),$$

$$(I2) \quad \zeta \rightarrow \zeta = 1,$$

$$(I3) \quad \zeta \rightarrow q = q' \rightarrow \zeta',$$

$$(I4) \quad \zeta \rightarrow q = q \rightarrow \zeta = 1 \Rightarrow \zeta = q,$$

$$(I5) \quad (\zeta \rightarrow q) \rightarrow q = (q \rightarrow \zeta) \rightarrow \zeta,$$

$$(L1) \quad (\zeta \vee q) \rightarrow \varphi = (\zeta \rightarrow \varphi) \wedge (q \rightarrow \varphi),$$

$$(L2) \quad (\zeta \wedge q) \rightarrow \varphi = (\zeta \rightarrow \varphi) \vee (q \rightarrow \varphi),$$

A partial ordering $\leq$ on ‘L’ defined by the condition $\zeta \leq q$ iff $\zeta \rightarrow q = 1$.

A LIA ‘L’ satisfies the propositions cited below (see [12]):

$$(a1) \quad 0 \rightarrow \zeta = 1, 1 \rightarrow \zeta = \zeta \text{ \& } \zeta \rightarrow 1 = 1.$$

$$(a2) \quad \zeta \rightarrow q \leq (q \rightarrow \varphi) \rightarrow (\zeta \rightarrow q).$$

$$(a3) \quad \zeta \leq q \text{ implies } q \rightarrow \varphi \leq \zeta \rightarrow \varphi \text{ and } \varphi \rightarrow \zeta \leq \varphi \rightarrow q.$$

$$(a4) \quad \zeta' = \zeta \rightarrow 0.$$

$$(a5) \quad \zeta \vee q = (\zeta \rightarrow q) \rightarrow q.$$

$$(a6) \quad ((q \rightarrow \zeta) \rightarrow \zeta')' = \zeta \wedge q = ((\zeta \rightarrow q) \rightarrow \zeta')'.$$

$$(a7) \quad \zeta \leq (\zeta \rightarrow q) \rightarrow q.$$

$$(a8) \quad ((\zeta \rightarrow q) \rightarrow q) \rightarrow q = \zeta \rightarrow q$$

Definition 2.2. ([9]) : A LI-ideal of ‘L’ is a nonempty subset χ of ‘L’ holds the propositions given by

$$(L3) \quad 0 \in \chi.$$

$$(L4) \quad (\forall \zeta \in 'L') (\forall q \in \chi) ((\zeta \rightarrow q) \in \chi \Rightarrow \zeta \in \chi).$$

MBJ-Neutrosophic LI-ideals:

We suppose ‘L’ as LIA except as otherwise indicated.

Definition 2.3. ([1]) : A MBJ-NLII of ‘L’ is an MBJ-Nset

$A = (M_A, \widetilde{B}_A, J_A)$ in ‘L’ if the following attributes are true.

$$(\forall \zeta \in L) (M_A(0) \geq M_A(\zeta), \widetilde{B}_A(0) \geq \widetilde{B}_A(\zeta), J_A(0) \leq J_A(\zeta))$$

$$\text{and } (\forall \varsigma, \varrho \in L) \begin{pmatrix} M_A(\varsigma) \geq \min\{M_A((\varsigma \rightarrow \varrho)'), M_A(\varrho)\} \\ \widetilde{B}_A(\varsigma) \geq \min\{\widetilde{B}_A((\varsigma \rightarrow \varrho)'), \widetilde{B}_A(\varrho)\} \\ J_A(\varsigma) \leq \max\{J_A((\varsigma \rightarrow \varrho)'), J_A(\varrho)\} \end{pmatrix}$$

We use the notation MBJ-NLI 'L' to represent the collection of all MBJ-NLI-ideals of 'L'.

Definition 2.4. ([2]) : An implicative LI-Ideal of 'L' is a nonempty subset A of 'L' which satisfies $0 \in L$ & $\forall \varsigma, \varrho, \varphi \in L, (((\varsigma \rightarrow \varrho)' \rightarrow \varrho)' \rightarrow \varphi)' \in A, \varphi \in A \Rightarrow (\varsigma \rightarrow \varrho)' \in A$.

Definition 2.5. ([2]) : **MBJ-Neutrosophic Implicative LI-ideal** : An MBJ-N set 'A' of 'L' is known as an MBJ-NILII of 'L' if it holds the inequalities given below for all $\varsigma \in L$, we have

$$(M_A(0) \geq M_A(\varsigma), \widetilde{B}_A(0) \geq \widetilde{B}_A(\varsigma), J_A(0) \leq J_A(\varsigma)), \forall \varsigma, \varrho, \varphi \in L \text{ and}$$

$$\begin{cases} M_A((\varsigma \rightarrow \varrho)') \geq \min\{M_A((((\varsigma \rightarrow \varrho)' \rightarrow \varrho)' \rightarrow \varphi)'), M_A(\varphi)\}, \\ \widetilde{B}_A((\varsigma \rightarrow \varrho)') \geq \min\{\widetilde{B}_A((((\varsigma \rightarrow \varrho)' \rightarrow \varrho)' \rightarrow \varphi)'), \widetilde{B}_A(\varphi)\}, \\ J_A((\varsigma \rightarrow \varrho)') \leq \max\{J_A((((\varsigma \rightarrow \varrho)' \rightarrow \varrho)' \rightarrow \varphi)'), J_A(\varphi)\}. \end{cases}$$

3. MBJ-Neutrosophic

3.1. MBJ-Neutrosophic positive implicative LI-ideals :

3.1.1. Definition :

Let X be the universe. A MBJ-Neutrosophic set $A = \{(\varsigma; M_A(\varsigma), \widetilde{B}_A(\varsigma), J_A(\varsigma)) / \varsigma \in X\}$ of a LIA 'L' is called a MBJ-Neutrosophic positive implicative LI-ideal of 'L' if it satisfies

$$M_A(0) \geq M_A(\varsigma), \widetilde{B}_A(0) \geq \widetilde{B}_A(\varsigma), J_A(0) \leq J_A(\varsigma)$$

$$M_A(\varrho) \geq \min\{M_A((\varrho \rightarrow (\varphi \rightarrow \varrho)')' \rightarrow \varsigma)'), M_A(\varsigma)\}$$

$$\widetilde{B}_A(\varrho) \geq \min\{\widetilde{B}_A((\varrho \rightarrow (\varphi \rightarrow \varrho)')' \rightarrow \varsigma)'), \widetilde{B}_A(\varsigma)\}$$

$$J_A(\varrho) \leq \max\{J_A((\varrho \rightarrow (\varphi \rightarrow \varrho)')' \rightarrow \varsigma)'), J_A(\varsigma)\}$$

3.1.2. Example : Let $L = \{0, a, b, c, 1\}$. Define partial order on L as $0 < a < b < c < 1$ and define $\varsigma \wedge \varrho = \min\{\varsigma, \varrho\}$ and $\varsigma \vee \varrho = \max\{\varsigma, \varrho\}$ for all $\varsigma, \varrho \in L$ and " ' " and " $\rightarrow$ " as follows

Table 1: Illustration of Complement of element

ς	ς'
0	1
a	c
b	b
c	a
1	0

Table 2: Illustration of Implication

$\rightarrow$	0	a	b	c	1
---------------	---	---	---	---	---

0	1	1	1	1	1
a	c	1	1	1	1
b	b	c	1	1	1
c	a	b	c	1	1
1	0	a	b	c	1

Then $(L, \vee, \wedge, ', \rightarrow)$ is LIA.

Define MBJ-Neutrosophic set A in L by Tabular representation of A.

Table 3: Illustration of Neutrosophic set $A = (M_A, \widetilde{B}_A, J_A)$

L	0	a	b	c	1
M_A	0.7	0.7	0.7	0.5	0.5
$\widetilde{B}_A$	[0.3, 0.5]	[0.3, 0.5]	[0.3, 0.5]	[0.2, 0.3]	[0.2, 0.3]
J_A	0.5	0.5	0.7	0.7	0.7

Then $A = (M_A, \widetilde{B}_A, J_A)$ is a MBJ-NPILII on L,

As it satisfies all the attributes of the MBJ-NPILII.

3.1.3. Theorem: Every MBJ-NPILII of a lattice implication algebra is a MBJ-NLII.

Proof : Let A be a MBJ-NPILII of a LIA 'L'.

$$\text{Then } M_A(0) \geq M_A(\varsigma), \widetilde{B}_A(0) \geq \widetilde{B}_A(\varsigma),$$

$$J_A(0) \leq J_A(\varsigma)$$

$$\text{And let } \varphi = q \text{ then } M_A(q) \geq \min\{M_A((q \rightarrow (q \rightarrow q)')' \rightarrow \varsigma)'), M_A(\varsigma)\}$$

$$= \min\{M_A((q \rightarrow 0)')' \rightarrow$$

$$\varsigma)'), M_A(\varsigma)\}$$

$$= \min\{M_A((q')' \rightarrow$$

$$\varsigma)'), M_A(\varsigma)\}$$

$$= \min\{M_A(q \rightarrow \varsigma)'), M_A(\varsigma)\}$$

$$q)'), \widetilde{B}_A(\varsigma)\}$$

$$\widetilde{B}_A(q) \geq r \min\{\widetilde{B}_A((q \rightarrow (q \rightarrow$$

$$\varsigma)'), \widetilde{B}_A(\varsigma)\}$$

$$\varsigma)'), \widetilde{B}_A(\varsigma)\}$$

$$= r \min\{\widetilde{B}_A((q')' \rightarrow$$

$$\varsigma)'), \widetilde{B}_A(\varsigma)\}$$

$$\begin{aligned}
& \zeta)' , \widetilde{B}_A(\zeta)\} \\
& \zeta)' \rightarrow \zeta)' , J_A(\zeta)\} \\
& \zeta)' , J_A(\zeta)\} \\
& \zeta)' , J_A(\zeta)\} \\
& = r \min\{\widetilde{B}_A(q \rightarrow \\
& J_A(q) \leq \max\{J_A((q \rightarrow (q \rightarrow \\
& = \max\{J_A((q \rightarrow 0)' \rightarrow \\
& = \max\{J_A((q')' \rightarrow \\
& = \max\{J_A(q \rightarrow \zeta)' , J_A(\zeta)\}
\end{aligned}$$

Therefore, A is a MBJ-NLII.

Note: Converse of the above theorem is need not be true. It will be followed by the below example.

3.1.4. Example: The Hasse diagram and Cayley tables of a poset, $L = \{0, a, b, c, d, 1\}$ are given below as follows:




Figure 1: Hasse diagram of L

Table 4: Illustration of compliment of element

a	b	ζ	ζ'
		0	1
		a	c
		b	d
		c	a
		d	b
		1	0

Table 5: Illustration of Implication

$\rightarrow$	0	a	b	c	d	1
0	1	1	1	1	1	1
a	c	1	b	c	b	1
b	d	a	1	b	a	1
c	a	a	1	1	a	1
d	b	1	1	b	1	1
1	0	a	b	c	d	1

The operations $\vee$ and $\wedge$ on L are defined as follows

$$\zeta \vee q = (\zeta \rightarrow q) \rightarrow q, \zeta \wedge q = ((\zeta' \rightarrow q') \rightarrow q')', \text{ for all } \zeta, q \in L.$$

Clearly we can observe that L is a LIA.

Then $(L, \vee, \wedge, ', \rightarrow)$ is LIA.

Define MBJ-Neutrosophic set A in L by

Table 6 : Illustration of Neutrosophic set $A = (M_A, \widetilde{B}_A, J_A)$

L	0	a	b	c	d	1
M_A	0.7	0.5	0.5	0.7	0.5	0.5
$\widetilde{B}_A$	[0.5, 0.6]	[0.3, 0.4]	[0.3, 0.4]	[0.3, 0.4]	[0.5, 0.6]	[0.3, 0.4]
J_A	0.3	0.5	0.5	0.5	0.3	0.5

Then A is a MBJ-NLII but it is not a M $\widetilde{B}$ J-NPILII on L, because

For $0, d, a \in L$, we can observe that

$$M_A(d) \not\geq \min\{M_A(((d \rightarrow (a \rightarrow (d)'))' \rightarrow 0)'), M_A(0)\}$$

$$0.5 \not\geq 0.7.$$

3.1.5. Theorem: Let A be a MBJ-NLII of a LIA 'L'. Then A is a M $\widetilde{B}$ J-NPILII of L if and only if it satisfies

$$M_A(\varsigma) \geq M_A((\varsigma \rightarrow (q \rightarrow \varsigma)'))'$$

$$\widetilde{B}_A(\varsigma) \geq \widetilde{B}_A((\varsigma \rightarrow (q \rightarrow \varsigma)'))'$$

$$J_A(\varsigma) \leq J_A((\varsigma \rightarrow (q \rightarrow \varsigma)'))'$$

Proof : Assume that A is a M $\widetilde{B}$ J-NPILII of L and let $\varsigma, q, \varphi \in L$. Let $\varphi = 0$, then

$$M_A(\varsigma) \geq \min\{M_A((\varsigma \rightarrow (q \rightarrow \varsigma)'))' \rightarrow 0)'), M_A(0)\}$$

$$= \min\{M_A(((\varsigma \rightarrow (q \rightarrow \varsigma)'))' \rightarrow 0)'), M_A(0)\}$$

$$= M_A((\varsigma \rightarrow (q \rightarrow \varsigma)'))'$$

$$\widetilde{B}_A(\varsigma) \geq \min\{\widetilde{B}_A((\varsigma \rightarrow (q \rightarrow \varsigma)'))' \rightarrow 0)'), \widetilde{B}_A(0)\}$$

$$= \min\{\widetilde{B}_A(((\varsigma \rightarrow (q \rightarrow \varsigma)'))' \rightarrow 0)'), \widetilde{B}_A(0)\}$$

$$= \widetilde{B}_A((\varsigma \rightarrow (q \rightarrow \varsigma)'))'$$

$$J_A(\varsigma) \leq \max\{J_A((\varsigma \rightarrow (q \rightarrow \varsigma)'))' \rightarrow 0)'), J_A(0)\}$$

$$= \max\{J_A(((\varsigma \rightarrow (q \rightarrow \varsigma)'))' \rightarrow 0)'), J_A(0)\}$$

$$= J_A((\varsigma \rightarrow (q \rightarrow \varsigma)'))'$$

Conversely suppose that

$$M_A(\varsigma) \geq M_A((\varsigma \rightarrow (q \rightarrow \varsigma)'))'$$

$$\widetilde{B}_A(\varsigma) \geq \widetilde{B}_A((\varsigma \rightarrow (q \rightarrow \varsigma)'))'$$

$$J_A(\varsigma) \leq J_A((\varsigma \rightarrow (q \rightarrow \varsigma)'))'$$

Since $M_A(\varsigma) \geq M_A((\varsigma \rightarrow (q \rightarrow \varsigma)'))'$

$$\begin{aligned}
&= \min\{M_A(((\zeta \rightarrow (q \rightarrow \zeta)')')'), M_A(0)\} \\
&= \min\{M_A(((\zeta \rightarrow (q \rightarrow \zeta)')' \rightarrow 0)'), M_A(0)\} \\
&\quad \widetilde{B}_A(\zeta) \geq \widetilde{B}_A((\zeta \rightarrow (q \rightarrow \zeta)')') \\
&= r \min\{\widetilde{B}_A(((\zeta \rightarrow (q \rightarrow \zeta)')')'), \widetilde{B}_A(0)\} \\
&= r \min\{\widetilde{B}_A(((\zeta \rightarrow (q \rightarrow \zeta)')' \rightarrow 0)'), \widetilde{B}_A(0)\} \\
J_A(\zeta) &\leq J_A(((\zeta \rightarrow (q \rightarrow \zeta)')')') \\
&= \max\{J_A(((\zeta \rightarrow (q \rightarrow \zeta)')')'), J_A(0)\} \\
&= \max\{J_A(((\zeta \rightarrow (q \rightarrow \zeta)')' \rightarrow 0)'), J_A(0)\}
\end{aligned}$$

Hence A is a MBJ-NPILII of L.

3.1.6. Theorem : Every MBJ-NPILII of a lattice implication algebra is a MBJ-NILII.

Proof : Let A be a MBJ-NPILII of a lattice implication algebra L.

$$M_A(\zeta) \geq \min\{M_A(((\zeta \rightarrow (q \rightarrow \zeta)')' \rightarrow \varphi)'), M_A(\varphi)\}$$

Put $\varphi = 0$ in the above equation, we get

$$\begin{aligned}
M_A(\zeta) &\geq \min\{M_A(((\zeta \rightarrow (q \rightarrow \zeta)')' \rightarrow 0)'), M_A(0)\} \\
&\quad = M_A(((\zeta \rightarrow (q \rightarrow \zeta)')' \rightarrow 0)') = M_A((\zeta \rightarrow (q \rightarrow \zeta)')')
\end{aligned}$$

$$\text{On the other hand } ((\zeta \rightarrow q)' \rightarrow ((\zeta \rightarrow (q \rightarrow q)')')')' = ((\zeta \rightarrow q)' \rightarrow q)'$$

$$\begin{aligned}
&\text{we obtain } M_A(\zeta \rightarrow q)' \geq M_A(((\zeta \rightarrow q)' \rightarrow \\
&\quad = M_A((\zeta \rightarrow q)' \rightarrow q)' \\
&\quad \geq \min\{M_A(((\zeta \rightarrow q)' \rightarrow q)' \rightarrow \varphi)'), M_A(\varphi)\}
\end{aligned}$$

$$\begin{aligned}
&\widetilde{B}_A(\zeta \rightarrow \varphi)' \geq \widetilde{B}_A(((\zeta \rightarrow \varphi)' \rightarrow \varphi)' \rightarrow (\zeta \rightarrow \varphi)') \\
&\quad = \widetilde{B}_A((\zeta \rightarrow q)' \rightarrow q)' \\
&\quad = r \min\{M_A(((\zeta \rightarrow q)' \rightarrow q)' \rightarrow \varphi)'), M_A(\varphi)\}
\end{aligned}$$

$$\begin{aligned}
J_A(\zeta \rightarrow \varphi)' &\leq J_A(((\zeta \rightarrow \varphi)' \rightarrow \varphi)' \rightarrow (\zeta \rightarrow \varphi)') \\
&\quad = J_A((\zeta \rightarrow q)' \rightarrow q)' \\
&\quad = \max\{M_A(((\zeta \rightarrow q)' \rightarrow q)' \rightarrow \varphi)'), M_A(\varphi)\}
\end{aligned}$$

Hence A is a MBJ-NILII.

3.2. MBJ-Neutrosophic associative LI-ideal :

3.2.1. Definition : Let ζ be a fixed element of a LIA 'L'. Let X be the universe. A MBJ-Neutrosophic set $A = \{(\zeta; M_A(\zeta), \widetilde{B}_A(\zeta), J_A(\zeta)/\zeta \in X)\}$ of a LIA L is called a MBJ-Neutrosophic associative LI-ideal of L with respect to ζ if it satisfies

$$\begin{aligned}
M_A(0) &\geq M_A(\zeta), \widetilde{B}_A(0) \geq \widetilde{B}_A(\zeta), I_A(0) \leq I_A(\zeta) \\
M_A(\varphi) &\geq \min\{M_A((\varphi \rightarrow \varrho)' \rightarrow \zeta)', M_A(\varrho \rightarrow \zeta)'\} \\
\widetilde{B}_A(\varphi) &\geq r \min\{\widetilde{B}_A((\varphi \rightarrow \varrho)' \rightarrow \zeta)', \widetilde{B}_A(\varrho \rightarrow \zeta)'\} \\
I_A(\varphi) &\leq \max\{I_A((\varphi \rightarrow \varrho)' \rightarrow \zeta)', I_A(\varrho \rightarrow \zeta)'\}
\end{aligned}$$

3.2.2. Proposition : Every MBJ-NALII of a LIA 'L' with respect to 0 is a MBJ-NLII.

Proof : Let A be a MBJ-NALII of a LIA 'L' with respect to 0. Then

$$\begin{aligned}
M_A(\varphi) &\geq \min\{M_A((\varphi \rightarrow \varrho)' \rightarrow 0)', M_A(\varrho \rightarrow 0)'\} = \min\{M_A(\varphi \rightarrow \varrho)', M_A(\varrho)\} \\
\widetilde{B}_A(\varphi) &\geq r \min\{\widetilde{B}_A((\varphi \rightarrow \varrho)' \rightarrow 0)', \widetilde{B}_A(\varrho \rightarrow 0)'\} = r \min\{\widetilde{B}_A(\varphi \rightarrow \varrho)', \widetilde{B}_A(\varrho)\} \\
I_A(\varphi) &\leq \max\{I_A((\varphi \rightarrow \varrho)' \rightarrow 0)', I_A(\varrho \rightarrow 0)'\} = \max\{I_A(\varphi \rightarrow \varrho)', I_A(\varrho)\}
\end{aligned}$$

Hence A is MBJ-LII.

3.2.3. Theorem : Let A be a MBJ-NLII of a LIA 'L'. Then A is a MBJ-NALII of a LIA 'L' if and only if it satisfies

$$\begin{aligned}
M_A(\varphi \rightarrow (\varrho \rightarrow \zeta)')' &\geq M_A((\varphi \rightarrow \varrho)' \rightarrow \zeta)') \\
\widetilde{B}_A(\varphi \rightarrow (\varrho \rightarrow \zeta)')' &\geq \widetilde{B}_A((\varphi \rightarrow \varrho)' \rightarrow \zeta)') \\
I_A(\varphi \rightarrow (\varrho \rightarrow \zeta)')' &\leq I_A((\varphi \rightarrow \varrho)' \rightarrow \zeta)')
\end{aligned}$$

Proof : Consider $M_A(\varphi \rightarrow (\varrho \rightarrow \zeta)')' \geq M_A((\varphi \rightarrow \varrho)' \rightarrow \zeta)')$

$$\begin{aligned}
\widetilde{B}_A(\varphi \rightarrow (\varrho \rightarrow \zeta)')' &\geq \widetilde{B}_A((\varphi \rightarrow \varrho)' \rightarrow \zeta)') \\
I_A(\varphi \rightarrow (\varrho \rightarrow \zeta)')' &\leq I_A((\varphi \rightarrow \varrho)' \rightarrow \zeta)')
\end{aligned}$$

Now we have to prove that A is a MBJ-NALII of a LIA 'L'.

Since A is a MBJ-NLII of a LIA 'L' then

$$\begin{aligned}
M_A(\varphi) &\geq \min\{M_A(\varphi \rightarrow (\varrho \rightarrow \zeta)')', M_A(\varrho \rightarrow \zeta)'\} \\
\widetilde{B}_A(\varphi) &\geq r \min\{\widetilde{B}_A(\varphi \rightarrow (\varrho \rightarrow \zeta)')', \widetilde{B}_A(\varrho \rightarrow \zeta)'\} \\
I_A(\varphi) &\leq \max\{I_A(\varphi \rightarrow (\varrho \rightarrow \zeta)')', I_A(\varrho \rightarrow \zeta)'\}
\end{aligned}$$

That implies $M_A(\varphi) \geq \min\{M_A((\varphi \rightarrow \varrho)' \rightarrow \zeta)'), M_A(\varrho \rightarrow \zeta)'\}$

$$\begin{aligned}
\widetilde{B}_A(\varphi) &\geq r \min\{\widetilde{B}_A((\varphi \rightarrow \varrho)' \rightarrow \zeta)'), \widetilde{B}_A(\varrho \rightarrow \zeta)'\} \\
I_A(\varphi) &\leq \max\{I_A((\varphi \rightarrow \varrho)' \rightarrow \zeta)'), I_A(\varrho \rightarrow \zeta)'\}
\end{aligned}$$

Hence A is a MBJ-NALII of a LIA 'L'.

Conversely consider A is a MBJ-NALII of a LIA 'L'.

Now we have to prove $M_A(\varphi \rightarrow (\varrho \rightarrow \zeta)')' \geq M_A((\varphi \rightarrow \varrho)' \rightarrow \zeta)')$

$$\begin{aligned}
\widetilde{B}_A(\varphi \rightarrow (\varrho \rightarrow \zeta)')' &\geq \widetilde{B}_A((\varphi \rightarrow \varrho)' \rightarrow \zeta)') \\
I_A(\varphi \rightarrow (\varrho \rightarrow \zeta)')' &\leq I_A((\varphi \rightarrow \varrho)' \rightarrow \zeta)')
\end{aligned}$$

Consider $((\varphi \rightarrow (\varrho \rightarrow \zeta)')' \rightarrow (\varphi \rightarrow \varrho)')' \rightarrow \zeta))$

$$\begin{aligned}
&= (\zeta' \rightarrow ((\varphi \rightarrow (\varrho \rightarrow \zeta)')' \rightarrow (\varphi \rightarrow \varrho)')) \\
&= \zeta' \rightarrow ((\varphi \rightarrow \varrho) \rightarrow (\varphi \rightarrow (\varrho \rightarrow \zeta)'))
\end{aligned}$$

$$\begin{aligned}
&= (\varphi \rightarrow q) \rightarrow (\zeta' \rightarrow ((q \rightarrow \zeta) \rightarrow \varphi')) \\
&= (\varphi \rightarrow q) \rightarrow ((q \rightarrow \zeta) \rightarrow (\zeta' \rightarrow \varphi')) \\
&= (\varphi \rightarrow q) \rightarrow (\varphi \rightarrow q) = 1
\end{aligned}$$

Then $M_A(\varphi \rightarrow (q \rightarrow \zeta)')'$

$$\begin{aligned}
&\geq \min \{M_A(((\varphi \rightarrow (q \rightarrow \zeta)')' \rightarrow (\varphi \rightarrow q)')' \rightarrow \zeta)), M_A((\varphi \rightarrow q)' \rightarrow \zeta')\} \\
&= \min \{M_A(0), M_A((\varphi \rightarrow q)' \rightarrow \zeta')\} \\
&= M_A((\varphi \rightarrow q)' \rightarrow \zeta')
\end{aligned}$$

Hence $M_A(\varphi \rightarrow (q \rightarrow \zeta)')' \geq M_A((\varphi \rightarrow q)' \rightarrow \zeta')$

$$\begin{aligned}
&\widetilde{B}_A(\varphi \rightarrow (q \rightarrow \zeta)')' \\
&\geq r \min \{\widetilde{B}_A(((\varphi \rightarrow (q \rightarrow \zeta)')' \rightarrow (\varphi \rightarrow q)')' \rightarrow \zeta)), \widetilde{B}_A((\varphi \rightarrow q)' \rightarrow \zeta')\} \\
&= r \min \{\widetilde{B}_A(0), \widetilde{B}_A((\varphi \rightarrow q)' \rightarrow \zeta')\} \\
&= \widetilde{B}_A((\varphi \rightarrow q)' \rightarrow \zeta')
\end{aligned}$$

Hence $\widetilde{B}_A(\varphi \rightarrow (q \rightarrow \zeta)')' \geq \widetilde{B}_A((\varphi \rightarrow q)' \rightarrow \zeta')$

$$\begin{aligned}
&I_A(\varphi \rightarrow (q \rightarrow \zeta)')' \\
&\leq \max \{I_A(((\varphi \rightarrow (q \rightarrow \zeta)')' \rightarrow (\varphi \rightarrow q)')' \rightarrow \zeta)), I_A((\varphi \rightarrow q)' \rightarrow \zeta')\} \\
&= \max \{I_A(0), I_A((\varphi \rightarrow q)' \rightarrow \zeta')\} \\
&= I_A((\varphi \rightarrow q)' \rightarrow \zeta')
\end{aligned}$$

Hence $I_A(\varphi \rightarrow (q \rightarrow \zeta)')' \leq I_A((\varphi \rightarrow q)' \rightarrow \zeta')$

3.2.4. Theorem : Let A be a MBJ-NLII of a LIA 'L'. Then A is a MBJ-NALII of a LIA 'L' if and only if it satisfies

$$\begin{aligned}
M_A(q) &\geq M_A((q \rightarrow \zeta)' \rightarrow \zeta') \\
\widetilde{B}_A(q) &\geq \widetilde{B}_A((q \rightarrow \zeta)' \rightarrow \zeta') \\
I_A(q) &\leq I_A((q \rightarrow \zeta)' \rightarrow \zeta')
\end{aligned}$$

Proof : Let A be a MBJ-NALII of a LIA 'L'. Then by above theorem we have

$$\begin{aligned}
M_A(\varphi \rightarrow (q \rightarrow \zeta)')' &\geq M_A((\varphi \rightarrow q)' \rightarrow \zeta') \\
\widetilde{B}_A(\varphi \rightarrow (q \rightarrow \zeta)')' &\geq \widetilde{B}_A((\varphi \rightarrow q)' \rightarrow \zeta') \\
I_A(\varphi \rightarrow (q \rightarrow \zeta)')' &\leq I_A((\varphi \rightarrow q)' \rightarrow \zeta')
\end{aligned}$$

Now take $\zeta = q$ then

$$\begin{aligned}
M_A(\varphi \rightarrow (q \rightarrow q)')' &\geq M_A((\varphi \rightarrow q)' \rightarrow q') \\
\widetilde{B}_A(\varphi \rightarrow (q \rightarrow q)')' &\geq \widetilde{B}_A((\varphi \rightarrow q)' \rightarrow q') \\
I_A(\varphi \rightarrow (q \rightarrow q)')' &\leq I_A((\varphi \rightarrow q)' \rightarrow q')
\end{aligned}$$

Since $q \rightarrow q = 1$ then

$$\begin{aligned}
M_A(\varphi \rightarrow (1)')' &\geq M_A((\varphi \rightarrow q)' \rightarrow q') \\
\widetilde{B}_A(\varphi \rightarrow (1)')' &\geq \widetilde{B}_A((\varphi \rightarrow q)' \rightarrow q')
\end{aligned}$$

$$I_A(\varphi \rightarrow (1)')' \leq I_A((\varphi \rightarrow q)' \rightarrow q)')$$

$$M_A(\varphi \rightarrow 0)' \geq M_A((\varphi \rightarrow q)' \rightarrow q)')$$

$$\widetilde{B}_A(\varphi \rightarrow 0)' \geq \widetilde{B}_A((\varphi \rightarrow q)' \rightarrow q)')$$

Since $\varphi \rightarrow 0 = \varphi'$ then

$$I_A(\varphi \rightarrow 0)' \leq I_A((\varphi \rightarrow q)' \rightarrow q)')$$

$$M_A(\varphi) \geq M_A((\varphi \rightarrow q)' \rightarrow q)')$$

$$\widetilde{B}_A(\varphi) \geq \widetilde{B}_A((\varphi \rightarrow q)' \rightarrow q)')$$

$$I_A(\varphi) \leq I_A((\varphi \rightarrow q)' \rightarrow q)')$$

Conversely suppose that

$$M_A(\varphi) \geq M_A((\varphi \rightarrow q)' \rightarrow q)')$$

$$\widetilde{B}_A(\varphi) \geq \widetilde{B}_A((\varphi \rightarrow q)' \rightarrow q)')$$

$$I_A(\varphi) \leq I_A((\varphi \rightarrow q)' \rightarrow q)')$$

Now we have to show that A is a MBJ-NALII of a LIA 'L'.

Consider $((\varphi \rightarrow \varsigma)' \rightarrow (q \rightarrow \varsigma)')' \rightarrow (\varphi \rightarrow q)' = 0$

And $((\varphi \rightarrow q)' \rightarrow (\varphi \rightarrow \varsigma)')' \leq (\varsigma \rightarrow q)'$

It follows that

$$(((\varphi \rightarrow (q \rightarrow \varsigma)')' \rightarrow \varsigma)' \rightarrow \varsigma)' \rightarrow ((\varphi \rightarrow q)' \rightarrow \varsigma)')' = 0$$

$$M_A((\varphi \rightarrow (q \rightarrow \varsigma)')')' \geq \{M_A(((\varphi \rightarrow (q \rightarrow \varsigma)')' \rightarrow \varsigma)' \rightarrow \varsigma)' \rightarrow ((\varphi \rightarrow q)' \rightarrow \varsigma)')', M_A((\varphi \rightarrow q)' \rightarrow \varsigma)')\}$$

$$\geq \min \{M_A(0), M_A((\varphi \rightarrow q)' \rightarrow \varsigma)')\} = M_A((\varphi \rightarrow q)' \rightarrow \varsigma)')$$

$$\widetilde{B}_A((\varphi \rightarrow (q \rightarrow \varsigma)')')' \geq \{\widetilde{B}_A(((\varphi \rightarrow (q \rightarrow \varsigma)')' \rightarrow \varsigma)' \rightarrow \varsigma)' \rightarrow ((\varphi \rightarrow q)' \rightarrow \varsigma)')', \widetilde{B}_A((\varphi \rightarrow q)' \rightarrow \varsigma)')\}$$

$$\geq r \min \{\widetilde{B}_A(0), \widetilde{B}_A((\varphi \rightarrow q)' \rightarrow \varsigma)')\} = \widetilde{B}_A((\varphi \rightarrow q)' \rightarrow \varsigma)')$$

$$I_A((\varphi \rightarrow (q \rightarrow \varsigma)')')' \leq \{I_A(((\varphi \rightarrow (q \rightarrow \varsigma)')' \rightarrow \varsigma)' \rightarrow \varsigma)' \rightarrow ((\varphi \rightarrow q)' \rightarrow \varsigma)')', I_A((\varphi \rightarrow q)' \rightarrow \varsigma)')\}$$

$$\leq \max \{I_A(0), I_A((\varphi \rightarrow q)' \rightarrow \varsigma)')\} = I_A((\varphi \rightarrow q)' \rightarrow \varsigma)')$$

Therefore A is a MBJ-NALII of a LIA 'L'.

3.3. MBJ-Neutrosophic Fantastic LI-ideal :

3.3.1. Definition : Let X be the universe. A MBJ-Neutrosophic set

$A = \{\langle \varsigma; M_A(\varsigma), \widetilde{B}_A(\varsigma), I_A(\varsigma) \rangle / \varsigma \in X\}$ of a LIA 'L' is called a MBJ-Neutrosophic fantastic LI-ideal of 'L' if it satisfies

$$M_A(0) \geq M_A(\varsigma), \widetilde{B}_A(0) \geq \widetilde{B}_A(\varsigma), I_A(0) \leq I_A(\varsigma) \text{ and}$$

$$M_A(\varsigma \rightarrow (q \rightarrow (q \rightarrow \varsigma)')')' \geq \min \{M_A((\varsigma \rightarrow q)' \rightarrow \varphi)', M_A(\varphi)\}$$

$$\widetilde{B}_A(\varsigma \rightarrow (q \rightarrow (q \rightarrow \varsigma)')')' \geq r \min \{\widetilde{B}_A((\varsigma \rightarrow q)' \rightarrow \varphi)', \widetilde{B}_A(\varphi)\}$$

$$I_A(\varsigma \rightarrow (q \rightarrow (q \rightarrow \varsigma)')')' \leq \max \{I_A((\varsigma \rightarrow q)' \rightarrow \varphi)', I_A(\varphi)\}$$

3.3.2. Example : The Hasse diagram and Cayley tables of a poset, $L = \{0, a, b, c, d, 1\}$ are given below as follows:




Figure 2: Hasse diagram of L

Table 7: Illustration of complement of element

a	b	ς	ς'
		0	1
	d	a	c
		b	d
		c	a
		d	b
		1	0

Table 8: Illustration of Implication

$\rightarrow$	0			a		
0	1	1	1	1	1	1
a	c	1	b	c	b	1
b	d	a	1	b	a	1
c	a	a	1	1	a	1
d	b	1	1	b	1	1
1	0	a	b	c	d	1

The operations $\vee$ and $\wedge$ on L are defined as follows

$$\varsigma \vee \varrho = (\varsigma \rightarrow \varrho) \rightarrow \varrho, \varsigma \wedge \varrho = ((\varsigma' \rightarrow \varrho') \rightarrow \varrho')', \text{ for all } \varsigma, \varrho \in L.$$

Clearly we can observe that L is a LIA.

Assume $A = (M_A, \widetilde{B}_A, J_A)$ is a MBJ-Neutrosophic set on L defined by Table given below

Table 9 : Illustration of Neutrosophic set $A = (M_A, \widetilde{B}_A, J_A)$

L	0	a	b	c	d	1
$M_A(\varsigma)$	0.8	0.4	0.4	0.6	0.4	0.4
$\widetilde{B}_A(\varsigma)$	[0.8, 0.9]	[0.3, 0.4]	[0.3, 0.4]	[0.3, 0.4]	[0.3, 0.4]	[0.2, 0.4]
$J_A(\varsigma)$	0.1	0.3	0.5	0.5	0.3	0.5

In general we verify that L is a MBJ-NFLII

3.3.3 Theorem : Every MBJ-NFLII is a MBJ-NLII.

Proof : Let A be a MBJ-NFLII of a LIA 'L'.

Taking $q = 0$ we get

$$M_A(\varsigma) = M_A(((\varsigma \rightarrow (0 \rightarrow (0 \rightarrow \varsigma)'))')' \geq \min\{M_A((\varsigma \rightarrow 0)' \rightarrow \varphi)'), M_A(\varphi)\}$$

$$= \min\{M_A(\varsigma \rightarrow \varphi)', M_A(\varphi)\}$$

$$\widetilde{B}_A(\varsigma) = \widetilde{B}_A(((\varsigma \rightarrow (0 \rightarrow (0 \rightarrow \varsigma)'))')' \geq r \min\{\widetilde{B}_A((\varsigma \rightarrow 0)' \rightarrow \varphi)'), \widetilde{B}_A(\varphi)\}$$

$$= r \min\{\widetilde{B}_A(\varsigma \rightarrow \varphi)', \widetilde{B}_A(\varphi)\}$$

$$I_A(\varsigma) = I_A(((\varsigma \rightarrow (0 \rightarrow (0 \rightarrow \varsigma)'))')' \leq \max\{I_A((\varsigma \rightarrow 0)' \rightarrow \varphi)'), I_A(\varphi)\}$$

$$= \max\{I_A(\varsigma \rightarrow \varphi)', I_A(\varphi)\}$$

Hence A is a MBJ-NLII.

Note : Converse of the above theorem is also true. It will be followed by the below example.

3.3.4. Example : Let $L = \{0, a, b, c, 1\}$. Define partial order on L as $0 < a < b < c < 1$ and define $\varsigma \wedge q = \min\{\varsigma, q\}$ and $\varsigma \vee q = \max\{\varsigma, q\}$ for all $\varsigma, q \in L$ and “'” and “ $\rightarrow$ ” as follows

Define MBJ-Neutrosophic set A in L by

Table 10 : Illustration of Neutrosophic set $A = (M_A, \widetilde{B}_A, I_A)$

L	0	a	b	c	1
M_A	0.5	0.4	0.4	0.4	0.2
$\widetilde{B}_A$	[0.4, 0.5]	[0.3, 0.4]	[0.3, 0.4]	[0.3, 0.4]	[0.2, 0.4]
I_A	0.2	0.4	0.4	0.4	0.5

Then A is a MBJ-NFLII on L, if A is a MBJ-NLII.

3.3.5. Theorem : A MBJ-NLII A of L is a MBJ-NFLII if and only if it satisfies

$$M_A(((\varsigma \rightarrow (q \rightarrow (q \rightarrow \varsigma)'))')' \geq M_A(\varsigma \rightarrow q)'$$

$$\widetilde{B}_A(((\varsigma \rightarrow (q \rightarrow (q \rightarrow \varsigma)'))')' \geq \widetilde{B}_A(\varsigma \rightarrow q)'$$

$$I_A(((\varsigma \rightarrow (q \rightarrow (q \rightarrow \varsigma)'))')' \geq I_A(\varsigma \rightarrow q)'$$

Proof : Let A be a MBJ-NFLII of a LIA 'L'.

Taking $q = 0$ in above definition

$$M_A(((\varsigma \rightarrow (q \rightarrow (q \rightarrow \varsigma)'))')' \geq \min\{M_A((\varsigma \rightarrow q)' \rightarrow 0)'), M_A(0)\} \geq M_A(\varsigma \rightarrow q)'$$

$$\widetilde{B}_A(((\varsigma \rightarrow (q \rightarrow (q \rightarrow \varsigma)'))')' \geq r \min\{\widetilde{B}_A((\varsigma \rightarrow q)' \rightarrow 0)'), \widetilde{B}_A(0)\} \geq \widetilde{B}_A(\varsigma \rightarrow q)'$$

$$I_A(((\varsigma \rightarrow (q \rightarrow (q \rightarrow \varsigma)'))')' \leq \max\{I_A((\varsigma \rightarrow q)' \rightarrow 0)'), I_A(0)\} \geq I_A(\varsigma \rightarrow q)'$$

Conversely suppose that A is a MBJ-NLII of 'L' and

$$\begin{aligned}
M_A(((\zeta \rightarrow (q \rightarrow (q \rightarrow \zeta)'))'))' &\geq M_A(\zeta \rightarrow q)' \geq \min \{ M_A((\zeta \rightarrow q)' \rightarrow \varphi)', M_A(\varphi) \} \\
\widetilde{B}_A(((\zeta \rightarrow (q \rightarrow (q \rightarrow \zeta)'))'))' &\geq \widetilde{B}_A(\zeta \rightarrow q)' \geq \text{rmin} \{ \widetilde{B}_A((\zeta \rightarrow q)' \rightarrow \varphi)', \widetilde{B}_A(\varphi) \} \\
I_A(((\zeta \rightarrow (q \rightarrow (q \rightarrow \zeta)'))'))' &\geq I_A(\zeta \rightarrow q)' \geq \max \{ I_A((\zeta \rightarrow q)' \rightarrow \varphi)', I_A(\varphi) \}
\end{aligned}$$

Hence A is a MBJ-NFLII of L.

3.3.6. Theorem : Every MBJ-NPILII of ‘L’ is a MBJ-NFLII.

Proof : Let A be a MBJ-NPILII of L.

Since every MBJ-NPILII of L is a MBJ-NLII

Since $\zeta \leq (\zeta \rightarrow (q \rightarrow (q \rightarrow \zeta)'))'$

we get $(q \rightarrow (\zeta \rightarrow (q \rightarrow (q \rightarrow \zeta)'))')' \leq (q \rightarrow \zeta)'$

Hence $(q \rightarrow (\zeta \rightarrow (q \rightarrow (q \rightarrow \zeta)'))')' \rightarrow (\zeta \rightarrow (q \rightarrow (q \rightarrow \zeta)'))'$

$$\begin{aligned}
&\geq (\zeta \rightarrow (q \rightarrow (q \rightarrow \zeta)'))' \rightarrow (q \rightarrow \zeta)' \\
&= (\zeta \rightarrow (q \rightarrow \zeta))' \rightarrow (q \rightarrow (q \rightarrow \zeta))'
\end{aligned}$$

$\geq (\zeta \rightarrow q)'$ and thus

$$\begin{aligned}
M_A(q \rightarrow \zeta)' &\geq M_A(q \rightarrow (\zeta \rightarrow (q \rightarrow (q \rightarrow \zeta)'))')' \rightarrow (\zeta \rightarrow (q \rightarrow (q \rightarrow \zeta)'))' \\
&\geq M_A(\zeta \rightarrow (q \rightarrow (q \rightarrow \zeta)'))' \\
\widetilde{B}_A(q \rightarrow \zeta)' &\geq \widetilde{B}_A(q \rightarrow (\zeta \rightarrow (q \rightarrow (q \rightarrow \zeta)'))')' \rightarrow (\zeta \rightarrow (q \rightarrow (q \rightarrow \zeta)'))' \\
&\geq \widetilde{B}_A(\zeta \rightarrow (q \rightarrow (q \rightarrow \zeta)'))' \\
I_A(q \rightarrow \zeta)' &\leq I_A(q \rightarrow (\zeta \rightarrow (q \rightarrow (q \rightarrow \zeta)'))')' \rightarrow (\zeta \rightarrow (q \rightarrow (q \rightarrow \zeta)'))' \\
&\leq I_A(\zeta \rightarrow (q \rightarrow (q \rightarrow \zeta)'))'
\end{aligned}$$

Therefore, by above theorem A is a MBJ-NFLII.

4. Conclusion:

In this paper we proved the relation between the ideals that “Every MBJ-NPILII is a MBJ-NLII”, “Every MBJ-NPILII is a MBJ-NILII”, “Every MBJ-NPILII is a MBJ-NFII”, “Every MBJ-NALII is a MBJ-NFII” and “Every MBJ-NFII is a MBJ-NLII”.

Funding: “This research received no external funding”.

Conflicts of Interest: “The authors declare no conflict of interest.”

References

- [1] Amarendra Babu. V, Abida Begum. K* and Siva Naga Malleswari. V on “MBJ-Neutrosophic LI-Ideals in Lattice Implication Algebras” published in Journal of Stochastic Modeling & Applications, ISSN: 0972-3641, Vol. 26 No. 3, page no.791-804, (January-June, Special Issues 2022 part-6).
- [2] Amarendra Babu. V, Abida Begum. K* and Siva Naga Malleswari. V on “MBJ-Neutrosophic Implicative LI-Ideals in Lattice Implication Algebras” published in Mathematical Statistician and Engineering Applications, ISSN: 2094-0343, 2326-9865, Vol.71, No. 3s2 (2022), page nos. 1022 - 1031.<http://philstat.org.ph>.
- [3] Borzooei R.A., Sabetkish M., and Jun Y.B., Neutrosophic LI-Ideal in lattice implication algebras, Neutrosophic Sets and Systems 31, 1 (2020). http://digitalrepository.unm.edu/nss_journal/vol/31/iss1/20.
- [4] Borzooei R.A., Zhang X.H., Smarandache F. and Jun Y.B., Commutative\generalized Neutrosophic ideals in BCK-algebras, Symmetry 10 (2018), 350 doi: 10.3390/sym10080350. Google Scholar.

- [5] Jun Y.B., Roh E.H. and Xu Y., LI-ideals in lattice implication algebras, Bull. Korean Math. Soc. 35(1) (1998), 13–24. Google Scholar.
- [6] Jun Y.B., Kim S.J. and Smarandache F., Interval Neutrosophic sets with Applications in BCK/BCI-algebra, Axioms 7 (2018), 23. doi: 10.3390/axioms7020023. Google Scholar.
- [7] Jun Y.B., Smarandache F. and Bordbar H., Neutrosophic N-structures applied to BCK/BCI-algebras, Information 8 (2017), 128. doi: 10.3390/info8040128. Google Scholar.
- [8] Jun Y.B., Smarandache F., Song S.Z. and Khan M., Neutrosophic positive Implicative N-Ideals in BCK/BCI-algebras, Axioms 7 (2018), 3. doi: 10.3390/axioms7010003. Google Scholar.
- [9] Jun Y.B. , E.H. Roh and Xu. Y, LI-Ideals in Lattice implication algebras, Bulletin of the Korean Mathematical Society, 35(1) (1998), 13–24.
- [10] Jun. Y.B, S. J. Kim and F. Smarandache, Interval Neutrosophic sets with applications in BCK / BCI algebra, Axioms 2018, 7, 23; doi:10.3390/axioms 7020023.
- [11] Mohseni Takallo. M, R. A. Borzooei, Young Bae Jun, MBJ-Neutrosophic structures and its applications in BCK/BCI-algebras, Neutrosophic sets and systems, Vol 23, 72-84, 2018.
- [12] Ozturk M.A. and Jun Y.B., Neutrosophic ideals in BCK/BCI-algebras based on Neutrosophic points, J. Inter. Math. Virtual Inst. 8 (2018), 1–17. Google Scholar.
- [13] Smarandache F., Neutrosophic set, a generalization of intuitionistic fuzzy sets, International Journal of Pure and Applied Mathematics 24(5) (2005), 287–297. Google Scholar.
- [14] Smarandache. F, A Unifying Field in Logics: Neutrosophic Logic, in Multiple Valued Logic / An International Journal, Vol. 8, No. 3, 385-438, 2002, www.gallup.unm.edu/~smarandache/eBook-neutrosophics4.pdf.
- [15] Smarandache. F, Definition of Neutrosophic Logic – A Generalization of the Intuitionistic Fuzzy Logic, Proceedings of the Third Conference of the European Society for Fuzzy Logic and Technology, EUSFLAT 2003, September 10-12, 2003, Zittau, Germany; University of Applied Sciences at Zittau/Goerlitz, 141-146.
- [16] Smarandache. F, Introduction to Neutrosophic Statistics, Sitech, 2014, <http://fs.gallup.unm.edu/NeutrosophicStatistics.pdf>, 2014.
- [17] Song S.Z., Smarandache. F., and Jun Y.B., Neutrosophic commutative N-Ideals in BCK-algebras, Information 8 (2017), 130. doi: 10.3390/info8040130. Google Scholar.
- [18] Xu Y., Lattice implication algebras, J. Southwest Jiaotong Univ. 1 (1993), 20–27. Google Scholar.