



On Some n-Refined Neutrosophic Groups For $3 \leq n \leq 5$

Abuobida M. Ahmed Alfahal^{1,*}, Sara Sawalmeh², Raja Abdullah Abdulfatah³, Yaser Ahmad Alhasan⁴

¹Deanship of the Preparatory Year, Prince Sattam bin Abdulaziz University, Alkharj, Saudi Arabia

²Mutah University, Faculty of Science, Mutah, Jordan

³Deanship the Preparatory Year, Prince Sattam bin Abdulaziz University, Alkharj, Saudi Arabia

⁴Deanship of the Preparatory Year, Prince Sattam bin Abdulaziz University, Alkharj, Saudi Arabia.

Emails: a.alfahal@psau.edu.sa; Sarawsawalmeh98@gmail.com; r.abdulfatah@psau.edu.sa;

y.alhasan@psau.edu.sa

*Corresponding author: a.alfahal@psau.edu.sa

Abstract

This paper is dedicated to studying some examples of n-refined neutrosophic groups and their algebraic substructures, where we deal with three different types of them, 3-refined, 4-refined, and 5-refined neutrosophic groups. Also, we present the algebraic structure of many substructures such as 3-refined neutrosophic AH-subgroups and kernels, 4-refined neutrosophic AH-homomorphisms and subgroups, and 5-refined neutrosophic AH-isomorphisms. On the other hand, many related examples will be provided to explain the algebraic concepts and their properties.

Keywords: 3-refined neutrosophic group; 4-refined neutrosophic group; 5-refined neutrosophic group.

1. Introduction

Neutrosophic algebra is the branch of neutrosophic applications that concerns the generalizations of famous classical algebraic structures into novel versions called neutrosophic algebraic structures [1-3,8-10].

In recent years, we have observed an accelerated development in the study of neutrosophic algebraic structures, since many of these structures have been studied and classified, for example, neutrosophic rings, neutrosophic matrices, neutrosophic vector spaces, and neutrosophic modules [4-7,11-13,19-22].

On the other hand, generalizations of neutrosophic algebraic structures have received a lot of attention from many researchers, as structures n-refined neutrosophic sets have been studied in many research papers. For example, n-refined neutrosophic rings, n-refined neutrosophic spaces, AH-substructures, and isometries [14-18,23-24].

n-refined neutrosophic groups have been studied in general in [25]. This has motivated us to study three different special cases, 3-refined neutrosophic groups, 4-refined neutrosophic groups, and 5-refined neutrosophic groups.

These structures will be studied in our work from many algebraic sides, especially those which are related to substructures such as AH-subgroups, AH-homomorphisms, and isomorphisms. Also, the classification of 3-refined, 4-refined, 5-refined neutrosophic groups in the case of additive abelian group will be proved and obtained.

2. Main results

Definition:

Let $(T,*)$ be a group, we define the 3-refined neutrosophic group $N_3(T)$ as follows:

Doi: <https://doi.org/10.54216/IJNS.220308>

Received: May 06, 2023 Revised: July 10, 2023 Accepted: October 13, 2023

$$N_3(T) = (\langle T \cup \{I_1, \dots, I_3\}, * \rangle) = \{(t_0, t_1 I_1, t_2 I_2, t_3 I_3); t_i \in T\}.$$

The set $N_3(T)$ is closed under $*$, and it is a semi-group but not a group since I_i is not invertible.

Remark:

If $(T, +)$ is an additive abelian group, then the addition operation on $N_3(T)$ is defined as follows:

Consider $x = (s_0, s_1 I_1, s_2 I_2, s_3 I_3), y = (r_0, r_1 I_1, r_2 I_2, r_3 I_3)$, we have

$$x + y = (s_0 + r_0, [s_1 + r_1]I_1, [s_2 + r_2]I_2, [s_3 + r_3]I_3). \text{ In this case } (N_3(T), +) \text{ is an abelian group.}$$

It can be classified as $N_3(T) \cong T \times T \times T \times T$.

Example:

Let $T = Z_2$ be the group of integers modulo 2, the corresponding 3-refined neutrosophic group is $N_3(T) = \{(0,0,0,0), (1,0,0,0), (0, I_1, 0,0), (0,0, I_2, 0), (1, I_1, 0,0), (1,0, I_2, 0), (0, I_1, I_2, 0), (1, I_1, I_2, 0), (1,0,0, I_1), (1,0,0, I_2), (1,0,0,1), (1, I_1, I_2, 1), (1, I_1, 0,1), (1, I_2, 0,1) \dots \dots \}$.

Remark:

If T is a multiplicative group, then group product on $N_3(T)$ can be defined as follows:

For $x = (s_0, s_1 I_1, s_2 I_2, s_3 I_3), y = (r_0, r_1 I_1, r_2 I_2, r_3 I_3)$, we have

$$xy = (l_0, l_1 I_1, l_2 I_2, l_3 I_3); t_s = \prod_{i,j=0}^3 (s_i r_j) I_i I_j; I_0 = e_G \text{ and } I_i I_j = I_{\min(i,j)}.$$

The identity element is $(e_G, e_G I_1, e_G I_2, e_G I_3)$.

In this case $N_3(T)$ is not isomorphic to the direct product of 4 copies of T , since it is not a group.

Example:

Let $T = S_3 = \{g_0 = I, g_1, g_2, g_3, g_4, g_5\}$ be symmetric group, the 3-refined neutrosophic group is $N_3(T) = \{(m, n I_1, s I_2, k I_3); m, n, s, k \in T\}$.

Definition:

(a) Let $N_3(T)$ be a 3-refined neutrosophic group. It is called abelian if $x * y = y * x$ for all $x, y \in N_3(T)$.

(b) The subset $Z(N_3(T)) = \{y \in N_3(T); y * x = x * y \text{ for all } x \in N_3(T)\}$ is called 3-refined neutrosophic center.

Theorem:

Let $N_3(T)$ be a 3-refined neutrosophic group. Then

(a) If G is abelian, $N_3(T)$ is abelian.

(b) $N_3(T)$ is abelian if and only if $N_3(T) = Z(N_3(T))$.

Definition:

Let $N_3(T)$ be a 3-refined neutrosophic group, K be a nonempty subset of $N_3(T)$, we say that K is a 3-refined neutrosophic subgroup if H contains a subgroup of T .

Definition:

Let $N_3(T)$ be a 3-refined neutrosophic group, the number of its elements is denoted by

$$O(N_3(T)).$$

If $N_3(T)$ is finite, then $O(N_3(T)) = k$, elsewhere $O(N_3(T)) = \infty$.

$O(N_3(T))$ is called the order of 3-refined neutrosophic group $N_3(T)$.

Theorem:

Let G be a finite group, $N_3(T)$ be its corresponding 3-refined neutrosophic group. Then if $O(G) = k$, we have $O(N_3(T)) = k^4$.

Definition:

Let $N_3(T), N_3(K)$ be two 3-refined neutrosophic groups, $h: N_3(T) \rightarrow N_3(K)$ be a well defined map, we say that h is a 3-refined neutrosophic homomorphism if:

- (a) $h(xy) = h(x)h(y)$ for all $x, y \in N_3(T)$.
 (b) $h(e_G, e_G, \dots, I_k, e_G, \dots, e_G) = (e_G, e_G, \dots, I_k, e_G, \dots, e_G)$.

Definition:

Let $N_3(T), N_3(K)$ be two 3-refined neutrosophic groups, $h: N_3(T) \rightarrow N_3(K)$ be a 3-refined neutrosophic homomorphism, we define:

- (a) $Ker(h) = \{x \in N_3(T); h(x) = e_{N_3(K)}\}$.
 (b) $Im(h) = \{y \in N_3(K); \exists x \in N_3(T): h(x) = y\}$.

Theorem:

Let $N_3(T), N_3(K)$ be two 3-refined neutrosophic groups, $h: N_3(T) \rightarrow N_3(K)$ be a 3-refined neutrosophic homomorphism, we have:

- (a) $Ker(h)$ is a 3-refined neutrosophic subgroup of $N_3(T)$.
 (b) $Im(h)$ is a 3-refined neutrosophic subgroup of $N_3(K)$.

Proof:

- (a) The restriction h_G of g is a homomorphism between

T and K , its kernel $Ker(h_G)$ will be a subset of $Ker(h)$, i.e $Ker(h)$ contains a subgroup of T , hence it is a 3-refined neutrosophic subgroup.

- (b) The proof is similar to (a).

Definition:

Let $N_3(T), N_3(K)$ be two 3-refined neutrosophic groups, $g: N_3(T) \rightarrow N_3(K)$ be a 3--refined neutrosophic homomorphism, we call it a 3--refined neutrosophic isomorphism if it is bijective.

Theorem:

Let $N_3(T)$ be a 3--refined neutrosophic group. The set $TI_k = \{x * I_k; 1 \leq k \leq 3\}$ has a group structure.

Proof:

We define the following operation on TI_k :

$(xI_k) \times (yI_k) = (x * y)I_k$. Also, \times is well defined operation, and it is associative with I_k as an identity, and for each

$xI_k \in TI_k$, there is an inverse $x^{-1}I_k$; x^{-1} is the inverse of x in T .

Theorem:

Let $N_3(T)$ be a 3--refined neutrosophic group, consider the previously defined group TI_k . Then $T \cong TI_k$.

Definition:

Let T, S be two groups, $T \times S$ be the direct product. We define the direct product of the related 3-refined neutrosophic groups as follows:

$N_3(T) \times N_3(S) = \langle T \times S \cup \{I_1, \dots, I_3\} \rangle$. We call it the 3-refined neutrosophic direct product.

Definition:

Let $N_3(T) = \{(t_0, t_1I_1, t_2I_2, t_3I_3); t_i \in G\}$ be a 3-refined neutrosophic group,

$N_3(S) = \{(m_0, m_1I_1, m_2I_2, m_3I_3); m_i \in S_i; S_i \text{ is a subgroup of } T \text{ for all } i\}$ is called an AH-subgroup of $N_3(T)$.

If $H_i \cong H_j$ for all $i \neq j$, then it is called an AHS-subgroup.

Definition:

Let $(T, *)$ be a group, we define the 4-refined neutrosophic group $N_4(T)$ as follows:

$$N_4(T) = (\langle T \cup \{I_1, \dots, I_4\}, * \rangle) = \{(t_0, t_1I_1, t_2I_2, t_3I_3, t_4I_4); t_i \in T\}.$$

The set $N_4(T)$ is closed under $*$, and it is a semi group but not a group since I_i is not invertible.

Remark:

If $(T, +)$ is an additive abelian group, then addition operation on $N_4(T)$ is defined as follows:

Consider $x = (s_0, s_1I_1, s_2I_2, s_3I_3, s_4I_4), y = (r_0, r_1I_1, r_2I_2, r_3I_3, r_4I_4)$, we have

$$x + y = (s_0 + r_0, [s_1 + r_1]I_1, [s_2 + r_2]I_2, [s_3 + r_3]I_3, [s_4 + r_4]I_4). \text{ In this case } (N_4(T), +) \text{ is an abelian group.}$$

It can be classified as $N_4(T) \cong T \times T \times T \times T \times T$.

Remark :

If T is a multiplicative group, then group product on $N_4(T)$ can be defined as follows:

For $x = (s_0, s_1I_1, s_2I_2, s_3I_3, s_4I_4), y = (r_0, r_1I_1, r_2I_2, r_3I_3, r_4I_4)$, we have

$$xy = (l_0, l_1I_1, l_2I_2, l_3I_3, l_4I_4); l_s = \prod_{i,j=0}^4 (s_i r_j) I_i I_j; I_0 = e_G \text{ and } I_i I_j = I_{\min(i,j)}.$$

The identity element is $(e_G, e_G I_1, e_G I_2, e_G I_3, e_G I_4)$.

In this case $N_4(T)$ is not isomorphic to the direct product of 5 copies of T , since it is not a group.

Definition:

(a) Let $N_4(T)$ be a 4-refined neutrosophic group. It is called abelian if $x * y = y * x$ for all $x, y \in N_4(T)$.

(b) The subset $Z(N_4(T)) = \{y \in N_4(T); y * x = x * y \text{ for all } x \in N_4(T)\}$ is called 4-refined neutrosophic center.

Theorem :

Let $N_4(T)$ be a 4-refined neutrosophic group. Then

(a) If G is abelian, $N_4(T)$ is abelian.

(b) $N_4(T)$ is abelian if and only if $N_4(T) = Z(N_4(T))$.

Definition:

Let $N_4(T)$ be an 4-refined neutrosophic group, K be a nonempty subset of $N_4(T)$, we say that K is a 4-refined neutrosophic subgroup if K contains a subgroup of T .

Definition:

Let $N_4(T)$ be a 4-refined neutrosophic group, the number of its elements is denoted by

$$O(N_4(T)).$$

If $N_4(T)$ is finite, then $O(N_4(T)) = k$, elsewhere $O(N_4(T)) = \infty$.

$O(N_4(T))$ is called the order of 4-refined neutrosophic group $N_4(T)$.

Theorem:

Let G be a finite group, $N_4(T)$ be its corresponding 4-refined neutrosophic group. Then if $O(G) = k$, we have $O(N_4(T)) = k^5$.

Definition:

Let $N_4(T), N_4(K)$ be two 4-refined neutrosophic groups, $h: N_4(T) \rightarrow N_4(K)$ be a well defined map, we say that h is a 4-refined neutrosophic homomorphism if:

- (a) $h(xy) = h(x)h(y)$ for all $x, y \in N_4(T)$.
 (b) $h(e_G, e_G, \dots, I_k, e_G, \dots, e_G) = (e_G, e_G, \dots, I_k, e_G, \dots, e_G)$.

Definition:

Let $N_4(T), N_4(K)$ be two 4-refined neutrosophic groups, $h: N_4(T) \rightarrow N_4(K)$ be a 4-refined neutrosophic homomorphism, we define:

- (a) $Ker(h) = \{x \in N_4(T); h(x) = e_{N_4(K)}\}$.
 (b) $Im(h) = \{y \in N_4(K); \exists x \in N_4(T): h(x) = y\}$.

Theorem:

Let $N_4(T), N_4(K)$ be two 4-refined neutrosophic groups, $h: N_4(T) \rightarrow N_4(K)$ be a 4-refined neutrosophic homomorphism, we have:

- (a) $Ker(h)$ is a 4-refined neutrosophic subgroup of $N_4(T)$.
 (b) $Im(h)$ is a 4-refined neutrosophic subgroup of $N_4(K)$.

Proof:

- (a) The restriction h_G of g is a homomorphism between

T and K , its kernel $Ker(h_G)$ will be a subset of $Ker(h)$, i.e $Ker(h)$ contains a subgroup of T , hence it is a 3-refined neutrosophic subgroup.

- (b) The proof is similar to (a).

Definition :

Let $N_4(T), N_4(K)$ be two 4-refined neutrosophic groups, $g: N_4(T) \rightarrow N_4(K)$ be a 4--refined neutrosophic homomorphism, we call it a 4--refined neutrosophic isomorphism if it is bijective.

Theorem:

Let $N_4(T)$ be a 4--refined neutrosophic group. The set $TI_k = \{x * I_k; 1 \leq k \leq 4\}$ has a group structure.

Proof:

We define the following operation on TI_k :

$(xI_k) \times (yI_k) = (x * y)I_k$. Also, \times is well defined operation, and it is associative with I_k as an identity, and for each

$xI_k \in TI_k$, there is an inverse $x^{-1}I_k$; x^{-1} is the inverse of x in T .

Theorem:

Let $N_4(T)$ be a 4--refined neutrosophic group, consider the previously defined group TI_k . Then $T \cong TI_k$.

Definition:

Let T, S be two groups, $T \times S$ be the direct product. We define the direct product of the related 4-refined neutrosophic groups as follows:

$N_4(T) \times N_4(S) = \langle T \times S \cup \{I_1, \dots, I_4\} \rangle$. We call it the 4-refined neutrosophic direct product.

Definition:

Let $N_4(T) = \{(t_0, t_1I_1, t_2I_2, t_3I_3, t_4I_4); t_i \in T\}$ be a 4-refined neutrosophic group,

$N_4(S) = \{(m_0, m_1I_1, m_2I_2, m_3I_3, m_4I_4); m_i \in S_i; S_i \text{ is a subgroup of } T \text{ for all } i\}$ is called an AH-subgroup of $N_4(T)$.

If $H_i \cong H_j$ for all $i \neq j$, then it is called an AHS-subgroup.

Definition:

Let $(T, *)$ be a group, we define the 5-refined neutrosophic group $N_5(T)$ as follows:

$$N_5(T) = (\langle T \cup \{I_1, \dots, I_5\}, * \rangle) = \{(t_0, t_1I_1, t_2I_2, t_3I_3, t_4I_4, t_5I_5); t_i \in T\}.$$

The set $N_5(T)$ is closed under $*$, and it is a semi-group but not a group since I_i is not invertible.

Remark:

If $(T, +)$ is an additive abelian group, then the addition operation on $N_5(T)$ is defined as follows:

Consider $x = (s_0, s_1I_1, s_2I_2, s_3I_3, s_4I_4, s_5I_5), y = (r_0, r_1I_1, r_2I_2, r_3I_3, r_4I_4, r_5I_5)$, we have

$x + y = (s_0 + r_0, [s_1 + r_1]I_1, [s_2 + r_2]I_2, [s_3 + r_3]I_3, [s_4 + r_4]I_4, [s_5 + r_5]I_5)$. In this case $(N_5(T), +)$ is an abelian group.

It can be classified as $N_5(T) \cong T \times T \times T \times T \times T \times T$.

Remark :

If T is a multiplicative group, then the group product on $N_5(T)$ can be defined as follows:

For $x = (s_0, s_1I_1, s_2I_2, s_3I_3, s_4I_4, s_5I_5), y = (r_0, r_1I_1, r_2I_2, r_3I_3, r_4I_4, r_5I_5)$, we have

$$xy = (l_0, l_1I_1, l_2I_2, l_3I_3, l_4I_4, l_5I_5); l_s = \prod_{i,j=0}^5 (s_i r_j) I_i I_j; l_0 = e_G \text{ and } I_i I_j = I_{\min(i,j)}.$$

The identity element is $(e_G, e_G I_1, e_G I_2, e_G I_3, e_G I_4, e_G I_5)$.

In this case $N_5(T)$ is not isomorphic to the direct product of 6 copies of T , since it is not a group.

Definition:

(a) Let $N_5(T)$ be a 4-refined neutrosophic group. It is called abelian if $x * y = y * x$ for all $x, y \in N_5(T)$.

(b) The subset $Z(N_5(T)) = \{y \in N_5(T); y * x = x * y \text{ for all } x \in N_5(T)\}$ is called 5-refined neutrosophic center.

Theorem :

Let $N_4(T)$ be a 5-refined neutrosophic group. Then

(a) If G is abelian, $N_5(T)$ is abelian.

(b) $N_5(T)$ is abelian if and only if $N_5(T) = Z(N_5(T))$.

Definition:

Let $N_5(T)$ be an 5-refined neutrosophic group, K be a nonempty subset of $N_5(T)$, we say that K is a 5-refined neutrosophic subgroup if K contains a subgroup of T .

Definition:

Let $N_5(T)$ be a 5-refined neutrosophic group, the number of its elements is denoted by

$$O(N_5(T)).$$

If $N_5(T)$ is finite, then $O(N_5(T)) = k$, elsewhere $O(N_5(T)) = \infty$.

$O(N_5(T))$ is called the order of 5-refined neutrosophic group $N_5(T)$.

Theorem:

Let G be a finite group, $N_5(T)$ be its corresponding 5-refined neutrosophic group. Then if $O(G) = k$, we have $O(N_5(T)) = k^6$.

Definition:

Let $N_5(T), N_5(K)$ be two 5-refined neutrosophic groups, $h: N_5(T) \rightarrow N_5(K)$ be a well defined map, we say that h is a 5-refined neutrosophic homomorphism if:

- (a) $h(xy) = h(x)h(y)$ for all $x, y \in N_5(T)$.
 (b) $h(e_G, e_G, \dots, I_k, e_G, \dots, e_G) = (e_G, e_G, \dots, I_k, e_G, \dots, e_G)$.

Definition:

Let $N_5(T), N_5(K)$ be two 5-refined neutrosophic groups, $h: N_5(T) \rightarrow N_5(K)$ be a 5-refined neutrosophic homomorphism, we define:

- (a) $Ker(h) = \{x \in N_5(T); h(x) = e_{N_5(K)}\}$.
 (b) $Im(h) = \{y \in N_5(K); \exists x \in N_5(T): h(x) = y\}$.

Theorem:

Let $N_5(T), N_5(K)$ be two 5-refined neutrosophic groups, $h: N_5(T) \rightarrow N_5(K)$ be a 5-refined neutrosophic homomorphism, we have:

- (a) $Ker(h)$ is a 5-refined neutrosophic subgroup of $N_5(T)$.
 (b) $Im(h)$ is a 5-refined neutrosophic subgroup of $N_5(K)$.

Proof:

- (a) The restriction h_G of g is a homomorphism between

T and K , its kernel $Ker(h_G)$ will be a subset of $Ker(h)$, i.e $Ker(h)$ contains a subgroup of T , hence it is a 3-refined neutrosophic subgroup.

- (b) The proof is similar to (a).

Definition :

Let $N_4(T), N_4(K)$ be two 4-refined neutrosophic groups, $g: N_4(T) \rightarrow N_4(K)$ be a 4-refined neutrosophic homomorphism, we call it a 4--refined neutrosophic isomorphism if it is bijective.

Theorem:

Let $N_5(T)$ be a 5--refined neutrosophic group. The set $TI_k = \{x * I_k; 1 \leq k \leq 5\}$ has a group structure.

Proof:

We define the following operation on TI_k :

$(xI_k) \times (yI_k) = (x * y)I_k$. Also, \times is well defined operation, and it is associative with I_k as an identity, and for each

$xI_k \in TI_k$, there is an inverse $x^{-1}I_k$; x^{-1} is the inverse of x in T .

Theorem:

Let $N_5(T)$ be a 5--refined neutrosophic group, consider the previously defined group TI_k . Then $T \cong TI_k$.

Definition:

Let T, S be two groups, $T \times S$ be the direct product. We define the direct product of the related 5-refined neutrosophic groups as follows:

$N_5(T) \times N_5(S) = \langle T \times S \cup \{I_1, \dots, I_5\} \rangle$. We call it the 5-refined neutrosophic direct product.

Definition:

Let $N_5(T) = \{(t_0, t_1I_1, t_2I_2, t_3I_3, t_4I_4, t_5I_5); t_i \in T\}$ be a 5-refined neutrosophic group,

$N_5(S) = \{(m_0, m_1I_1, m_2I_2, m_3I_3, m_4I_4, m_5I_5); m_i \in S_i; S_i \text{ is a subgroup of } T \text{ for all } i\}$ is called an AH-subgroup of $N_5(T)$.

If $H_i \cong H_j$ for all $i \neq j$, then it is called an AHS-subgroup.

3. Conclusion

In this paper, we have defined the concept of 3,4,5-refined neutrosophic groups. Also, we have introduced some of their substructures such as 3-refined neutrosophic subgroup/homomorphism, 4-refined neutrosophic homomorphism/subgroup, and 5-refined neutrosophic isomorphism/homomorphism. Many examples were produced to explain the abstract algebraic concepts and theorems.

Acknowledgments " This study is supported via funding from Prince sattam bin Abdulaziz University project number (PSAU/2023/R/1445)".

References

- [1] Agboola, A.A.A., Akwu, A.D., and Oyebo, Y.T., "Neutrosophic Groups and Subgroups", International J .Math.Combin, Vol. 3, pp. 1-9, 2012.
- [2] Agboola, A.A.A., Akinola, A.D., and Oyebola, O.Y., " NeutrosophicRings I ", International J.Mathcombin, Vol. 4, pp. 1-14, 2011.
- [3] Agboola, A.A.A., "On Refined Neutrosophic Algebraic Structures", Neutrosophic Sets and Systems, Vol. 10, pp. 99-102, 2015.
- [4] Abobala, M., On Refined Neutrosophic Matrices and Their Applications In Refined Neutrosophic Algebraic Equations, Journal Of Mathematics, Hindawi, 2021
- [5] Merkepci, H., and Ahmad, K., " On The Conditions Of Imperfect Neutrosophic Duplets and Imperfect Neutrosophic Triplets", Galoistica Journal Of Mathematical Structures And Applications, Vol.2, 2022.
- [6] Abobala, M., Hatip, A., Olgun, N., Broumi, S., Salama, A.A., and Khaled, E, H., The algebraic creativity In The Neutrosophic Square Matrices, Neutrosophic Sets and Systems, Vol. 40, pp. 1-11, 2021.
- [7] Hatip, A., and Abobala, M., "AH-Substructures In Strong Refined Neutrosophic Modules", International Journal of Neutrosophic Science, Vol. 9, pp. 110-116 . 2020.
- [8] Kandasamy, V.W.B., and Smarandache, F., "Some Neutrosophic Algebraic Structures and Neutrosophic N-Algebraic Structures", Hexis, Phonex, Arizona 2006.
- [9] Smarandache, F., "Symbolic Neutrosophic Theory", EuropaNova asbl, Bruxelles, 2015.
- [10] Smarandache, F., "*n*-Valued Refined Neutrosophic Logic and Its Applications in Physics", Progress in Physics, Vol. 4, pp.143-146, 2013.
- [11] Abobala, M., and Hatip, A., " An Algebraic Approach To Neutrosophic Euclidean Geometry", Neutrosophic Sets and Systems, 2021.
- [12] Abobala, M., "On The Characterization of Maximal and Minimal Ideals In Several Neutrosophic Rings", Neutrosophic Sets and Systems, Vol. 45, 2021.
- [13] Abobala, M., "Neutrosophic Real Inner Product Spaces", Neutrosophic Sets and Systems, Vol. 43, 2021
- [14] Sarkis, M., " On The Solutions Of Fermat's Diophantine Equation In 3-refined Neutrosophic Ring of Integers", Neoma Journal of Mathematics and Computer Science, 2023.
- [15] Smarandache, F., and Abobala, M., "n-Refined Neutrosophic Rings", International Journal Of Neutrosophic Science, 2020.

- [16] Ahmad, K., Thjeel, N., AbdulZahra, M., and Jaleel, R., " On The Classification of n -Refined Neutrosophic Rings And Its Applications In Matrix Computing Algorithms And Linear Systems", *International Journal Of Neutrosophic Science*, 2022.
- [17] Celik, M., and Hatip, A., " On The Refined AH-Isometry And Its Applications In Refined Neutrosophic Surfaces", *Galoitica Journal Of Mathematical Structures And Applications*, 2022.
- [18] Abobala, M., "On Some Algebraic Properties of n -Refined Neutrosophic Elements and n -Refined Neutrosophic Linear Equations", *Mathematical Problems in Engineering*, Hindawi, 2021.
- [19] Adeleke, E.O., Agboola, A.A.A., and Smarandache, F., "Refined Neutrosophic Rings II", *International Journal of Neutrosophic Science*, Vol. 2(2), pp. 89-94. 2020.
- [20] Ibrahim, M.A., Agboola, A.A.A, Badmus, B.S. and Akinleye, S.A., "On refined Neutrosophic Vector Spaces I", *International Journal of Neutrosophic Science*, Vol. 7, pp. 97-109. 2020.
- [21] Al Aswad, M., and Dalla, R., " Neutrosophic Divisor Point of A Straight Line Segment With A Given Ratio", *Pure Mathematics for Theoretical Computer Science*, Vol.2, 2023.
- [22] Smarandache F., and Abobala, M., " n -Refined Neutrosophic Vector Spaces", *International Journal of Neutrosophic Science*, Vol. 7, pp. 47-54. 2020.
- [23] Sankari, H., and Abobala, M." n -Refined Neutrosophic Modules", *Neutrosophic Sets and Systems*, Vol. 36, pp. 1-11. 2020.
- [24] Abobala, M., "A Study of AH-Substructures in n -Refined Neutrosophic Vector Spaces", *International Journal of Neutrosophic Science*", Vol. 9, pp.74-85. 2020.
- [25] Abobala, M., " n -Refined Neutrosophic Groups I", *International Journal of Neutrosophic Science*, Vol. 0, pp. 27-34. 2020.