



Numerical Procedures for Computing the Exact Solutions to Systems of Ordinary Differential Equations

Nidal Anakira^{1,2,*}, Osama Oqilat³, Adel Almalki⁴, Irianto Irianto⁵, Saad Meqdad⁶, Ala Amourah¹

¹Faculty of Education and Arts, Mathematics Section, Sohar University, Sohar 3111, Sultanat of Oman

²Jadara University Research Center, Jadara University, Jordan

³Department of Basic Sciences, Faculty of Arts and Science, Al-Ahliyya Amman University, Amman 19328, Jordan

⁴Department of Mathematics, Al-Qunfudhah University College, Umm Al-Qura University, Mecca, Saudi Arabia

⁵Department General Education, Faculty of Resilience, Rabdan Academy, Abu Dhabi 22401, United Arab Emirates

⁶Applied Science Private University, Amman, Jordan

Email: nanakira@su.edu.om; o.oqilat@ammanu.edu.jo; aaamalki@uqu.edu.sa; iharny@ra.ac.ae; s_meq75@yahoo.com; AAmourah@su.edu.om

Abstract

This paper presents a modified homotopy perturbation method (HPM), which aimed at solving systems of ordinary differential equations (ODEs). The MHPM, which combines the HPM, Laplace transform, and Padé approximants, offers an alternative approach to address the challenges associated with solving such problems. By employing this method, it becomes feasible to overcome these challenges and obtain a dependable approximation for the exact solution. The effectiveness and applicability of the proposed scheme are demonstrated through preliminary results derived from illustrative examples, all of which correspond to exact solutions.

Keywords: Numerical Approximation; HPM; MHPM; Laplace transformation; Padé approximants

1 Introduction

Solving systems of ODEs is a critical task in various fields of applied science and engineering. Whether one is modelling physical phenomena or optimizing industrial processes, accurately solving these equations is essential for understanding the behaviour of dynamic systems. However, finding analytical solutions can be challenging and even impractical, especially for complex systems with nonlinearities. In such cases, numerical methods offer a powerful range of tools for approximating the solution. There are different numerical schemes in fields of numerical analysis that are used in finding approximate solutions for various types of equations [1-20, 28-30]. The solutions resulting from these schemes are excellent and agree with the exact solutions in all cases. However, the difference between them lies only in the structure of the method, which reflects the ease of calculations or requires more effort and time.

In this study, we present a novel enhancement to the utilization of the HPM for resolving systems of ODEs. Our approach can be applied to any given problem, providing exact solutions, opposite the HPM solution that converge towards the exact solution as the number of approximation terms increases. It is noteworthy to acknowledge that the accuracy of our approach does not depend about the approximation employed, which may require additional computational resources and time, especially when dealing with nonlinear problems. Consequently, researchers are continuously striving to develop or adapt numerical techniques to achieve higher accuracy or exact solutions. The primary objective of this paper is to improve the accuracy of the HPM through the application of an alternative methodology. This methodology entails modifying the series solution of the HPM by utilizing the Laplace

transform on the truncated HPM solution. Subsequently, the transformed series is converted into a meromorphic function using Padé approximants. Lastly, the inverse Laplace transform is employed to derive the desired solution for the given problem. This approach is simple, requiring minimal effort and is highly efficient in obtaining accurate results.

This work organized as follows: Section 2 introduces the fundamental concept of the HPM, along with a brief explanation of the Pade approximants. In Section 3, numerical examples are presented to demonstrate the effectiveness of the discussed procedure in obtaining the exact solution for systems of ODEs. The results highlight that accurate solutions can be obtained with only a few terms. The final section summarizes the conclusions of this work.

2. Fundamental Idea of HPM Procedure

To demonstrate the fundamental concept of the HPM procedure [21-25], let's examine the following equation.

$$A(u) - f(r) = 0, \quad r \in \Omega, \quad (1)$$

where A is the integral operator consist of the linear and nonlinear operator L and N , respectively, while B is a boundary operator, $f(r)$ is a known function, and Γ is the boundary of the domain Ω . Eq. (1) can be rewritten as

$$L(u) - N(u) - f(r) = 0. \quad (2)$$

A homotopy equation $v : \Omega[0, 1] \rightarrow \mathbb{R}$ which satisfies

$$H(v; p) = L(v) - L(v_0) + pL(v_0) + p[N(v) - f(r)] = 0, \quad (3)$$

or

$$H(v; p) = (1 - p)[L(v) - L(v_0)] + p[A(v_0) - F(r)] = 0, \quad (4)$$

is constructed, here, $r \in \Omega$, $p \in [0, 1]$ is the homotopy parameter, and $v_0(x)$ is an initial approximation of Eq. (1). It is noted that

$$(v; 0) = L(u) - L(v_0) = 0, \quad H(v; 1) = A(v) - F(r) = 0. \quad (5)$$

The changing process of p from 0 to 1, include the changing of $H(v; p)$ from $L(u) - L(v_0)$ to $A(v) - F(r)$ that is called deformation. Moreover, $L(u) - L(v_0)$ and $A(v) - F(r)$ are called homotopic. Note that, $0 \leq p \leq 1$ it is considered as a small parameter, the solution of Eqs. (3) or (4) can be expressed as a series in p , as follows:

$$v = v_0 + pv_1 + p^2v_2 + p^3v_3 + \dots \quad (6)$$

when $p \rightarrow 1$, the approximate solution of Eq. (1). i.e.,

$$u(x) = \lim_{p \rightarrow 1} v(x) = v_1 + v_2 + v_3 + \dots \quad (7)$$

3. Padé approximation

For the function $u(x)$ [26, 27], the Padé approximation of order $\left[\frac{L}{M}\right]$, can be formulated as follows,

$$\left[\frac{L}{M}\right] = \frac{P_L(x)}{Q_M(x)},$$

where $P_L(x)$ and $Q_M(x)$, are two polynomials of the highest degree L and M . The power series is given in form of

$$u(x) = \sum_{i=1}^{\infty} a_i x^i.$$

The coefficients of the polynomials $P_L(x)$ and $Q_M(x)$, can be obtained from

$$u(t) - \frac{P_L(x)}{Q_M(x)} = O(x^{L+M+1}). \quad (8)$$

When the denominator and numerator's functions $\frac{P_L(x)}{Q_M(x)}$ is multiplied by a constant that is not zero, the fractional values stay the same, such that we can set up the normalization requirement as

$$Q_M(0) = 1. \quad (9)$$

It observed that the polynomial for the functions $P_L(x)$ and $Q_M(x)$ has no public factors. The coefficients of the polynomial $Q_M(x)$ and $P_L(x)$ are given by

$$\begin{aligned} P_L(t) &= P_0 + P_1 t + P_2 t^2 + \cdots + P_L t^L, \\ Q_M(t) &= q_0 + q_1 t + q_2 t^2 + \cdots + q_M t^M, \end{aligned} \quad (10)$$

the following linear systems of coefficients can be obtained by multiplying Eq. (8) by $Q_M(x)$ in light of Eq. (8).

$$\begin{cases} a_{L+1} + a_L q_1 + \cdots + a_{L-M+1} q_M = 0 \\ a_{L+2} + a_{L+1} q_1 + \cdots + a_{L-M+2} q_M = 0 \\ \vdots \\ a_{L+M} + a_{L+M-1} q_1 + \cdots + a_L q_M = 0 \end{cases}, \quad (11)$$

$$\begin{cases} a_0 = P_0 \\ a_0 + a_0 q_1 = P_1 \\ a_2 + a_1 q_1 + a_0 q_2 = P_2 \\ \vdots \\ a_L + a_{L-1} q_1 + \cdots + a_0 q_L = P_L \end{cases}, \quad (12)$$

These equations will be solved using Eq. (11). It is seen as a set of linear formulas for the unidentified q 's. When the q 's are identified, then (12) have an explicit formula for the unknown p 's, this concludes the solution to the problem. If (11) and (12) are non-singular, then we can solve them directly and get Eq. (13).

$$\begin{bmatrix} L \\ M \end{bmatrix} = \frac{\det \begin{bmatrix} a_{L-M+1} & a_{L-M+2} & \cdots & a_{L+1} \\ \vdots & \vdots & \ddots & \vdots \\ a_L & a_{L+1} & \cdots & a_{L+M} \\ \sum_{j=M}^L a_{j-M} X^j & \sum_{j=M-1}^L a_{j-M+1} X^j & \cdots & \sum_{j=0}^L a_j X^j \end{bmatrix}}{\det \begin{bmatrix} a_{L-M+1} & a_{L-M+2} & \cdots & a_{L+1} \\ \vdots & \vdots & \ddots & \vdots \\ a_L & a_{L+1} & \cdots & a_{L+M} \\ X^M & X^{M-1} & \cdots & 1 \end{bmatrix}}, \quad (13)$$

4. Applications of HPM

In the following section, we present two illustrative examples of systems of ordinary differential equations (ODEs), both linear and nonlinear. The purpose of displaying these examples is to demonstrate the effectiveness and reliability of the Modified HPM procedure.

Example 4. 1 Consider the following system of first-order linear equations [23].

$$\begin{aligned} u_1'(t) &= u_2(t), \\ u_2'(t) &= u_3(t), \\ u_3'(t) &= \frac{1}{t} u_1(t) + u_3(t), \end{aligned} \quad (14)$$

Subject to

$$u_1(0) = 0, u_2(0) = 1, \text{ and } u_3(0) = 2, \text{ with exact solutions } u = (u_1(t), u_2(t)) = (te^t, e^t(1+t), e^t(2+t)),$$

We will now formulate the following set of homotopy equations, following the algorithm outlined in Section 2.

$$\begin{aligned}(1-q) \frac{dv_1(t;p)}{dt} &= (h;q) \left(\frac{dv_1(t;p)}{dt} - v_2(t;p) \right) \\ (1-q) \frac{dv_2(t;p)}{dt} &= (h;q) \left[\frac{dv_1(t;p)}{dt} - v_3(t;p) \right], \\ (1-p) \left[\frac{dv_3(t;p)}{dt} \right] &= (h;q) \left[\frac{dv_3(t;p)}{dt} - \frac{1}{t} v_1(t;p) - v_3(t;p) \right]\end{aligned}\quad (15)$$

Following the same process in example one, we have the 5th -order HPM approximate solution

$$\begin{aligned}\tilde{u}_1(t) &= t + t^2 + \frac{t^3}{2} + \frac{t^4}{6} + \frac{t^5}{24} + \frac{t^6}{120} + \frac{167t^7}{151200} + \frac{t^8}{20160}, \\ \tilde{u}_2(t) &= 1 + 2t + \frac{3t^2}{2} + \frac{2t^3}{3} + \frac{5t^4}{24} + \frac{t^5}{20} + \frac{13t^6}{1350} + \frac{59t^7}{50400} + \frac{t^8}{20160}, \\ \tilde{u}_3(t) &= 2 + 3t + 2t^2 + \frac{5t^3}{6} + \frac{t^4}{4} + \frac{7t^5}{120} + \frac{1403t^6}{129600} + \frac{433t^7}{352800} + \frac{t^8}{20160}.\end{aligned}\quad (16)$$

Tables (1), (2) and (2) present a comparison between HPM process and exact solutions. We observed that the accuracy of the results varied depending on the order of the approximations. Therefore, to enhance the precision of the HPM procedure, we will begin by applying the Laplace transformation to the initial terms in the HPM series solutions. Next, we will utilize the Pade approximants and finally, we will conclude by implementing the inverse Laplace transformation. The process is outlined below

$$\begin{aligned}L(\tilde{u}_1(t)) &= \frac{2}{s^9} + \frac{167}{30s^8} + \frac{6}{s^7} + \frac{5}{s^6} + \frac{4}{s^5} + \frac{3}{s^4} + \frac{2}{s^3} + \frac{1}{s^2}, \\ L(\tilde{u}_2(t)) &= \frac{2}{s^9} + \frac{59}{10s^8} + \frac{104}{15s^7} + \frac{6}{s^6} + \frac{5}{s^5} + \frac{4}{s^4} + \frac{3}{s^3} + \frac{2}{s^2} + \frac{1}{s}, \\ L(\tilde{u}_3(t)) &= \frac{2}{s^9} + \frac{433}{70s^8} + \frac{1403}{180s^7} + \frac{7}{s^6} + \frac{6}{s^5} + \frac{5}{s^4} + \frac{4}{s^3} + \frac{3}{s^2} + \frac{2}{s},\end{aligned}\quad (17)$$

Use $s = \frac{1}{z}$, leads to

$$\begin{aligned}L(\tilde{u}_1(t)) &= z^2 + 2z^3 + 3z^4 + 4z^5 + 5z^6 + 6z^7 + \frac{167z^8}{30} + 2z^9, \\ L(\tilde{u}_2(t)) &= z + 2z^2 + 3z^3 + 4z^4 + 5z^5 + 6z^6 + \frac{104z^7}{15} + \frac{59z^8}{10} + 2z^9, \\ L(\tilde{u}_3(t)) &= 2z + 3z^2 + 4z^3 + 5z^4 + 6z^5 + 7z^6 + \frac{1403z^7}{180} + \frac{433z^8}{70} + 2z^9,\end{aligned}\quad (18)$$

The Pade approximants of order $\left[\frac{3}{3} \right]$ in term of $x = \frac{1}{s}$, gives

$$\begin{aligned}\left[\frac{3}{3} \right] &= \frac{1}{\left(1 + \frac{1}{s^2} - \frac{2}{s} \right) s^2}, \\ \left[\frac{3}{3} \right] &= \frac{1}{\left(1 + \frac{1}{s^2} - \frac{2}{s} \right) s}, \\ \left[\frac{3}{3} \right] &= -\frac{1}{\left(1 + \frac{1}{s^2} - \frac{2}{s} \right) s^2} + \frac{2}{\left(1 + \frac{1}{s^2} - \frac{2}{s} \right) s}\end{aligned}\quad (19)$$

The exact solutions $u = (u_1(t), u_2(t), u_3(t)) = (te^t, e^t(1+t), e^t(2+t))$, are obtained by applying the inverse Laplace transform to the $\left[\frac{3}{3}\right]$ Pade approximate.

Example 4. 2: Given the system of nonlinear first-order equations [24],

$$\begin{aligned} u_1'(t) &= 2u_2^2(t), \\ u_2'(t) &= e^{-t}u_1(t), \\ u_3'(t) &= u_2(t) + u_3(t), \end{aligned} \quad (20)$$

subject to

$$u_1(0) = 1, u_2(0) = 1, \text{ and } u_3(0) = 0, \text{ with exact solutions } u = (u_1(t), u_2(t), u_3(t)) = (e^{2t}, e^t, te^t).$$

We will now construct the following homotopy equation, which is based on the algorithm presented in Section 2.

$$\begin{aligned} (1-p)\left[\frac{dv_1(t;p)}{dt} - (h;q)\left[\frac{dv_1(t;p)}{dt} - 2v_2^2(t;p)\right]\right], \\ (1-p)\left[\frac{dv_2(t;p)}{dt} - (h;q)\left[\frac{dv_2(t;p)}{dt} - e^{-t}v_1(t;p)\right]\right] \\ (1-p)\left[\frac{dv_3(t;p)}{dt} - (h;q)\left[\frac{dv_3(t;p)}{dt} - v_{21}(t;p) - v_3(t;p)\right]\right] \end{aligned} \quad (21)$$

The problem of zeroth order is expressed in Eqs. (22) as given below

$$\begin{aligned} u'_{1,0}(t) &= 0, & u_{1,0}(0) &= 1, \\ u'_{2,0}(t) &= 0, & u_{2,0}(0) &= 1, \\ u'_{3,0}(t) &= 0, & u_{3,0}(0) &= 0, \end{aligned} \quad (22)$$

and their solutions are

$$\begin{aligned} u_{1,0}(t) &= 1, \\ u_{2,0}(t) &= 1, \\ u_{3,0}(t) &= 0. \end{aligned} \quad (23)$$

According to the HPM procedure, we have the approximate solution of the 8'th order.

$$\begin{aligned} \tilde{u}_1(t) &= 1 + 2t + 2t^2 + \frac{4t^3}{3} + \frac{2t^4}{3} + \frac{4t^5}{15} + \frac{4t^6}{45} + \frac{8t^7}{315} + \frac{2t^8}{315}, \\ \tilde{u}_2(t) &= 1 + t + \frac{t^2}{2} + \frac{t^3}{6} + \frac{t^4}{24} + \frac{t^5}{120} + \frac{t^6}{720} + \frac{t^7}{5040} + \frac{t^8}{40320}, \\ \tilde{u}_3(t) &= t + t^2 + \frac{t^3}{2} + \frac{t^4}{6} + \frac{t^5}{24} + \frac{t^6}{120} + \frac{t^7}{720} + \frac{t^8}{5040}, \end{aligned} \quad (24)$$

which provides the precise solution of Eq. (20) as the number of terms approaches infinity, i.e. $\lim_{n \rightarrow \infty} \tilde{u}_n(t) = (e^{2t}, e^t, te^t)$, which are the. In order to enhance the accuracy of the HPM solution, we apply the Laplace transform to the first few terms of HPM solutions (24), as follows:

$$\begin{aligned} L(\tilde{u}_1(t)) &= \frac{128}{s^8} + \frac{64}{s^7} + \frac{32}{s^6} + \frac{16}{s^5} + \frac{8}{s^4} + \frac{4}{s^3} + \frac{2}{s^2} + \frac{1}{s}, \\ L(\tilde{u}_2(t)) &= \frac{1}{s^9} + \frac{1}{s^8} + \frac{1}{s^7} + \frac{1}{s^6} + \frac{1}{s^5} + \frac{1}{s^4} + \frac{1}{s^3} + \frac{1}{s^2} + \frac{1}{s}, \\ L(\tilde{u}_3(t)) &= \frac{8}{s^9} + \frac{7}{s^8} + \frac{6}{s^7} + \frac{5}{s^6} + \frac{4}{s^5} + \frac{3}{s^4} + \frac{2}{s^3} + \frac{1}{s^2}. \end{aligned} \quad (25)$$

Taking $s = \frac{1}{z}$, gives

$$\begin{aligned}
L(\tilde{u}_1(t)) &= z + 2z^2 + 4z^3 + 8z^4 + 16z^5 + 32z^6 + 64z^7 + 128z^8, \\
L(\tilde{u}_2(t)) &= z + z^2 + z^3 + z^4 + z^5 + z^6 + z^7 + z^8 + z^9, \\
L(\tilde{u}_3(t)) &= z^2 + 2z^3 + 3z^4 + 4z^5 + 5z^6 + 6z^7 + 7z^8 + 8z^9, \quad (26)
\end{aligned}$$

in term of $z = \frac{1}{s}$, the Pade approximates of $\left[\frac{3}{3}\right]$, are

$$\begin{aligned}
\left[\frac{3}{3}\right] \frac{1}{\left(1 - \frac{2}{s}\right)s}, \\
\left[\frac{3}{3}\right] &= \frac{1}{\left(1 - \frac{1}{s}\right)s}, \\
\left[\frac{3}{3}\right] &= \frac{1}{\left(1 + \frac{1}{s^2} - \frac{2}{s}\right)s^2}, \quad (27)
\end{aligned}$$

By applying the inverse, Laplace transforms into Eqs. (27), respectively, we obtain the exact solutions.

5. Results and Discussion

From the HPM solutions, we tabulated and formulated numerical results and discussions in Tables 1-6 and Figures 1-6. Our observations indicate that the accuracy of the solutions depends approximately the approximation terms, and the solutions converge to the exact ones when an infinitely large number of terms are considered. This implies that achieving higher accuracy necessitates additional computational work and effort. Consequently, to obtain accurate results, modifications should be made to the HPM procedure by incorporating Laplace transformations and Pade approximations, which provides exact solutions without the need to increase the number of approximation terms in the standard HPM.

Table 1: Numerical result of example 1

x	Exact Solution $u_1(t) = te^t$	Approximate Solution	HPM Absolute Error
0.0	0.0000000000	0.0000000000	0.000000000
0.2	0.2442805516	0.2442805476	4.04×10^{-9}
0.4	0.5967298791	0.5967293088	5.70×10^{-7}
0.6	1.0932712802	1.0932605520	1.07×10^{-5}
0.8	1.7804327428	1.7803444852	8.83×10^{-5}
1.0	2.7182818285	2.7178207672	4.61×10^{-4}

Table 2: Numerical result of example 1

t	Exact Solution $u_1(t) = (t + 1)e^t$	Approximate Solution	HPM Absolute Error
0.0	1.0000000000	1.0000000000	0.000000000
0.2	1.4656833098	1.465683299	1.17×10^{-8}
0.4	2.0885545767	2.0885533934	1.18×10^{-6}
0.6	2.9153900806	2.9153708834	1.92×10^{-5}
0.8	4.0059736713	4.0058288383	1.45×10^{-4}
1.0	5.4365636569	5.4358498677	7.14×10^{-4}

Table 3: Numerical result of example 1

t	Exact Solution $u_1(t) = (t+2)e^t$	Approximate Solution	HPM Absolute Error
0.0	2.0000000000	2.0000000000	0.00000000
0.2	2.6870860680	2.687086042	2.59×10^{-8}
0.4	3.5803792743	3.580377052	2.22×10^{-6}
0.6	4.737508881	4.737476270	3.26×10^{-5}
0.8	6.2315145998	6.231284915	2.30×10^{-4}
1.0	8.1548454854	8.153769211	1.07×10^{-3}

Table 4: Numerical result of example2

t	Exact Solution $u_1(t) = e^{2t}$	Approximate Solution	HPM Absolute Error
0.0	1.0000000000	1.0000000000	0.00
0.2	1.4918246976	1.4918246969	7.52×10^{-10}
0.4	2.2255409285	2.2255405267	4.02×10^{-7}
0.6	3.3201169227	3.3201007909	1.61×10^{-5}
0.8	4.9530324244	4.9528076759	2.25×10^{-5}
1.0	7.3890560989	7.3873015873	1.75×10^{-3}

Table 5: Numerical result of example 2

t	Exact Solution $u_1(t) = e^t$	Approximate Solution	HPM Absolute Error
0.0	1.000000000	1.000000000	0.0
0.2	1.2214027581	1.2214027581	1.44×10^{-12}
0.4	1.4918246976	1.4918246969	7.53×10^{-10}
0.6	1.8221188004	1.8221187709	2.95×10^{-8}
0.8	2.2255409285	2.2255405267	4.02×10^{-7}
1.0	2.7182818285	2.7182787698	3.06×10^{-6}

Table 6: Numerical result of example 1

t	Exact Solution $u_1(t) = te^t$	Approximate Solution	HPM Absolute Error
0.0	0.0000000000	0.0000000000	0.00
0.2	0.2442805516	0.2442805516	1.30×10^{-11}
0.4	0.5967298791	0.5967298723	6.80×10^{-9}
0.6	1.0932712802	1.0932710126	2.67×10^{-7}
0.8	1.7804327428	1.7804290926	3.65×10^{-6}
1.0	2.7182818285	2.7182539683	2.79×10^{-5}




Figure 1. Plot of a) Exact and approximate solutions b) Absolute errors for example 1




Figure 2. Plot of a) Exact and approximate solutions b) Absolute errors for example 2

6. Conclusion

In this research study, we present a novel methodology based on the HPM for solving a system of ordinary differential equations. This methodology not only demonstrates effectiveness and reliability, but also offers distinct advantages over alternative techniques. It is capable of accurately providing solutions for complex systems, making it a valuable tool for researchers and practitioners involved in the analysis of dynamic phenomena governed by such systems. Through illustrative examples and comparisons with numerical outcomes obtained from existing literature, we demonstrate that this methodology can achieve the exact analytical solution by using only a limited number of terms from the HPM truncated series solution. In conclusion, we firmly assert that this methodology proposes a robust and promising approach for handling various types of differential equations.

Funding: "This research received no external funding"

Conflicts of Interest: "The authors declare no conflict of interest."

References

- [1] Altaie, Sarmad A., Nidal Anakira, Ali Jameel, Osama Ababneh, Ahmad Qazza, and Abdel Kareem Alomari. "Homotopy analysis method analytical scheme for developing a solution to partial differential equations in fuzzy environment." *Fractal and Fractional* 6, no. 8 (2022): 419.
- [2] Marinca V, Herişanu N. Application of optimal homotopy asymptotic method for solving nonlinear equations arising in heat transfer. *International communications in heat and mass transfer*. 2008 Jul 1;35(6):710-5.
- [3] Jameela A, Anakira NR, Alomari AK, Hashim I, Shakhatreh MA. Numerical solution of n'th order fuzzy initial value problems by six stages. *Journal of Nonlinear Science Applications*. 2016;9(2):627-40. Biazar J, Ghazvini H, Eslami M. He's homotopy perturbation method for systems of integro-differential equations. *Chaos, Solitons & Fractals*. 2009 Feb 15;39(3):1253-8.
- [4] Alahmad, S., Anakira, N. R., Mamat, M., Sulaiman, I., & AlAhmad, R. (2022). Modified differential transformation method for solving classes of non-linear differential equations. *TWMS Journal of Applied and Engineering Mathematics*, 12(1), 107.
- [5] Hemeda AA. Homotopy perturbation method for solving systems of nonlinear coupled equations. *Applied Mathematical Sciences*. 2012;6(96):4787-800.
- [6] Amourah AA, Yousef F. Some properties of a class of analytic functions involving a new generalized differential operator. *Boletim da Sociedade Paranaense de Matematica*. 2020;38(6):33-42.
- [7] Jameel AF, Shather AH, Anakira NR, Alomari AK, Saaban A. Comparison for the approximate solution of the second-order fuzzy nonlinear differential equation with fuzzy initial conditions. *Math Stat* 8: 527–534.
- [8] Ramos JJ. On the variational iteration method and other iterative techniques for nonlinear differential equations. *Applied Mathematics and Computation*. 2008 May 15;199(1):39-69.
- [9] Anakira NR, Alomari AK, Hashim I. Application of optimal homotopy asymptotic method for solving linear delay differential equations. In *AIP Conference Proceedings* 2013 Nov 27 (Vol. 1571, No. 1, pp. 1013-1019). American Institute of Physics.
- [10] Shang X, Han D. Application of the variational iteration method for solving nth-order integro-differential equations. *Journal of Computational and Applied Mathematics*. 2010 Jul 1;234(5):1442-7.
- [11] Anakira N, Jameel A, Hijazi M, Alomari AK, Man N. A new approach for solving multi-pantograph type delay differential equations. *International Journal of Electrical & Computer Engineering* (2088-8708). 2022 Apr 1;12(2).
- [12] Abualhomos, M. "Numerical ways for solving fuzzy differential equations." *Int. J. Appl. Engin. Res* 13 (2018): 4610-4613. Al-Ahmad S, Anakira NR, Mamat M, Suliman IM, AlAhmad R. Modified differential transformation scheme for solving classes of non-linear differential equations. *TWMS J. Appl. Eng. Math*. 2021.
- [13] Ayaz F. Solutions of the system of differential equations by differential transform method. *Applied Mathematics and Computation*. 2004 Jan 12;147(2):547-67.
- [14] Stević S. On a system of difference equations. *Applied Mathematics and Computation*. 2011 Dec 1;218(7):3372-8.
- [15] Jameel, Ali Fareed, Nidal Ratib Anakira, A. K. Alomari, D. M. Alsharo, and Azizan Saaban. "New semi-analytical method for solving two point nth order fuzzy boundary value problem." *International Journal of Mathematical Modelling and Numerical Optimisation* 9, no. 1 (2019): 12-31.

- [16] Burqan, Aliaa, Mona Khandaqji, Zeyad Al-Zhour, Ahmad El-Ajou, and Tasneem Alrahamneh. "Analytical Approximate Solutions of Caputo Fractional KdV-Burgers Equations Using Laplace Residual Power Series Technique." *Journal of Applied Mathematics* 2024, no. 1 (2024): 7835548.
- [17] Amourah AA, Yousef F, Al-Hawary T, Darus M. A certain fractional derivative operator for p-valent functions and new class of analytic functions with negative coefficients. *Far East Journal of Mathematical Sciences*. 2016 Jan 1;99(1):75.
- [18] Subramanian M, Manigandan M, Tunç C, Gopal TN, Alzabut J. On system of nonlinear coupled differential equations and inclusions involving Caputo-type sequential derivatives of fractional order. *Journal of Taibah University for Science*. 2022 Dec 31;16(1):1-23.
- [19] Awawdeh, Fadi, M. Khandaqji, Z. Mustafa, and Mohammad Younis. "A new approach for the solution of the electrostatic potential differential equations." *Mathematical Problems in Engineering* 2009 (2009): 23.
- [20] Jameel, Ali, N. R. Anakira, A. K. Alomari, and Noraziah H. Man. "Solution and analysis of the fuzzy Volterra integral equations via homotopy analysis method." *Computer Modeling in Engineering & Sciences* 127, no. 3 (2021): 875-899.
- [21] Shams M, Kausar N, Alayyash K, Al-Shamiri MM, Arif N, Ismail R. Semi-analytical scheme for solving intuitionistic fuzzy system of differential equations. *IEEE Access*. 2023 Feb 1;11:33205-23.
- [22] He JH. Homotopy perturbation method: a new nonlinear analytical technique. *Applied Mathematics and computation*. 2003 Feb 15;135(1):73-9.
- [23] He JH. Homotopy perturbation technique. *Computer methods in applied mechanics and engineering*. 1999 Aug 1;178(3-4):257-62.
- [24] Shakeri, F., & Dehghan, M. (2008). Solution of delay differential equations via a homotopy perturbation method. *Mathematical and computer Modelling*, 48(3-4), 486-498.
- [25] Al-Ahmad S, Sulaiman IM, Mamat M, Ghazali PL. Modified differential transform scheme for solving systems of first order ordinary differential equations. *J. Math. Comput. Sci*. 2021;22:73-84.
- [26] Al-Ahmad, S., I. M. Sulaiman, M. Mamat, and L. G. Puspa. "A Modification of Differential Transform Method for Solving Systems of Second Order Ordinary Differential Equations." (2020).
- [27] Jameel, A. F., Anakira, N. R., Rashidi, M. M., Alomari, A. K., Saaban, A., & Shakhathreh, M. A. (2018). Differential Transformation Method For Solving High Order Fuzzy Initial Value Problems. *Italian Journal of Pure and Applied Mathematics*, 39, 194-208.
- [28] Jameel, A. F., Anakira, N. R., Shather, A. H., Saaban, A., & Alomari, A. K. (2020). Numerical algorithm for solving second order nonlinear fuzzy initial value problems. *International Journal of Electrical and Computer Engineering (IJECE)*, 10(6), 6497-6506.
- [29] Anakira, N. R. (2018). Optimal homotopy asymptotic method for solving multi-pantograph type delay differential equations. *Adv. Differ. Equ. Control Proc.* 19 (3), 191-204.
- [30] Anakira, N. R., Alomari, A. K., Jameel, A. F., & Hashim, I. (2017). Multistage optimal homotopy asymptotic method for solving boundary value problems with robin boundary conditions. *Far East Journal of Mathematical Sciences*, 102(8), 1727-1744.