



## Complex cubic neutrosophic set applied to subbisemiring and its extension of bisemiring

Brikena Vrioni<sup>1,\*</sup>, Nasreen Kausar<sup>2</sup>, Murugan Palanikumar<sup>3</sup>, Ervin Hoxha<sup>1</sup>

<sup>1</sup>School of Arts and Sciences, American International University, Kuwait

<sup>2</sup>Department of Mathematics, Faculty of Arts and Science, Yildiz Technical University, Esenler, 34220, Istanbul, Turkey

<sup>3</sup>Department of Mathematics, Saveetha School of Engineering, Saveetha Institute of Medical and Technical Sciences, Chennai-602105, India

Emails: [brikena.vrioni@yahoo.com](mailto:brikena.vrioni@yahoo.com); [kausar.nasreen57@gmail.com](mailto:kausar.nasreen57@gmail.com); [palanimaths86@gmail.com](mailto:palanimaths86@gmail.com); [ervin.hoxha1990@gmail.com](mailto:ervin.hoxha1990@gmail.com)

### Abstract

We construct the concept of complex cubic neutrosophic subbisemiring (ComCNSBS). We analyze the important properties and homomorphic aspects of ComCNSBS. For bisemirings, we propose the ComCNSBS level sets. A complex neutrosophic subset  $\Gamma$  if and only if each non-empty level set  $R^{(\varphi, \kappa)}$ , where  $R = (\widehat{\Re}_\Gamma^T \cdot e^{i\theta\Im_\Gamma^T}, \widehat{\Re}_\Gamma^I \cdot e^{i\theta\Im_\Gamma^I}, \widehat{\Re}_\Gamma^F \cdot e^{i\theta\Im_\Gamma^F}, \Re_\Gamma^T \cdot e^{i\theta\Im_\Gamma^T}, \Re_\Gamma^I \cdot e^{i\theta\Im_\Gamma^I}, \Re_\Gamma^F \cdot e^{i\theta\Im_\Gamma^F})$  is a ComCNSBS. We show that homomorphic images of all ComCNSBSs are ComCNSBSs, and homomorphic pre-images of all ComCNSBSs are ComCNSBSs. We illustrate the practical significance of our findings.

**Keywords:** ComCNSBS; ComCNNSBS; SBS; homomorphism

### 1 Introduction

Fuzzy set (FS) theory was developed by Zadeh<sup>1</sup> and is most effective in managing ambiguity and uncertainty. An element in an FS is considered a member if it contains a single value inside the interval. However, since resistance can exist in real-world situations, the degree of non-membership might not necessarily equal one minus the degree of membership. As FS theory advances quickly, more and more hybrid fuzzy models are being developed. Numerous uncertain theories, including FS,<sup>1</sup> intuitionistic FS (IFS),<sup>2</sup> Pythagorean FS (PFS),<sup>3</sup> and spherical FS (SFS),<sup>4</sup> have been developed as a result of the uncertainties. Sets with grades ranging from 0 to 1, referred to as MG, comprise an FS. IFS is classified as MG, despite Atanassov<sup>2</sup> stating that non-membership grades (NMG) can only have a value of 1. There is a chance that, in a decision-making process, the sum of MGs and NMGs will sometimes exceed 1. The generalized MG and NMG logic, which has a value not exceeding 1 and is determined by the square of the MGs and NMGs, was constructed by Yager<sup>3</sup> using

PFS logic. These ideas are unable to explain the neutral state, which is neither positive nor negative. Cuong<sup>5</sup> discussed the picture fuzzy set. Three grading points were utilized by FS: positive, neutral, and negative. These grades added together could not total more than 1. For certain reasons, it is also equivalent to PFS or IFS. It is an independent generalization of three models that deals with the truth, indeterminacy, and falsity of FS and IFS. Numerous recent studies have been discussed by Palanikumar et al.<sup>6-8</sup>

Smarandache<sup>9</sup> was developed by neutrosophic set (NS) to deal with contradicting and ambiguous data. This reasoning establishes the degree to which a statement is true, ambiguous, or untrue. Ramot et al.<sup>10</sup> presented the notion of the complex fuzzy set (CFS). There is a wide range of possible values for the membership functions of CFSs deals. The unit circle of the complex plane is extended to  $[0, 1]$ , but the unit circle of a fuzzy membership function is fixed. Rather than extending exclusively to  $[0, 1]$ , the membership function  $\mu_X(x)$  of the CFS  $X$  extends to the unit circle in the complex plane. Thus,  $\mu_X(x)$  is a complex-valued function that assigns a MG of the form  $\eta_X(x) \cdot e^{i\tau_X(x)}$ , where  $i = \sqrt{-1}$ , to any element  $x$  in the universal set. The value of  $\mu_X(x)$  is defined by the two real-valued variables,  $\eta_X(x)$  and  $\tau_X(x)$ , where  $\eta_X(x) \in [0, 1]$ . Golan<sup>11</sup> established the concept of semiring logic and its applications. Hussian and colleagues<sup>12</sup> talked about the concept and expansion of bisemirings. The topic of bipolar-valued FSs and related techniques is covered by Lee.<sup>13</sup> Ahsan et al.<sup>14</sup> investigated fuzzy semirings. The notion of bisemirings was first presented by Sen et al.<sup>16</sup> Recently, the new type of normal subbisemiring (SBS) were presented by Palanikumar et al.,<sup>17</sup> bipolar-valued neutrosophic normal SBS.<sup>18</sup> In this study, the following contributions are made:

1. Every ComCNSBS's intersection results in another ComCNSBS.
2. Let  $\Upsilon$  be the strongest complex neutrosophic relation of  $S$ , and let  $\Gamma$  be a ComCNSBS of  $S$ . If and only if  $\Upsilon$  is a ComCNSBS of  $S$ , then  $\Gamma$  is a ComCNSBS of bisemiring  $S \times S$ .
3.  $R = (\widehat{\mathfrak{R}}_T^T \cdot e^{i\theta\widehat{\mathfrak{S}}_T^T}, \widehat{\mathfrak{R}}_T^I \cdot e^{i\theta\widehat{\mathfrak{S}}_T^I}, \widehat{\mathfrak{R}}_T^F \cdot e^{i\theta\widehat{\mathfrak{S}}_T^F}, \widehat{\mathfrak{R}}_T^T \cdot e^{i\theta\widehat{\mathfrak{S}}_T^T}, \widehat{\mathfrak{R}}_T^I \cdot e^{i\theta\widehat{\mathfrak{S}}_T^I}, \widehat{\mathfrak{R}}_T^F \cdot e^{i\theta\widehat{\mathfrak{S}}_T^F})$  is a SBS of  $S$  for all  $\varphi, \varkappa \in \mathbb{D}[0, 1]$ .
4. Every ComCNSBS has a homomorphic image that is also a ComCNSBS, and each ComCNSBS has a homomorphic preimage that is also a ComCNSBS.

We will look at certain aspects of the SBS and ComCNSBS concepts and make some conclusions. The following five sections make up the article. In Section 1, semirings and SBS are introduced. Section 2 contains information on semiring and SBS preparation. Listing of ComCNSBS properties is done in Section 3. Utilizing numerical examples are recommended for ComCNSBS evaluation. Section 4 indicates the outcome and future direction.

## 2 Preliminaries

**Definition 2.1.**<sup>16</sup> The structure  $(S, \uplus, \ominus, \odot)$  is a bisemiring, if the conditions such as  $(S, \uplus, \ominus)$  and  $(S, \ominus, \odot)$  both semirings. That is,  $(S, \uplus)$ ,  $(S, \ominus)$  and  $(S, \odot)$  both semigroups and

1.  $\ell_v \ominus (\ell_\zeta \uplus \ell_\eta) = (\ell_v \ominus \ell_\zeta) \uplus (\ell_v \ominus \ell_\eta)$ ,
2.  $(\ell_\zeta \uplus \ell_\eta) \ominus \ell_v = (\ell_\zeta \ominus \ell_v) \uplus (\ell_\eta \ominus \ell_v)$ ,
3.  $\ell_v \odot (\ell_\zeta \ominus \ell_\eta) = (\ell_v \odot \ell_\zeta) \ominus (\ell_v \odot \ell_\eta)$ ,
4.  $(\ell_\zeta \ominus \ell_\eta) \odot \ell_v = (\ell_\zeta \odot \ell_v) \ominus (\ell_\eta \odot \ell_v), \forall \ell_v, \ell_\zeta, \ell_\eta \in S$ .

**Definition 2.2.**<sup>9</sup> A NS  $v$  in the universe  $\mathcal{U}$  is  $v = \{j, u_v^T(x), u_v^I(x), u_v^F(x) | x \in \mathcal{U}\}$ , where  $u_v^T(x), u_v^I(x), u_v^F(x)$  represents the truth, intermediate and false grade of  $v$  respectively, and  $u_v^T : \mathcal{U} \rightarrow [0, 1], u_v^I : \mathcal{U} \rightarrow [0, 1], u_v^F : \mathcal{U} \rightarrow [0, 1]$  and  $0 \leq u_v^T(x) + u_v^I(x) + u_v^F(x) \leq 3$ .

**Definition 2.3.**<sup>9</sup> Let  $\psi_1 = \langle u_{\psi_1}^T, u_{\psi_1}^I, u_{\psi_1}^F \rangle, \psi_2 = \langle u_{\psi_2}^T, u_{\psi_2}^I, u_{\psi_2}^F \rangle$  and  $\psi_3 = \langle u_{\psi_3}^T, u_{\psi_3}^I, u_{\psi_3}^F \rangle$  are the neutrosophic numbers of  $\mathcal{U}$ . Then

1.  $\psi_2 \sqcup \psi_3 = \left\langle \max(j_{\psi_2}^T, u_{\psi_3}^T), \min(j_{\psi_2}^I, u_{\psi_3}^I), \min(j_{\psi_2}^F, u_{\psi_3}^F) \right\rangle$ ,
2.  $\psi_2 \sqcap \psi_3 = \left\langle \min(j_{\psi_2}^T, u_{\psi_3}^T), \max(j_{\psi_2}^I, u_{\psi_3}^I), \max(j_{\psi_2}^F, u_{\psi_3}^F) \right\rangle$ ,
3.  $\psi_2 \geq \psi_3$  iff  $u_{\psi_2}^T \geq u_{\psi_3}^T$  and  $u_{\psi_2}^I \leq u_{\psi_3}^I$  and  $u_{\psi_2}^F \leq u_{\psi_3}^F$ ,
4.  $\psi_2 = \psi_3$  iff  $u_{\psi_2}^T = u_{\psi_3}^T$  and  $u_{\psi_2}^I = u_{\psi_3}^I$  and  $u_{\psi_2}^F = u_{\psi_3}^F$ .

**Definition 2.4.** <sup>9</sup> If  $\psi = \{x, j_v^T(x), j_v^I(x), j_v^F(x)\}$  of  $\mathcal{U}$  be the NS, then  $(\zeta, \tau)$ -cut is described as  $\{x \in U | j_v^T(x) \geq \zeta, j_v^I(x) \geq \zeta, j_v^F(x) \leq \tau\}$ .

**Definition 2.5.** <sup>9</sup> If  $V$  and  $W$  be NSs of  $S$ , then Cartesian product is described as  $V \times W = \{j_{V \times W}^T(j, \psi), j_{V \times W}^I(j, \psi), j_{V \times W}^F(j, \psi) | j \in S\}$ , where

$$j_{V \times W}^T(j, \psi) = \min\{j_V^T(x), j_W^T(\psi)\}, j_{V \times W}^I(j, \psi) = \frac{j_V^I(x) + j_W^I(\psi)}{2}, j_{V \times W}^F(j, \psi) = \max\{j_V^F(x), j_W^F(\psi)\}.$$

### 3 Complex cubic neutrosophic SBS

Unless otherwise indicated,  $S$  denotes the bisemiring,  $\Re$  means the real part, and  $\Im$  refers the imaginary part and  $\theta = 2\pi$ .

**Definition 3.1.** The complex cubic NS (ComCNS)  $\Gamma$  in universal set  $\mathcal{D}$ ,

$\Gamma = \{j, \widehat{\Re}_\Gamma^T(j) \cdot e^{i\theta \widehat{\Im}_\Gamma^T(j)}, \widehat{\Re}_\Gamma^I(j) \cdot e^{i\theta \widehat{\Im}_\Gamma^I(j)}, \widehat{\Re}_\Gamma^F(j) \cdot e^{i\theta \widehat{\Im}_\Gamma^F(j)}, \Re_\Gamma^T(j) \cdot e^{i\theta \Im_\Gamma^T(j)}, \Re_\Gamma^I(j) \cdot e^{i\theta \Im_\Gamma^I(j)}, \Re_\Gamma^F(j) \cdot e^{i\theta \Im_\Gamma^F(j)} : j \in \mathcal{D}\}$ , where  $\widehat{\Re}_\Gamma^T(j) = [\Re_\Gamma^{TL}, \Re_\Gamma^{TU}]$ ,  $\widehat{\Re}_\Gamma^I(j) = [\Re_\Gamma^{IL}, \Re_\Gamma^{IU}]$ ,  $\widehat{\Re}_\Gamma^F(j) = [\Re_\Gamma^{FL}, \Re_\Gamma^{FU}]$  and  $\widehat{\Re}_\Gamma^T(j) \cdot e^{i\theta \widehat{\Im}_\Gamma^T(j)}, \widehat{\Re}_\Gamma^I(j) \cdot e^{i\theta \widehat{\Im}_\Gamma^I(j)}, \widehat{\Re}_\Gamma^F(j) \cdot e^{i\theta \widehat{\Im}_\Gamma^F(j)} : \mathcal{D} \rightarrow D[0, 1]$  and  $\Re_\Gamma^T(j) \cdot e^{i\theta \Im_\Gamma^T(j)}, \Re_\Gamma^I(j) \cdot e^{i\theta \Im_\Gamma^I(j)}, \Re_\Gamma^F(j) \cdot e^{i\theta \Im_\Gamma^F(j)} : \mathcal{D} \rightarrow [0, 1]$  represents the degree of truth, indeterminacy and false, respectively. For simplicity the symbol  $\widehat{\Re}_\Gamma^T, \widehat{\Re}_\Gamma^I, \widehat{\Re}_\Gamma^F$  is ComCNS  $\Gamma = \{j, \widehat{\Re}_\Gamma^T(j) \cdot e^{i\theta \widehat{\Im}_\Gamma^T(j)}, \widehat{\Re}_\Gamma^I(j) \cdot e^{i\theta \widehat{\Im}_\Gamma^I(j)}, \widehat{\Re}_\Gamma^F(j) \cdot e^{i\theta \widehat{\Im}_\Gamma^F(j)}, \Re_\Gamma^T(j) \cdot e^{i\theta \Im_\Gamma^T(j)}, \Re_\Gamma^I(j) \cdot e^{i\theta \Im_\Gamma^I(j)}, \Re_\Gamma^F(j) \cdot e^{i\theta \Im_\Gamma^F(j)} : j \in \mathcal{D}\}$ .

**Definition 3.2.** Let  $\Gamma = \{j, \widehat{\Re}_\Gamma^T(j) \cdot e^{i\theta \widehat{\Im}_\Gamma^T(j)}, \widehat{\Re}_\Gamma^I(j) \cdot e^{i\theta \widehat{\Im}_\Gamma^I(j)}, \widehat{\Re}_\Gamma^F(j) \cdot e^{i\theta \widehat{\Im}_\Gamma^F(j)}, \Re_\Gamma^T(j) \cdot e^{i\theta \Im_\Gamma^T(j)}, \Re_\Gamma^I(j) \cdot e^{i\theta \Im_\Gamma^I(j)}, \Re_\Gamma^F(j) \cdot e^{i\theta \Im_\Gamma^F(j)}\}$  and  $\Lambda = \{j, \widehat{\Re}_\Lambda^T(j) \cdot e^{i\theta \widehat{\Im}_\Lambda^T(j)}, \widehat{\Re}_\Lambda^I(j) \cdot e^{i\theta \widehat{\Im}_\Lambda^I(j)}, \widehat{\Re}_\Lambda^F(j) \cdot e^{i\theta \widehat{\Im}_\Lambda^F(j)}, \Re_\Lambda^T(j) \cdot e^{i\theta \Im_\Lambda^T(j)}, \Re_\Lambda^I(j) \cdot e^{i\theta \Im_\Lambda^I(j)}, \Re_\Lambda^F(j) \cdot e^{i\theta \Im_\Lambda^F(j)}\}$  be two ComCNSs of  $\mathcal{D}$ . Then we define the intersection and union operation is defined as

(i)  $\Gamma \cap \Lambda = \left\{ \left( j, \min\{\widehat{\Re}_\Gamma^T(j) \cdot e^{i\theta \widehat{\Im}_\Gamma^T(j)}, \widehat{\Re}_\Lambda^T(j) \cdot e^{i\theta \widehat{\Im}_\Lambda^T(j)}\}, \min\{\widehat{\Re}_\Gamma^I(j) \cdot e^{i\theta \widehat{\Im}_\Gamma^I(j)}, \widehat{\Re}_\Lambda^I(j) \cdot e^{i\theta \widehat{\Im}_\Lambda^I(j)}\}, \max\{\widehat{\Re}_\Gamma^F(j) \cdot e^{i\theta \widehat{\Im}_\Gamma^F(j)}, \widehat{\Re}_\Lambda^F(j) \cdot e^{i\theta \widehat{\Im}_\Lambda^F(j)}\}, \min\{\Re_\Gamma^T(j) \cdot e^{i\theta \Im_\Gamma^T(j)}, \Re_\Lambda^T(j) \cdot e^{i\theta \Im_\Lambda^T(j)}\}, \min\{\Re_\Gamma^I(j) \cdot e^{i\theta \Im_\Gamma^I(j)}, \Re_\Lambda^I(j) \cdot e^{i\theta \Im_\Lambda^I(j)}\}, \max\{\Re_\Gamma^F(j) \cdot e^{i\theta \Im_\Gamma^F(j)}, \Re_\Lambda^F(j) \cdot e^{i\theta \Im_\Lambda^F(j)}\} \right) | j \in \mathcal{D} \right\}$ .

(ii)  $\Gamma \cup \Lambda = \left\{ \left( j, \max\{\widehat{\Re}_\Gamma^T(j) \cdot e^{i\theta \widehat{\Im}_\Gamma^T(j)}, \widehat{\Re}_\Lambda^T(j) \cdot e^{i\theta \widehat{\Im}_\Lambda^T(j)}\}, \max\{\widehat{\Re}_\Gamma^I(j) \cdot e^{i\theta \widehat{\Im}_\Gamma^I(j)}, \widehat{\Re}_\Lambda^I(j) \cdot e^{i\theta \widehat{\Im}_\Lambda^I(j)}\}, \min\{\widehat{\Re}_\Gamma^F(j) \cdot e^{i\theta \widehat{\Im}_\Gamma^F(j)}, \widehat{\Re}_\Lambda^F(j) \cdot e^{i\theta \widehat{\Im}_\Lambda^F(j)}\}, \max\{\Re_\Gamma^T(j) \cdot e^{i\theta \Im_\Gamma^T(j)}, \Re_\Lambda^T(j) \cdot e^{i\theta \Im_\Lambda^T(j)}\}, \max\{\Re_\Gamma^I(j) \cdot e^{i\theta \Im_\Gamma^I(j)}, \Re_\Lambda^I(j) \cdot e^{i\theta \Im_\Lambda^I(j)}\}, \min\{\Re_\Gamma^F(j) \cdot e^{i\theta \Im_\Gamma^F(j)}, \Re_\Lambda^F(j) \cdot e^{i\theta \Im_\Lambda^F(j)}\} \right) | j \in \mathcal{D} \right\}$ .

**Definition 3.3.** The ComCNS  $\Gamma = \{j, \widehat{\Re}_\Gamma^T(j) \cdot e^{i\theta \widehat{\Im}_\Gamma^T(j)}, \widehat{\Re}_\Gamma^I(j) \cdot e^{i\theta \widehat{\Im}_\Gamma^I(j)}, \widehat{\Re}_\Gamma^F(j) \cdot e^{i\theta \widehat{\Im}_\Gamma^F(j)}, \Re_\Gamma^T(j) \cdot e^{i\theta \Im_\Gamma^T(j)}, \Re_\Gamma^I(j) \cdot e^{i\theta \Im_\Gamma^I(j)}, \Re_\Gamma^F(j) \cdot e^{i\theta \Im_\Gamma^F(j)}\}$  of  $\mathcal{D}$ . Then  $(\wp, \varkappa)$ -cut is described as  $\{j \in \mathcal{D} | \widehat{\Re}_\Gamma^T(j) \cdot e^{i\theta \widehat{\Im}_\Gamma^T(j)} \geq \wp, \widehat{\Re}_\Gamma^I(j) \cdot e^{i\theta \widehat{\Im}_\Gamma^I(j)} \geq \wp, \widehat{\Re}_\Gamma^F(j) \cdot e^{i\theta \widehat{\Im}_\Gamma^F(j)} \leq \varkappa, \Re_\Gamma^T(j) \cdot e^{i\theta \Im_\Gamma^T(j)} \geq \wp, \Re_\Gamma^I(j) \cdot e^{i\theta \Im_\Gamma^I(j)} \geq \wp, \Re_\Gamma^F(j) \cdot e^{i\theta \Im_\Gamma^F(j)} \leq \varkappa\}$ .

**Definition 3.4.** The Cartesian product of  $\Gamma$  and  $\Lambda$  is described as

$$\Gamma \times \Lambda = \left\{ \widehat{\Re}_{\Gamma \times \Lambda}^T((j, \psi)) \cdot e^{i\theta \widehat{\Im}_{\Gamma \times \Lambda}^T((j, \psi))}, \widehat{\Re}_{\Gamma \times \Lambda}^I((j, \psi)) \cdot e^{i\theta \widehat{\Im}_{\Gamma \times \Lambda}^I((j, \psi))}, \widehat{\Re}_{\Gamma \times \Lambda}^F(j, \psi) \cdot e^{i\theta \widehat{\Im}_{\Gamma \times \Lambda}^F((j, \psi))}, \Re_{\Gamma \times \Lambda}^T((j, \psi)), \Re_{\Gamma \times \Lambda}^I((j, \psi)), \Re_{\Gamma \times \Lambda}^F(j, \psi) \right\}.$$

$e^{i\theta\mathfrak{S}_{\Gamma\times\Lambda}^T((j,\psi))}, \mathfrak{R}_{\Gamma\times\Lambda}^I((j,\psi)) \cdot e^{i\theta\mathfrak{S}_{\Gamma\times\Lambda}^I((j,\psi))}, \mathfrak{R}_{\Gamma\times\Lambda}^F(j,\psi) \cdot e^{i\theta\mathfrak{S}_{\Gamma\times\Lambda}^F((j,\psi))} \mid \text{for all } j, \psi \in S\}$ , where  $\Gamma$  and  $\Lambda$  be the ComCNS of  $\mathfrak{D}$ , where

$$\left\{ \begin{array}{l} \widehat{\mathfrak{R}_{\Gamma\times\Lambda}^T((j,\psi))} \cdot e^{i\theta\widehat{\mathfrak{S}_{\Gamma\times\Lambda}^T((j,\psi))}} = \min \left\{ \widehat{\mathfrak{R}_{\Gamma}^T(j)} \cdot e^{i\theta\widehat{\mathfrak{S}_{\Gamma}^T(j)}}, \widehat{\mathfrak{R}_{\Lambda}^T(\psi)} \cdot e^{i\theta\widehat{\mathfrak{S}_{\Lambda}^T(\psi)}} \right\} \\ \widehat{\mathfrak{R}_{\Gamma\times\Lambda}^I((j,\psi))} \cdot e^{i\theta\widehat{\mathfrak{S}_{\Gamma\times\Lambda}^I((j,\psi))}} = \frac{\widehat{\mathfrak{R}_{\Gamma}^I(j)} \cdot e^{i\theta\widehat{\mathfrak{S}_{\Gamma}^I(j)}} + \widehat{\mathfrak{R}_{\Lambda}^I(\psi)} \cdot e^{i\theta\widehat{\mathfrak{S}_{\Lambda}^I(\psi)}}}{2} \\ \widehat{\mathfrak{R}_{\Gamma\times\Lambda}^F((j,\psi))} \cdot e^{i\theta\widehat{\mathfrak{S}_{\Gamma\times\Lambda}^F((j,\psi))}} = \max \left\{ \widehat{\mathfrak{R}_{\Gamma}^F(j)} \cdot e^{i\theta\widehat{\mathfrak{S}_{\Gamma}^F(j)}}, \widehat{\mathfrak{R}_{\Lambda}^F(\psi)} \cdot e^{i\theta\widehat{\mathfrak{S}_{\Lambda}^F(\psi)}} \right\} \end{array} \right\}$$

$$\left\{ \begin{array}{l} \mathfrak{R}_{\Gamma\times\Lambda}^T((j,\psi)) \cdot e^{i\theta\mathfrak{S}_{\Gamma\times\Lambda}^T((j,\psi))} = \min \left\{ \mathfrak{R}_{\Gamma}^T(j) \cdot e^{i\theta\mathfrak{S}_{\Gamma}^T(j)}, \mathfrak{R}_{\Lambda}^T(\psi) \cdot e^{i\theta\mathfrak{S}_{\Lambda}^T(\psi)} \right\} \\ \mathfrak{R}_{\Gamma\times\Lambda}^I((j,\psi)) \cdot e^{i\theta\mathfrak{S}_{\Gamma\times\Lambda}^I((j,\psi))} = \frac{\mathfrak{R}_{\Gamma}^I(j) \cdot e^{i\theta\mathfrak{S}_{\Gamma}^I(j)} + \mathfrak{R}_{\Lambda}^I(\psi) \cdot e^{i\theta\mathfrak{S}_{\Lambda}^I(\psi)}}{2} \\ \mathfrak{R}_{\Gamma\times\Lambda}^F((j,\psi)) \cdot e^{i\theta\mathfrak{S}_{\Gamma\times\Lambda}^F((j,\psi))} = \max \left\{ \mathfrak{R}_{\Gamma}^F(j) \cdot e^{i\theta\mathfrak{S}_{\Gamma}^F(j)}, \mathfrak{R}_{\Lambda}^F(\psi) \cdot e^{i\theta\mathfrak{S}_{\Lambda}^F(\psi)} \right\} \end{array} \right\}$$

**Definition 3.5.** Let  $\Gamma$  be the ComCNS of  $S$  is called as ComCNSBS if

$$\left\{ \begin{array}{l} \widehat{\mathfrak{R}_{\Gamma}^T((j \boxplus_1 \psi))} \cdot e^{i\theta\widehat{\mathfrak{S}_{\Gamma}^T((j \boxplus_1 \psi))}} \geq \min \{ \widehat{\mathfrak{R}_{\Gamma}^T(j)} \cdot e^{i\theta\widehat{\mathfrak{S}_{\Gamma}^T(j)}}, \widehat{\mathfrak{R}_{\Gamma}^T(\psi)} \cdot e^{i\theta\widehat{\mathfrak{S}_{\Gamma}^T(\psi)}} \} \\ \widehat{\mathfrak{R}_{\Gamma}^T((j \boxplus_2 \psi))} \cdot e^{i\theta\widehat{\mathfrak{S}_{\Gamma}^T((j \boxplus_2 \psi))}} \geq \min \{ \widehat{\mathfrak{R}_{\Gamma}^T(j)} \cdot e^{i\theta\widehat{\mathfrak{S}_{\Gamma}^T(j)}}, \widehat{\mathfrak{R}_{\Gamma}^T(\psi)} \cdot e^{i\theta\widehat{\mathfrak{S}_{\Gamma}^T(\psi)}} \} \\ \widehat{\mathfrak{R}_{\Gamma}^T((j \boxplus_3 \psi))} \cdot e^{i\theta\widehat{\mathfrak{S}_{\Gamma}^T((j \boxplus_3 \psi))}} \geq \min \{ \widehat{\mathfrak{R}_{\Gamma}^T(j)} \cdot e^{i\theta\widehat{\mathfrak{S}_{\Gamma}^T(j)}}, \widehat{\mathfrak{R}_{\Gamma}^T(\psi)} \cdot e^{i\theta\widehat{\mathfrak{S}_{\Gamma}^T(\psi)}} \} \end{array} \right\}$$

$$\left\{ \begin{array}{l} \widehat{\mathfrak{R}_{\Gamma}^I((j \boxplus_1 \psi))} \cdot e^{i\theta\widehat{\mathfrak{S}_{\Gamma}^I((j \boxplus_1 \psi))}} \geq \frac{\widehat{\mathfrak{R}_{\Gamma}^I(j)} \cdot e^{i\theta\widehat{\mathfrak{S}_{\Gamma}^I(j)}} + \widehat{\mathfrak{R}_{\Gamma}^I(\psi)} \cdot e^{i\theta\widehat{\mathfrak{S}_{\Gamma}^I(\psi)}}}{2} \\ \text{OR} \\ \widehat{\mathfrak{R}_{\Gamma}^I((j \boxplus_2 \psi))} \cdot e^{i\theta\widehat{\mathfrak{S}_{\Gamma}^I((j \boxplus_2 \psi))}} \geq \frac{\widehat{\mathfrak{R}_{\Gamma}^I(j)} \cdot e^{i\theta\widehat{\mathfrak{S}_{\Gamma}^I(j)}} + \widehat{\mathfrak{R}_{\Gamma}^I(\psi)} \cdot e^{i\theta\widehat{\mathfrak{S}_{\Gamma}^I(\psi)}}}{2} \\ \text{OR} \\ \widehat{\mathfrak{R}_{\Gamma}^I((j \boxplus_3 \psi))} \cdot e^{i\theta\widehat{\mathfrak{S}_{\Gamma}^I((j \boxplus_3 \psi))}} \geq \frac{\widehat{\mathfrak{R}_{\Gamma}^I(j)} \cdot e^{i\theta\widehat{\mathfrak{S}_{\Gamma}^I(j)}} + \widehat{\mathfrak{R}_{\Gamma}^I(\psi)} \cdot e^{i\theta\widehat{\mathfrak{S}_{\Gamma}^I(\psi)}}}{2} \end{array} \right\}$$

$$\left\{ \begin{array}{l} \widehat{\mathfrak{R}_{\Gamma}^F((j \boxplus_1 \psi))} \cdot e^{i\theta\widehat{\mathfrak{S}_{\Gamma}^F((j \boxplus_1 \psi))}} \leq \max \{ \widehat{\mathfrak{R}_{\Gamma}^F(j)} \cdot e^{i\theta\widehat{\mathfrak{S}_{\Gamma}^F(j)}}, \widehat{\mathfrak{R}_{\Gamma}^F(\psi)} \cdot e^{i\theta\widehat{\mathfrak{S}_{\Gamma}^F(\psi)}} \} \\ \widehat{\mathfrak{R}_{\Gamma}^F((j \boxplus_2 \psi))} \cdot e^{i\theta\widehat{\mathfrak{S}_{\Gamma}^F((j \boxplus_2 \psi))}} \leq \max \{ \widehat{\mathfrak{R}_{\Gamma}^F(j)} \cdot e^{i\theta\widehat{\mathfrak{S}_{\Gamma}^F(j)}}, \widehat{\mathfrak{R}_{\Gamma}^F(\psi)} \cdot e^{i\theta\widehat{\mathfrak{S}_{\Gamma}^F(\psi)}} \} \\ \widehat{\mathfrak{R}_{\Gamma}^F((j \boxplus_3 \psi))} \cdot e^{i\theta\widehat{\mathfrak{S}_{\Gamma}^F((j \boxplus_3 \psi))}} \leq \max \{ \widehat{\mathfrak{R}_{\Gamma}^F(j)} \cdot e^{i\theta\widehat{\mathfrak{S}_{\Gamma}^F(j)}}, \widehat{\mathfrak{R}_{\Gamma}^F(\psi)} \cdot e^{i\theta\widehat{\mathfrak{S}_{\Gamma}^F(\psi)}} \} \end{array} \right\}$$

$$\left\{ \begin{array}{l} \mathfrak{R}_{\Gamma}^T((j \boxplus_1 \psi)) \cdot e^{i\theta\mathfrak{S}_{\Gamma}^T((j \boxplus_1 \psi))} \geq \min \{ \mathfrak{R}_{\Gamma}^T(j) \cdot e^{i\theta\mathfrak{S}_{\Gamma}^T(j)}, \mathfrak{R}_{\Gamma}^T(\psi) \cdot e^{i\theta\mathfrak{S}_{\Gamma}^T(\psi)} \} \\ \mathfrak{R}_{\Gamma}^T((j \boxplus_2 \psi)) \cdot e^{i\theta\mathfrak{S}_{\Gamma}^T((j \boxplus_2 \psi))} \geq \min \{ \mathfrak{R}_{\Gamma}^T(j) \cdot e^{i\theta\mathfrak{S}_{\Gamma}^T(j)}, \mathfrak{R}_{\Gamma}^T(\psi) \cdot e^{i\theta\mathfrak{S}_{\Gamma}^T(\psi)} \} \\ \mathfrak{R}_{\Gamma}^T((j \boxplus_3 \psi)) \cdot e^{i\theta\mathfrak{S}_{\Gamma}^T((j \boxplus_3 \psi))} \geq \min \{ \mathfrak{R}_{\Gamma}^T(j) \cdot e^{i\theta\mathfrak{S}_{\Gamma}^T(j)}, \mathfrak{R}_{\Gamma}^T(\psi) \cdot e^{i\theta\mathfrak{S}_{\Gamma}^T(\psi)} \} \end{array} \right\}$$

$$\left\{ \begin{array}{l} \mathfrak{R}_{\Gamma}^I((j \boxplus_1 \psi)) \cdot e^{i\theta\mathfrak{S}_{\Gamma}^I((j \boxplus_1 \psi))} \geq \frac{\mathfrak{R}_{\Gamma}^I(j) \cdot e^{i\theta\mathfrak{S}_{\Gamma}^I(j)} + \mathfrak{R}_{\Gamma}^I(\psi) \cdot e^{i\theta\mathfrak{S}_{\Gamma}^I(\psi)}}{2} \\ \text{OR} \\ \mathfrak{R}_{\Gamma}^I((j \boxplus_2 \psi)) \cdot e^{i\theta\mathfrak{S}_{\Gamma}^I((j \boxplus_2 \psi))} \geq \frac{\mathfrak{R}_{\Gamma}^I(j) \cdot e^{i\theta\mathfrak{S}_{\Gamma}^I(j)} + \mathfrak{R}_{\Gamma}^I(\psi) \cdot e^{i\theta\mathfrak{S}_{\Gamma}^I(\psi)}}{2} \\ \text{OR} \\ \mathfrak{R}_{\Gamma}^I((j \boxplus_3 \psi)) \cdot e^{i\theta\mathfrak{S}_{\Gamma}^I((j \boxplus_3 \psi))} \geq \frac{\mathfrak{R}_{\Gamma}^I(j) \cdot e^{i\theta\mathfrak{S}_{\Gamma}^I(j)} + \mathfrak{R}_{\Gamma}^I(\psi) \cdot e^{i\theta\mathfrak{S}_{\Gamma}^I(\psi)}}{2} \end{array} \right\}$$

$$\left\{ \begin{array}{l} \mathfrak{R}_{\Gamma}^F((j \boxplus_1 \psi)) \cdot e^{i\theta\mathfrak{S}_{\Gamma}^F((j \boxplus_1 \psi))} \leq \max \{ \mathfrak{R}_{\Gamma}^F(j) \cdot e^{i\theta\mathfrak{S}_{\Gamma}^F(j)}, \mathfrak{R}_{\Gamma}^F(\psi) \cdot e^{i\theta\mathfrak{S}_{\Gamma}^F(\psi)} \} \\ \mathfrak{R}_{\Gamma}^F((j \boxplus_2 \psi)) \cdot e^{i\theta\mathfrak{S}_{\Gamma}^F((j \boxplus_2 \psi))} \leq \max \{ \mathfrak{R}_{\Gamma}^F(j) \cdot e^{i\theta\mathfrak{S}_{\Gamma}^F(j)}, \mathfrak{R}_{\Gamma}^F(\psi) \cdot e^{i\theta\mathfrak{S}_{\Gamma}^F(\psi)} \} \\ \mathfrak{R}_{\Gamma}^F((j \boxplus_3 \psi)) \cdot e^{i\theta\mathfrak{S}_{\Gamma}^F((j \boxplus_3 \psi))} \leq \max \{ \mathfrak{R}_{\Gamma}^F(j) \cdot e^{i\theta\mathfrak{S}_{\Gamma}^F(j)}, \mathfrak{R}_{\Gamma}^F(\psi) \cdot e^{i\theta\mathfrak{S}_{\Gamma}^F(\psi)} \} \end{array} \right\}$$

for all  $j, \psi \in S$ .

**Example 3.6.** Consider  $S = \{\tau, v, \zeta, \eta\}$ :

|              |        |        |         |         |              |         |        |         |        |              |        |        |         |        |
|--------------|--------|--------|---------|---------|--------------|---------|--------|---------|--------|--------------|--------|--------|---------|--------|
| $\boxplus_1$ | $\tau$ | $v$    | $\zeta$ | $\eta$  | $\boxplus_2$ | $\tau$  | $v$    | $\zeta$ | $\eta$ | $\boxplus_3$ | $\tau$ | $v$    | $\zeta$ | $\eta$ |
| $\tau$       | $\tau$ | $\tau$ | $\tau$  | $\tau$  | $\tau$       | $\tau$  | $v$    | $\zeta$ | $\eta$ | $\tau$       | $\tau$ | $\tau$ | $\tau$  | $\tau$ |
| $v$          | $\tau$ | $v$    | $\tau$  | $v$     | $v$          | $v$     | $v$    | $\eta$  | $\eta$ | $v$          | $\tau$ | $v$    | $\zeta$ | $\eta$ |
| $\zeta$      | $\tau$ | $\tau$ | $\zeta$ | $\zeta$ | $\zeta$      | $\zeta$ | $\eta$ | $\zeta$ | $\eta$ | $\zeta$      | $\eta$ | $\eta$ | $\eta$  | $\eta$ |
| $\eta$       | $\tau$ | $v$    | $\zeta$ | $\eta$  | $\eta$       | $\eta$  | $\eta$ | $\eta$  | $\eta$ | $\eta$       | $\eta$ | $\eta$ | $\eta$  | $\eta$ |

|                                                                         |                                            |                                            |
|-------------------------------------------------------------------------|--------------------------------------------|--------------------------------------------|
|                                                                         | $(w) = \tau$                               | $(w) = v$                                  |
| $(\widehat{\mathfrak{R}}_\Gamma^T, \widehat{\mathfrak{S}}_\Gamma^T)(w)$ | $[0.8e^{i2\pi(0.65)}, 0.9e^{i2\pi(0.75)}]$ | $[0.7e^{i2\pi(0.55)}, 0.8e^{i2\pi(0.65)}]$ |
| $(\widehat{\mathfrak{R}}_\Gamma^I, \widehat{\mathfrak{S}}_\Gamma^I)(w)$ | $[0.5e^{i2\pi(0.35)}, 0.6e^{i2\pi(0.45)}]$ | $[0.4e^{i2\pi(0.25)}, 0.5e^{i2\pi(0.35)}]$ |
| $(\widehat{\mathfrak{R}}_\Gamma^F, \widehat{\mathfrak{S}}_\Gamma^F)(w)$ | $[0.6e^{i2\pi(0.45)}, 0.7e^{i2\pi(0.55)}]$ | $[0.7e^{i2\pi(0.55)}, 0.8e^{i2\pi(0.6)}]$  |

|                                                                         |                                            |                                            |
|-------------------------------------------------------------------------|--------------------------------------------|--------------------------------------------|
|                                                                         | $(w) = \zeta$                              | $(w) = \eta$                               |
| $(\widehat{\mathfrak{R}}_\Gamma^T, \widehat{\mathfrak{S}}_\Gamma^T)(w)$ | $[0.4e^{i2\pi(0.25)}, 0.5e^{i2\pi(0.35)}]$ | $[0.6e^{i2\pi(0.45)}, 0.7e^{i2\pi(0.55)}]$ |
| $(\widehat{\mathfrak{R}}_\Gamma^I, \widehat{\mathfrak{S}}_\Gamma^I)(w)$ | $[0.2e^{i2\pi(0.15)}, 0.3e^{i2\pi(0.25)}]$ | $[0.3e^{i2\pi(0.15)}, 0.4e^{i2\pi(0.25)}]$ |
| $(\widehat{\mathfrak{R}}_\Gamma^F, \widehat{\mathfrak{S}}_\Gamma^F)(w)$ | $[0.9e^{i2\pi(0.75)}, 0.8e^{i2\pi(0.65)}]$ | $[0.85e^{i2\pi(0.7)}, 0.9e^{i2\pi(0.75)}]$ |

|                                                                         |                      |                      |
|-------------------------------------------------------------------------|----------------------|----------------------|
|                                                                         | $(w) = \tau$         | $(w) = v$            |
| $(\widehat{\mathfrak{R}}_\Gamma^T, \widehat{\mathfrak{S}}_\Gamma^T)(w)$ | $0.7e^{i2\pi(0.55)}$ | $0.6e^{i2\pi(0.45)}$ |
| $(\widehat{\mathfrak{R}}_\Gamma^I, \widehat{\mathfrak{S}}_\Gamma^I)(w)$ | $0.4e^{i2\pi(0.25)}$ | $0.3e^{i2\pi(0.15)}$ |
| $(\widehat{\mathfrak{R}}_\Gamma^F, \widehat{\mathfrak{S}}_\Gamma^F)(w)$ | $0.5e^{i2\pi(0.35)}$ | $0.6e^{i2\pi(0.45)}$ |

|                                                                         |                      |                      |
|-------------------------------------------------------------------------|----------------------|----------------------|
|                                                                         | $(w) = \zeta$        | $(w) = \eta$         |
| $(\widehat{\mathfrak{R}}_\Gamma^T, \widehat{\mathfrak{S}}_\Gamma^T)(w)$ | $0.3e^{i2\pi(0.15)}$ | $0.5e^{i2\pi(0.35)}$ |
| $(\widehat{\mathfrak{R}}_\Gamma^I, \widehat{\mathfrak{S}}_\Gamma^I)(w)$ | $0.1e^{i2\pi(0.05)}$ | $0.2e^{i2\pi(0.05)}$ |
| $(\widehat{\mathfrak{R}}_\Gamma^F, \widehat{\mathfrak{S}}_\Gamma^F)(w)$ | $0.8e^{i2\pi(0.65)}$ | $0.75e^{i2\pi(0.6)}$ |

Hence,  $\Gamma$  is a ComCNSBS of  $S$  as a result.

**Theorem 3.7.** Every ComCNSBS's intersection results in another ComCNSBS of  $S$ .

**Proof.** Let  $\{\widehat{\Upsilon}_i : i \in I\}$  be the collection of ComCNSBSs and  $\Gamma = \bigcap_{i \in I} \widehat{\Upsilon}_i$ . Let  $j, \psi \in S$ .

Now,

$$\begin{aligned}
 \widehat{\mathfrak{R}}_\Gamma^T((j \boxplus_1 \psi)) \cdot e^{i\theta \widehat{\mathfrak{S}}_\Gamma^T((j \boxplus_1 \psi))} &= \inf_{i \in I} \widehat{\mathfrak{R}}_{\Upsilon_i}^T((j \boxplus_1 \psi)) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Upsilon_i}^T((j \boxplus_1 \psi))} \\
 &\geq \inf_{i \in I} \min\{\widehat{\mathfrak{R}}_{\Upsilon_i}^T(j) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Upsilon_i}^T(j)}, \widehat{\mathfrak{R}}_{\Upsilon_i}^T(\psi) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Upsilon_i}^T(\psi)}\} \\
 &= \min\left\{\inf_{i \in I} \widehat{\mathfrak{R}}_{\Upsilon_i}^T(j) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Upsilon_i}^T(j)}, \inf_{i \in I} \widehat{\mathfrak{R}}_{\Upsilon_i}^T(\psi) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Upsilon_i}^T(\psi)}\right\} \\
 &= \min\{\widehat{\mathfrak{R}}_\Gamma^T(j) \cdot e^{i\theta \widehat{\mathfrak{S}}_\Gamma^T(j)}, \widehat{\mathfrak{R}}_\Gamma^T(\psi) \cdot e^{i\theta \widehat{\mathfrak{S}}_\Gamma^T(\psi)}\}
 \end{aligned}$$

Similarly,

$$\begin{aligned}
 \widehat{\mathfrak{R}}_\Gamma^T((j \boxplus_2 \psi)) \cdot e^{i\theta \widehat{\mathfrak{S}}_\Gamma^T((j \boxplus_2 \psi))} &\geq \min\{\widehat{\mathfrak{R}}_\Gamma^T(j) \cdot e^{i\theta \widehat{\mathfrak{S}}_\Gamma^T(j)}, \widehat{\mathfrak{R}}_\Gamma^T(\psi) \cdot e^{i\theta \widehat{\mathfrak{S}}_\Gamma^T(\psi)}\}, \\
 \widehat{\mathfrak{R}}_\Gamma^T((j \boxplus_3 \psi)) \cdot e^{i\theta \widehat{\mathfrak{S}}_\Gamma^T((j \boxplus_3 \psi))} &\geq \min\{\widehat{\mathfrak{R}}_\Gamma^T(j) \cdot e^{i\theta \widehat{\mathfrak{S}}_\Gamma^T(j)}, \widehat{\mathfrak{R}}_\Gamma^T(\psi) \cdot e^{i\theta \widehat{\mathfrak{S}}_\Gamma^T(\psi)}\}.
 \end{aligned}$$

Now,

$$\begin{aligned}
\widehat{\mathfrak{R}}_I^I((j \boxplus_1 \psi)) \cdot e^{i\theta \widehat{\mathfrak{S}}_I^I((j \boxplus_1 \psi))} &= \inf_{i \in I} \widehat{\mathfrak{R}}_{\Upsilon_i}^I((j \boxplus_1 \psi)) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Upsilon_i}^I((j \boxplus_1 \psi))} \\
&\geq \inf_{i \in I} \frac{\widehat{\mathfrak{R}}_{\Upsilon_i}^I(j) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Upsilon_i}^I(j)} + \widehat{\mathfrak{R}}_{\Upsilon_i}^I(\psi) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Upsilon_i}^I(\psi)}}{2} \\
&= \frac{\inf_{i \in I} \widehat{\mathfrak{R}}_{\Upsilon_i}^I(j) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Upsilon_i}^I(j)} + \inf_{i \in I} \widehat{\mathfrak{R}}_{\Upsilon_i}^I(\psi) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Upsilon_i}^I(\psi)}}{2} \\
&= \frac{\widehat{\mathfrak{R}}_I^I(j) \cdot e^{i\theta \widehat{\mathfrak{S}}_I^I(j)} + \widehat{\mathfrak{R}}_I^I(\psi) \cdot e^{i\theta \widehat{\mathfrak{S}}_I^I(\psi)}}{2}
\end{aligned}$$

Similarly,

$$\begin{aligned}
\widehat{\mathfrak{R}}_I^I((j \boxplus_2 \psi)) \cdot e^{i\theta \widehat{\mathfrak{S}}_I^I((j \boxplus_2 \psi))} &\geq \frac{\widehat{\mathfrak{R}}_I^I(j) \cdot e^{i\theta \widehat{\mathfrak{S}}_I^I(j)} + \widehat{\mathfrak{R}}_I^I(\psi) \cdot e^{i\theta \widehat{\mathfrak{S}}_I^I(\psi)}}{2} \text{ and} \\
\widehat{\mathfrak{R}}_I^I((j \boxplus_3 \psi)) \cdot e^{i\theta \widehat{\mathfrak{S}}_I^I((j \boxplus_3 \psi))} &\geq \frac{\widehat{\mathfrak{R}}_I^I(j) \cdot e^{i\theta \widehat{\mathfrak{S}}_I^I(j)} + \widehat{\mathfrak{R}}_I^I(\psi) \cdot e^{i\theta \widehat{\mathfrak{S}}_I^I(\psi)}}{2}.
\end{aligned}$$

Now,

$$\begin{aligned}
\widehat{\mathfrak{R}}_I^F((j \boxplus_1 \psi)) \cdot e^{i\theta \widehat{\mathfrak{S}}_I^F((j \boxplus_1 \psi))} &= \sup_{i \in I} \widehat{\mathfrak{R}}_{\Upsilon_i}^F((j \boxplus_1 \psi)) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Upsilon_i}^F((j \boxplus_1 \psi))} \\
&\leq \sup_{i \in I} \max\{\widehat{\mathfrak{R}}_{\Upsilon_i}^F(j) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Upsilon_i}^F(j)}, \widehat{\mathfrak{R}}_{\Upsilon_i}^F(\psi) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Upsilon_i}^F(\psi)}\} \\
&= \max\left\{\sup_{i \in I} \widehat{\mathfrak{R}}_{\Upsilon_i}^F(j) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Upsilon_i}^F(j)}, \sup_{i \in I} \widehat{\mathfrak{R}}_{\Upsilon_i}^F(\psi) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Upsilon_i}^F(\psi)}\right\} \\
&= \max\{\widehat{\mathfrak{R}}_I^F(j) \cdot e^{i\theta \widehat{\mathfrak{S}}_I^F(j)}, \widehat{\mathfrak{R}}_I^F(\psi) \cdot e^{i\theta \widehat{\mathfrak{S}}_I^F(\psi)}\}
\end{aligned}$$

Similarly,

$$\begin{aligned}
\widehat{\mathfrak{R}}_I^F((j \boxplus_2 \psi)) \cdot e^{i\theta \widehat{\mathfrak{S}}_I^F((j \boxplus_2 \psi))} &\leq \max\{\widehat{\mathfrak{R}}_I^F(j) \cdot e^{i\theta \widehat{\mathfrak{S}}_I^F(j)}, \widehat{\mathfrak{R}}_I^F(\psi) \cdot e^{i\theta \widehat{\mathfrak{S}}_I^F(\psi)}\} \text{ and} \\
\widehat{\mathfrak{R}}_I^F((j \boxplus_3 \psi)) \cdot e^{i\theta \widehat{\mathfrak{S}}_I^F((j \boxplus_3 \psi))} &\leq \max\{\widehat{\mathfrak{R}}_I^F(j) \cdot e^{i\theta \widehat{\mathfrak{S}}_I^F(j)}, \widehat{\mathfrak{R}}_I^F(\psi) \cdot e^{i\theta \widehat{\mathfrak{S}}_I^F(\psi)}\}.
\end{aligned}$$

Let  $\{\Upsilon_i : i \in I\}$  be the collection of ComCNSBSs of  $S$  and  $\Gamma = \bigcap_{i \in I} \Upsilon_i$ . Let  $j, \psi \in S$ .

Now,

$$\begin{aligned}
\mathfrak{R}_I^T((j \boxplus_1 \psi)) \cdot e^{i\theta \mathfrak{S}_I^T((j \boxplus_1 \psi))} &= \inf_{i \in I} \mathfrak{R}_{\Upsilon_i}^T((j \boxplus_1 \psi)) \cdot e^{i\theta \mathfrak{S}_{\Upsilon_i}^T((j \boxplus_1 \psi))} \\
&\geq \inf_{i \in I} \min\{\mathfrak{R}_{\Upsilon_i}^T(j) \cdot e^{i\theta \mathfrak{S}_{\Upsilon_i}^T(j)}, \mathfrak{R}_{\Upsilon_i}^T(\psi) \cdot e^{i\theta \mathfrak{S}_{\Upsilon_i}^T(\psi)}\} \\
&= \min\left\{\inf_{i \in I} \mathfrak{R}_{\Upsilon_i}^T(j) \cdot e^{i\theta \mathfrak{S}_{\Upsilon_i}^T(j)}, \inf_{i \in I} \mathfrak{R}_{\Upsilon_i}^T(\psi) \cdot e^{i\theta \mathfrak{S}_{\Upsilon_i}^T(\psi)}\right\} \\
&= \min\{\mathfrak{R}_I^T(j) \cdot e^{i\theta \mathfrak{S}_I^T(j)}, \mathfrak{R}_I^T(\psi) \cdot e^{i\theta \mathfrak{S}_I^T(\psi)}\}
\end{aligned}$$

Similarly,

$$\begin{aligned}
\mathfrak{R}_I^T((j \boxplus_2 \psi)) \cdot e^{i\theta \mathfrak{S}_I^T((j \boxplus_2 \psi))} &\geq \min\{\mathfrak{R}_I^T(j) \cdot e^{i\theta \mathfrak{S}_I^T(j)}, \mathfrak{R}_I^T(\psi) \cdot e^{i\theta \mathfrak{S}_I^T(\psi)}\}, \\
\mathfrak{R}_I^T((j \boxplus_3 \psi)) \cdot e^{i\theta \mathfrak{S}_I^T((j \boxplus_3 \psi))} &\geq \min\{\mathfrak{R}_I^T(j) \cdot e^{i\theta \mathfrak{S}_I^T(j)}, \mathfrak{R}_I^T(\psi) \cdot e^{i\theta \mathfrak{S}_I^T(\psi)}\}.
\end{aligned}$$

Now,

$$\begin{aligned}
\mathfrak{R}_I^I((j \boxplus_1 \psi)) \cdot e^{i\theta \mathfrak{S}_I^I((j \boxplus_1 \psi))} &= \inf_{i \in I} \mathfrak{R}_{\Upsilon_i}^I((j \boxplus_1 \psi)) \cdot e^{i\theta \mathfrak{S}_{\Upsilon_i}^I((j \boxplus_1 \psi))} \\
&\geq \inf_{i \in I} \frac{\mathfrak{R}_{\Upsilon_i}^I(j) \cdot e^{i\theta \mathfrak{S}_{\Upsilon_i}^I(j)} + \mathfrak{R}_{\Upsilon_i}^I(\psi) \cdot e^{i\theta \mathfrak{S}_{\Upsilon_i}^I(\psi)}}{2} \\
&= \frac{\inf_{i \in I} \mathfrak{R}_{\Upsilon_i}^I(j) \cdot e^{i\theta \mathfrak{S}_{\Upsilon_i}^I(j)} + \inf_{i \in I} \mathfrak{R}_{\Upsilon_i}^I(\psi) \cdot e^{i\theta \mathfrak{S}_{\Upsilon_i}^I(\psi)}}{2} \\
&= \frac{\mathfrak{R}_I^I(j) \cdot e^{i\theta \mathfrak{S}_I^I(j)} + \mathfrak{R}_I^I(\psi) \cdot e^{i\theta \mathfrak{S}_I^I(\psi)}}{2}
\end{aligned}$$

Similarly,

$$\mathfrak{R}_I^I((j \boxplus_2 \psi)) \cdot e^{i\theta \mathfrak{S}_I^I((j \boxplus_2 \psi))} \geq \frac{\mathfrak{R}_I^I(j) \cdot e^{i\theta \mathfrak{S}_I^I(j)} + \mathfrak{R}_I^I(\psi) \cdot e^{i\theta \mathfrak{S}_I^I(\psi)}}{2} \text{ and}$$

$$\mathfrak{R}_\Gamma^I((j \boxplus_3 \psi)) \cdot e^{i\theta \mathfrak{S}_\Gamma^I((j \boxplus_3 \psi))} \geq \frac{\mathfrak{R}_\Gamma^I(j) \cdot e^{i\theta \mathfrak{S}_\Gamma^I(j)} + \mathfrak{R}_\Gamma^I(\psi) \cdot e^{i\theta \mathfrak{S}_\Gamma^I(\psi)}}{2}.$$

Now,

$$\begin{aligned} \mathfrak{R}_\Gamma^F((j \boxplus_1 \psi)) \cdot e^{i\theta \mathfrak{S}_\Gamma^F((j \boxplus_1 \psi))} &= \sup_{i \in I} \mathfrak{R}_{\Gamma_i}^F((j \boxplus_1 \psi)) \cdot e^{i\theta \mathfrak{S}_{\Gamma_i}^F((j \boxplus_1 \psi))} \\ &\leq \sup_{i \in I} \max\{\mathfrak{R}_{\Gamma_i}^F(j) \cdot e^{i\theta \mathfrak{S}_{\Gamma_i}^F(j)}, \mathfrak{R}_{\Gamma_i}^F(\psi) \cdot e^{i\theta \mathfrak{S}_{\Gamma_i}^F(\psi)}\} \\ &= \max\left\{\sup_{i \in I} \mathfrak{R}_{\Gamma_i}^F(j) \cdot e^{i\theta \mathfrak{S}_{\Gamma_i}^F(j)}, \sup_{i \in I} \mathfrak{R}_{\Gamma_i}^F(\psi) \cdot e^{i\theta \mathfrak{S}_{\Gamma_i}^F(\psi)}\right\} \\ &= \max\{\mathfrak{R}_\Gamma^F(j) \cdot e^{i\theta \mathfrak{S}_\Gamma^F(j)}, \mathfrak{R}_\Gamma^F(\psi) \cdot e^{i\theta \mathfrak{S}_\Gamma^F(\psi)}\} \end{aligned}$$

Similarly,

$$\mathfrak{R}_\Gamma^F((j \boxplus_2 \psi)) \cdot e^{i\theta \mathfrak{S}_\Gamma^F((j \boxplus_2 \psi))} \leq \max\{\mathfrak{R}_\Gamma^F(j) \cdot e^{i\theta \mathfrak{S}_\Gamma^F(j)}, \mathfrak{R}_\Gamma^F(\psi) \cdot e^{i\theta \mathfrak{S}_\Gamma^F(\psi)}\} \text{ and}$$

$$\mathfrak{R}_\Gamma^F((j \boxplus_3 \psi)) \cdot e^{i\theta \mathfrak{S}_\Gamma^F((j \boxplus_3 \psi))} \leq \max\{\mathfrak{R}_\Gamma^F(j) \cdot e^{i\theta \mathfrak{S}_\Gamma^F(j)}, \mathfrak{R}_\Gamma^F(\psi) \cdot e^{i\theta \mathfrak{S}_\Gamma^F(\psi)}\}.$$

Thus,  $\Gamma$  is a ComCNSBS of  $S$ .

**Theorem 3.8.** If  $\Gamma$  and  $\Lambda$  be the ComCNSBSs of  $S_1$  and  $S_2$  respectively, then  $\widehat{\Gamma \times \Lambda}$  is a ComCNSBS of  $S_1 \times S_2$ .

**Proof.** Let  $j_1, j_2 \in S_1$  and  $\psi_1, \psi_2 \in S_2$ . Then  $(j_1, \psi_1)$  and  $(j_2, \psi_2)$  are in  $S_1 \times S_2$ . Now

$$\begin{aligned} &\widehat{\mathfrak{R}_{\Gamma \times \Lambda}^T}([(j_1, \psi_1) \boxplus_1 (j_2, \psi_2)]) \cdot e^{i\theta \widehat{\mathfrak{S}_{\Gamma \times \Lambda}^T}([(j_1, \psi_1) \boxplus_1 (j_2, \psi_2)])} \\ &= \widehat{\mathfrak{R}_{\Gamma \times \Lambda}^T}((j_1 \boxplus_1 j_2, \psi_1 \boxplus_1 \psi_2)) \cdot e^{i\theta \widehat{\mathfrak{S}_{\Gamma \times \Lambda}^T}((j_1 \boxplus_1 j_2, \psi_1 \boxplus_1 \psi_2))} \\ &= \min\{\widehat{\mathfrak{R}_\Gamma^T}((j_1 \boxplus_1 j_2)) \cdot e^{i\theta \widehat{\mathfrak{S}_\Gamma^T}((j_1 \boxplus_1 j_2))}, \widehat{\mathfrak{R}_\Lambda^T}((\psi_1 \boxplus_1 \psi_2)) \cdot e^{i\theta \widehat{\mathfrak{S}_\Lambda^T}((\psi_1 \boxplus_1 \psi_2))}\} \\ &\geq \min\{\min\{\widehat{\mathfrak{R}_\Gamma^T}(j_1) \cdot e^{i\theta \widehat{\mathfrak{S}_\Gamma^T}(j_1)}, \widehat{\mathfrak{R}_\Gamma^T}(j_2) \cdot e^{i\theta \widehat{\mathfrak{S}_\Gamma^T}(j_2)}\}, \min\{\widehat{\mathfrak{R}_\Lambda^T}(\psi_1) \cdot e^{i\theta \widehat{\mathfrak{S}_\Lambda^T}(\psi_1)}, \widehat{\mathfrak{R}_\Lambda^T}(\psi_2) \cdot e^{i\theta \widehat{\mathfrak{S}_\Lambda^T}(\psi_2)}\}\} \\ &= \min\{\min\{\widehat{\mathfrak{R}_\Gamma^T}(j_1) \cdot e^{i\theta \widehat{\mathfrak{S}_\Gamma^T}(j_1)}, \widehat{\mathfrak{R}_\Lambda^T}(\psi_1) \cdot e^{i\theta \widehat{\mathfrak{S}_\Lambda^T}(\psi_1)}\}, \min\{\widehat{\mathfrak{R}_\Gamma^T}(j_2) \cdot e^{i\theta \widehat{\mathfrak{S}_\Gamma^T}(j_2)}, \widehat{\mathfrak{R}_\Lambda^T}(\psi_2) \cdot e^{i\theta \widehat{\mathfrak{S}_\Lambda^T}(\psi_2)}\}\} \\ &= \min\{\widehat{\mathfrak{R}_{\Gamma \times \Lambda}^T}((j_1, \psi_1)) \cdot e^{i\theta \widehat{\mathfrak{S}_{\Gamma \times \Lambda}^T}((j_1, \psi_1))}, \widehat{\mathfrak{R}_{\Gamma \times \Lambda}^T}((j_2, \psi_2)) \cdot e^{i\theta \widehat{\mathfrak{S}_{\Gamma \times \Lambda}^T}((j_2, \psi_2))}\} \end{aligned}$$

$$\begin{aligned} &\text{Also } \widehat{\mathfrak{R}_{\Gamma \times \Lambda}^T}([(j_1, \psi_1) \boxplus_2 (j_2, \psi_2)]) \cdot e^{i\theta \widehat{\mathfrak{S}_{\Gamma \times \Lambda}^T}([(j_1, \psi_1) \boxplus_2 (j_2, \psi_2)])} \\ &\geq \min\{\widehat{\mathfrak{R}_{\Gamma \times \Lambda}^T}((j_1, \psi_1)) \cdot e^{i\theta \widehat{\mathfrak{S}_{\Gamma \times \Lambda}^T}((j_1, \psi_1))}, \widehat{\mathfrak{R}_{\Gamma \times \Lambda}^T}((j_2, \psi_2)) \cdot e^{i\theta \widehat{\mathfrak{S}_{\Gamma \times \Lambda}^T}((j_2, \psi_2))}\} \end{aligned}$$

$$\begin{aligned} &\text{and } \widehat{\mathfrak{R}_{\Gamma \times \Lambda}^T}([(j_1, \psi_1) \boxplus_3 (j_2, \psi_2)]) \cdot e^{i\theta \widehat{\mathfrak{S}_{\Gamma \times \Lambda}^T}([(j_1, \psi_1) \boxplus_3 (j_2, \psi_2)])} \\ &\geq \min\{\widehat{\mathfrak{R}_{\Gamma \times \Lambda}^T}((j_1, \psi_1)) \cdot e^{i\theta \widehat{\mathfrak{S}_{\Gamma \times \Lambda}^T}((j_1, \psi_1))}, \widehat{\mathfrak{R}_{\Gamma \times \Lambda}^T}((j_2, \psi_2)) \cdot e^{i\theta \widehat{\mathfrak{S}_{\Gamma \times \Lambda}^T}((j_2, \psi_2))}\}. \end{aligned}$$

Now,

$$\begin{aligned} &\widehat{\mathfrak{R}_{\Gamma \times \Lambda}^I}([(j_1, \psi_1) \boxplus_1 (j_2, \psi_2)]) \cdot e^{i\theta \widehat{\mathfrak{S}_{\Gamma \times \Lambda}^I}([(j_1, \psi_1) \boxplus_1 (j_2, \psi_2)])} \\ &= \widehat{\mathfrak{R}_{\Gamma \times \Lambda}^I}((j_1 \boxplus_1 j_2, \psi_1 \boxplus_1 \psi_2)) \cdot e^{i\theta \widehat{\mathfrak{S}_{\Gamma \times \Lambda}^I}((j_1 \boxplus_1 j_2, \psi_1 \boxplus_1 \psi_2))} \\ &= \frac{\widehat{\mathfrak{R}_\Gamma^I}((j_1 \boxplus_1 j_2)) \cdot e^{i\theta \widehat{\mathfrak{S}_{\Gamma \times \Lambda}^I}((j_1 \boxplus_1 j_2))} + \widehat{\mathfrak{R}_\Lambda^I}((\psi_1 \boxplus_1 \psi_2)) \cdot e^{i\theta \widehat{\mathfrak{S}_{\Gamma \times \Lambda}^I}((\psi_1 \boxplus_1 \psi_2))}}{2} \\ &\geq \frac{1}{2} \left[ \frac{\widehat{\mathfrak{R}_\Gamma^I}(j_1) \cdot e^{i\theta \widehat{\mathfrak{S}_\Gamma^I}(j_1)} + \widehat{\mathfrak{R}_\Gamma^I}(j_2) \cdot e^{i\theta \widehat{\mathfrak{S}_\Gamma^I}(j_2)}}{2} + \frac{\widehat{\mathfrak{R}_\Lambda^I}(\psi_1) \cdot e^{i\theta \widehat{\mathfrak{S}_\Lambda^I}(\psi_1)} + \widehat{\mathfrak{R}_\Lambda^I}(\psi_2) \cdot e^{i\theta \widehat{\mathfrak{S}_\Lambda^I}(\psi_2)}}{2} \right] \\ &= \frac{1}{2} \left[ \frac{\widehat{\mathfrak{R}_\Gamma^I}(j_1) \cdot e^{i\theta \widehat{\mathfrak{S}_\Gamma^I}(j_1)} + \widehat{\mathfrak{R}_\Lambda^I}(\psi_1) \cdot e^{i\theta \widehat{\mathfrak{S}_\Lambda^I}(\psi_1)}}{2} + \frac{\widehat{\mathfrak{R}_\Gamma^I}(j_2) \cdot e^{i\theta \widehat{\mathfrak{S}_\Gamma^I}(j_2)} + \widehat{\mathfrak{R}_\Lambda^I}(\psi_2) \cdot e^{i\theta \widehat{\mathfrak{S}_\Lambda^I}(\psi_2)}}{2} \right] \\ &= \frac{1}{2} \left[ \widehat{\mathfrak{R}_{\Gamma \times \Lambda}^I}((j_1, \psi_1)) \cdot e^{i\theta \widehat{\mathfrak{S}_{\Gamma \times \Lambda}^I}((j_1, \psi_1))} + \widehat{\mathfrak{R}_{\Gamma \times \Lambda}^I}((j_2, \psi_2)) \cdot e^{i\theta \widehat{\mathfrak{S}_{\Gamma \times \Lambda}^I}((j_2, \psi_2))} \right] \end{aligned}$$

Also

$$\widehat{\mathfrak{R}_{\Gamma \times \Lambda}^I}([(j_1, \psi_1) \boxplus_2 (j_2, \psi_2)]) \cdot e^{i\theta \widehat{\mathfrak{S}_{\Gamma \times \Lambda}^I}([(j_1, \psi_1) \boxplus_2 (j_2, \psi_2)])} \geq \frac{1}{2} \left[ \widehat{\mathfrak{R}_{\Gamma \times \Lambda}^I}((j_1, \psi_1)) \cdot e^{i\theta \widehat{\mathfrak{S}_{\Gamma \times \Lambda}^I}((j_1, \psi_1))} + \widehat{\mathfrak{R}_{\Gamma \times \Lambda}^I}((j_2, \psi_2)) \cdot e^{i\theta \widehat{\mathfrak{S}_{\Gamma \times \Lambda}^I}((j_2, \psi_2))} \right].$$





Also

$$\mathfrak{R}_{\Gamma \times \Lambda}^I[((j_1, \psi_1) \boxplus_2 (j_2, \psi_2))] \cdot e^{i\theta \mathfrak{S}_{\Gamma \times \Lambda}^I[((j_1, \psi_1) \boxplus_2 (j_2, \psi_2))]} \geq \frac{1}{2} \left[ \mathfrak{R}_{\Gamma \times \Lambda}^I((j_1, \psi_1)) \cdot e^{i\theta \mathfrak{S}_{\Gamma \times \Lambda}^I((j_1, \psi_1))} + \mathfrak{R}_{\Gamma \times \Lambda}^I((j_2, \psi_2)) \cdot e^{i\theta \mathfrak{S}_{\Gamma \times \Lambda}^I((j_2, \psi_2))} \right]$$

and

$$\mathfrak{R}_{\Gamma \times \Lambda}^I[((j_1, \psi_1) \boxplus_3 (j_2, \psi_2))] \cdot e^{i\theta \mathfrak{S}_{\Gamma \times \Lambda}^I[((j_1, \psi_1) \boxplus_3 (j_2, \psi_2))]} \geq \frac{1}{2} \left[ \mathfrak{R}_{\Gamma \times \Lambda}^I((j_1, \psi_1)) \cdot e^{i\theta \mathfrak{S}_{\Gamma \times \Lambda}^I((j_1, \psi_1))} + \mathfrak{R}_{\Gamma \times \Lambda}^I((j_2, \psi_2)) \cdot e^{i\theta \mathfrak{S}_{\Gamma \times \Lambda}^I((j_2, \psi_2))} \right].$$

Now,

$$\begin{aligned} & \mathfrak{R}_{\Gamma \times \Lambda}^F(((j_1, \psi_1) \boxplus_1 (j_2, \psi_2))) \cdot e^{i\theta \mathfrak{S}_{\Gamma \times \Lambda}^F(((j_1, \psi_1) \boxplus_1 (j_2, \psi_2)))} \\ &= \mathfrak{R}_{\Gamma \times \Lambda}^F((j_1 \boxplus_1 j_2, \psi_1 \boxplus_1 \psi_2)) \cdot e^{i\theta \mathfrak{S}_{\Gamma \times \Lambda}^F((j_1 \boxplus_1 j_2, \psi_1 \boxplus_1 \psi_2))} \\ &= \max\{\mathfrak{R}_{\Gamma}^F((j_1 \boxplus_1 j_2)) \cdot e^{i\theta \mathfrak{S}_{\Gamma \times \Lambda}^F((j_1 \boxplus_1 j_2))}, \mathfrak{R}_{\Lambda}^F((\psi_1 \boxplus_1 \psi_2)) \cdot e^{i\theta \mathfrak{S}_{\Gamma \times \Lambda}^F((\psi_1 \boxplus_1 \psi_2))}\} \\ &\leq \max\{\max\{\mathfrak{R}_{\Gamma}^F(j_1) \cdot e^{i\theta \mathfrak{S}_{\Gamma}^F(j_1)}, \mathfrak{R}_{\Gamma}^F(j_2) \cdot e^{i\theta \mathfrak{S}_{\Gamma}^F(j_2)}\}, \max\{\mathfrak{R}_{\Lambda}^F(\psi_1) \cdot e^{i\theta \mathfrak{S}_{\Lambda}^F(\psi_1)}, \mathfrak{R}_{\Lambda}^F(\psi_2) \cdot e^{i\theta \mathfrak{S}_{\Lambda}^F(\psi_2)}\}\} \\ &= \max\{\max\{\mathfrak{R}_{\Gamma}^F(j_1) \cdot e^{i\theta \mathfrak{S}_{\Gamma}^F(j_1)}, \mathfrak{R}_{\Lambda}^F(\psi_1) \cdot e^{i\theta \mathfrak{S}_{\Lambda}^F(\psi_1)}\}, \max\{\mathfrak{R}_{\Gamma}^F(j_2) \cdot e^{i\theta \mathfrak{S}_{\Gamma}^F(j_2)}, \mathfrak{R}_{\Lambda}^F(\psi_2) \cdot e^{i\theta \mathfrak{S}_{\Lambda}^F(\psi_2)}\}\} \\ &= \max\{\mathfrak{R}_{\Gamma \times \Lambda}^F((j_1, \psi_1)) \cdot e^{i\theta \mathfrak{S}_{\Gamma \times \Lambda}^F((j_1, \psi_1))}, \mathfrak{R}_{\Gamma \times \Lambda}^F((j_2, \psi_2)) \cdot e^{i\theta \mathfrak{S}_{\Gamma \times \Lambda}^F((j_2, \psi_2))}\} \end{aligned}$$

$$\begin{aligned} & \text{Also } \mathfrak{R}_{\Gamma \times \Lambda}^F(((j_1, \psi_1) \boxplus_2 (j_2, \psi_2))) \cdot e^{i\theta \mathfrak{S}_{\Gamma \times \Lambda}^F(((j_1, \psi_1) \boxplus_2 (j_2, \psi_2)))} \\ &\leq \max\{\mathfrak{R}_{\Gamma \times \Lambda}^F((j_1, \psi_1)) \cdot e^{i\theta \mathfrak{S}_{\Gamma \times \Lambda}^F((j_1, \psi_1))}, \mathfrak{R}_{\Gamma \times \Lambda}^F((j_2, \psi_2)) \cdot e^{i\theta \mathfrak{S}_{\Gamma \times \Lambda}^F((j_2, \psi_2))}\}, \\ &\mathfrak{R}_{\Gamma \times \Lambda}^F(((j_1, \psi_1) \boxplus_3 (j_2, \psi_2))) \cdot e^{i\theta \mathfrak{S}_{\Gamma \times \Lambda}^F(((j_1, \psi_1) \boxplus_3 (j_2, \psi_2)))} \\ &\leq \max\{\mathfrak{R}_{\Gamma \times \Lambda}^F((j_1, \psi_1)) \cdot e^{i\theta \mathfrak{S}_{\Gamma \times \Lambda}^F((j_1, \psi_1))}, \mathfrak{R}_{\Gamma \times \Lambda}^F((j_2, \psi_2)) \cdot e^{i\theta \mathfrak{S}_{\Gamma \times \Lambda}^F((j_2, \psi_2))}\}. \end{aligned}$$

Thus,  $\widehat{\Gamma \times \Lambda}$  is a ComCNSBS.

**Definition 3.9.** The strongest ComCN connection on  $S$ , given  $\Gamma \subseteq S$  is

$$\left\{ \begin{aligned} \widehat{\mathfrak{R}}_{\Gamma}^T((j, \psi)) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Gamma}^T((j, \psi))} &= \min\{\widehat{\mathfrak{R}}_{\Gamma}^T(j) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Gamma}^T(j)}, \widehat{\mathfrak{R}}_{\Gamma}^T(\psi) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Gamma}^T(\psi)}\} \\ \widehat{\mathfrak{R}}_{\Gamma}^I((j, \psi)) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Gamma}^I((j, \psi))} &= \frac{\widehat{\mathfrak{R}}_{\Gamma}^I(j) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Gamma}^I(j)} + \widehat{\mathfrak{R}}_{\Gamma}^I(\psi) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Gamma}^I(\psi)}}{2} \\ \widehat{\mathfrak{R}}_{\Gamma}^F((j, \psi)) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Gamma}^F((j, \psi))} &= \max\{\widehat{\mathfrak{R}}_{\Gamma}^F(j) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Gamma}^F(j)}, \widehat{\mathfrak{R}}_{\Gamma}^F(\psi) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Gamma}^F(\psi)}\} \end{aligned} \right\}$$

**Theorem 3.10.** Let  $\Upsilon$  be the strongest complex cubic neutrosophic relation of  $S$ , and let  $\Gamma$  be a ComCNSBS of  $S$ . If and only if  $\Upsilon$  is a ComCNSBS of  $S \times S$ , then  $\Gamma$  is a ComCNSBS of  $S \times S$ .

**Proof.** Let  $\widehat{\Upsilon}$  be the strongest complex cubic neutrosophic relation of  $S$ , and suppose  $\Gamma$  is a ComCNSBS of  $S \times S$ .

For  $j = (j_1, j_2), \psi = (\psi_1, \psi_2) \in S \times S$ . Now,

$$\begin{aligned} & \widehat{\mathfrak{R}}_{\Gamma}^T((j \boxplus_1 \psi)) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Gamma}^T((j \boxplus_1 \psi))} \\ &= \widehat{\mathfrak{R}}_{\Gamma}^T((((j_1, j_2) \boxplus_1 (\psi_1, \psi_2)))) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Gamma}^T((((j_1, j_2) \boxplus_1 (\psi_1, \psi_2))))} \\ &= \widehat{\mathfrak{R}}_{\Gamma}^T(j_1 \boxplus_1 \psi_1, j_2 \boxplus_1 \psi_2) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Gamma}^T(j_1 \boxplus_1 \psi_1, j_2 \boxplus_1 \psi_2)} \\ &= \min\{\widehat{\mathfrak{R}}_{\Gamma}^T((j_1 \boxplus_1 \psi_1)) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Gamma}^T((j_1 \boxplus_1 \psi_1))}, \widehat{\mathfrak{R}}_{\Gamma}^T((j_2 \boxplus_1 \psi_2)) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Gamma}^T((j_2 \boxplus_1 \psi_2))}\} \\ &\geq \min\{\min\{\widehat{\mathfrak{R}}_{\Gamma}^T(j_1) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Gamma}^T(j_1)}, \widehat{\mathfrak{R}}_{\Gamma}^T(\psi_1) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Gamma}^T(\psi_1)}\}, \\ &\quad \min\{\widehat{\mathfrak{R}}_{\Gamma}^T(j_2) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Gamma}^T(j_2)}, \widehat{\mathfrak{R}}_{\Gamma}^T(\psi_2) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Gamma}^T(\psi_2)}\}\} \\ &= \min\{\min\{\widehat{\mathfrak{R}}_{\Gamma}^T(j_1) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Gamma}^T(j_1)}, \widehat{\mathfrak{R}}_{\Gamma}^T(j_2) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Gamma}^T(j_2)}\}, \\ &\quad \min\{\widehat{\mathfrak{R}}_{\Gamma}^T(\psi_1) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Gamma}^T(\psi_1)}, \widehat{\mathfrak{R}}_{\Gamma}^T(\psi_2) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Gamma}^T(\psi_2)}\}\} \\ &= \min\{\widehat{\mathfrak{R}}_{\Gamma}^T((j_1, j_2)) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Gamma}^T((j_1, j_2))}, \widehat{\mathfrak{R}}_{\Gamma}^T((\psi_1, \psi_2)) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Gamma}^T((\psi_1, \psi_2))}\} \\ &= \min\{\widehat{\mathfrak{R}}_{\Gamma}^T(j) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Gamma}^T(j)}, \widehat{\mathfrak{R}}_{\Gamma}^T(\psi) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Gamma}^T(\psi)}\} \end{aligned}$$

$$\begin{aligned} & \text{Also } \widehat{\mathfrak{R}}_{\Gamma}^T((j \boxplus_2 \psi)) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Gamma}^T((j \boxplus_2 \psi))} \geq \min\{\widehat{\mathfrak{R}}_{\Gamma}^T(j) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Gamma}^T(j)}, \widehat{\mathfrak{R}}_{\Gamma}^T(\psi) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Gamma}^T(\psi)}\}, \\ & \widehat{\mathfrak{R}}_{\Gamma}^T((j \boxplus_3 \psi)) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Gamma}^T((j \boxplus_3 \psi))} \geq \min\{\widehat{\mathfrak{R}}_{\Gamma}^T(j) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Gamma}^T(j)}, \widehat{\mathfrak{R}}_{\Gamma}^T(\psi) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Gamma}^T(\psi)}\}. \end{aligned}$$

Now,  $\widehat{\mathfrak{R}}_I^I(j \boxplus \psi) \cdot e^{i\theta\widehat{\mathfrak{S}}_I^I(j \boxplus \psi)}$

$$\begin{aligned}
 &= \widehat{\mathfrak{R}}_I^I[(((j_1, j_2)) \boxplus (\psi_1, \psi_2))] \cdot e^{i\theta\widehat{\mathfrak{S}}_I^I[(((j_1, j_2)) \boxplus (\psi_1, \psi_2))]} \\
 &= \widehat{\mathfrak{R}}_I^I((j_1 \boxplus \psi_1, j_2 \boxplus \psi_2)) \cdot e^{i\theta\widehat{\mathfrak{S}}_I^I((j_1 \boxplus \psi_1, j_2 \boxplus \psi_2))} \\
 &= \frac{\widehat{\mathfrak{R}}_I^I((j_1 \boxplus \psi_1)) \cdot e^{i\theta\widehat{\mathfrak{S}}_I^I((j_1 \boxplus \psi_1))} + \widehat{\mathfrak{R}}_I^I((j_2 \boxplus \psi_2)) \cdot e^{i\theta\widehat{\mathfrak{S}}_I^I((j_2 \boxplus \psi_2))}}{2} \\
 &\geq \frac{1}{2} \left[ \frac{\widehat{\mathfrak{R}}_I^I(j_1) \cdot e^{i\theta\widehat{\mathfrak{S}}_I^I(j_1)} + \widehat{\mathfrak{R}}_I^I(\psi_1) \cdot e^{i\theta\widehat{\mathfrak{S}}_I^I(\psi_1)}}{2} + \frac{\widehat{\mathfrak{R}}_I^I(j_2) \cdot e^{i\theta\widehat{\mathfrak{S}}_I^I(j_2)} + \widehat{\mathfrak{R}}_I^I(\psi_2) \cdot e^{i\theta\widehat{\mathfrak{S}}_I^I(\psi_2)}}{2} \right] \\
 &= \frac{1}{2} \left[ \frac{\widehat{\mathfrak{R}}_I^I(j_1) \cdot e^{i\theta\widehat{\mathfrak{S}}_I^I(j_1)} + \widehat{\mathfrak{R}}_I^I(j_2) \cdot e^{i\theta\widehat{\mathfrak{S}}_I^I(j_2)}}{2} + \frac{\widehat{\mathfrak{R}}_I^I(\psi_1) \cdot e^{i\theta\widehat{\mathfrak{S}}_I^I(\psi_1)} + \widehat{\mathfrak{R}}_I^I(\psi_2) \cdot e^{i\theta\widehat{\mathfrak{S}}_I^I(\psi_2)}}{2} \right] \\
 &= \frac{\widehat{\mathfrak{R}}_I^I((j_1, j_2)) \cdot e^{i\theta\widehat{\mathfrak{S}}_I^I((j_1, j_2))} + \widehat{\mathfrak{R}}_I^I((\psi_1, \psi_2)) \cdot e^{i\theta\widehat{\mathfrak{S}}_I^I((\psi_1, \psi_2))}}{2} \\
 &= \frac{\widehat{\mathfrak{R}}_I^I(j) \cdot e^{i\theta\widehat{\mathfrak{S}}_I^I(j)} + \widehat{\mathfrak{R}}_I^I(\psi) \cdot e^{i\theta\widehat{\mathfrak{S}}_I^I(\psi)}}{2}
 \end{aligned}$$

$$\text{Also } \widehat{\mathfrak{R}}_I^I((j \boxplus \psi)) \cdot e^{i\theta\widehat{\mathfrak{S}}_I^I((j \boxplus \psi))} \geq \frac{\widehat{\mathfrak{R}}_I^I(j) \cdot e^{i\theta\widehat{\mathfrak{S}}_I^I(j)} + \widehat{\mathfrak{R}}_I^I(\psi) \cdot e^{i\theta\widehat{\mathfrak{S}}_I^I(\psi)}}{2}$$

$$\text{and } \widehat{\mathfrak{R}}_I^I((j \boxplus \psi)) \cdot e^{i\theta\widehat{\mathfrak{S}}_I^I((j \boxplus \psi))} \geq \frac{\widehat{\mathfrak{R}}_I^I(j) \cdot e^{i\theta\widehat{\mathfrak{S}}_I^I(j)} + \widehat{\mathfrak{R}}_I^I(\psi) \cdot e^{i\theta\widehat{\mathfrak{S}}_I^I(\psi)}}{2}.$$

$$\text{Similarly, } \widehat{\mathfrak{R}}_I^F((j \boxplus \psi)) \cdot e^{i\theta\widehat{\mathfrak{S}}_I^F((j \boxplus \psi))} \leq \max\{\widehat{\mathfrak{R}}_I^F(j) \cdot e^{i\theta\widehat{\mathfrak{S}}_I^F(j)}, \widehat{\mathfrak{R}}_I^F(\psi) \cdot e^{i\theta\widehat{\mathfrak{S}}_I^F(\psi)}\},$$

$$\widehat{\mathfrak{R}}_I^F((j \boxplus \psi)) \cdot e^{i\theta\widehat{\mathfrak{S}}_I^F((j \boxplus \psi))} \leq \max\{\widehat{\mathfrak{R}}_I^F(j) \cdot e^{i\theta\widehat{\mathfrak{S}}_I^F(j)}, \widehat{\mathfrak{R}}_I^F(\psi) \cdot e^{i\theta\widehat{\mathfrak{S}}_I^F(\psi)}\} \text{ and}$$

$$\widehat{\mathfrak{R}}_I^F((j \boxplus \psi)) \cdot e^{i\theta\widehat{\mathfrak{S}}_I^F((j \boxplus \psi))} \leq \max\{\widehat{\mathfrak{R}}_I^F(j) \cdot e^{i\theta\widehat{\mathfrak{S}}_I^F(j)}, \widehat{\mathfrak{R}}_I^F(\psi) \cdot e^{i\theta\widehat{\mathfrak{S}}_I^F(\psi)}\}.$$

For any  $j = (j_1, j_2), \psi = (\psi_1, \psi_2) \in S \times S$ . Now,

$$\begin{aligned}
 &\mathfrak{R}_I^T((j \boxplus \psi)) \cdot e^{i\theta\widehat{\mathfrak{S}}_I^T((j \boxplus \psi))} \\
 &= \mathfrak{R}_I^T[(((j_1, j_2)) \boxplus (\psi_1, \psi_2))] \cdot e^{i\theta\widehat{\mathfrak{S}}_I^T[(((j_1, j_2)) \boxplus (\psi_1, \psi_2))]} \\
 &= \mathfrak{R}_I^T(j_1 \boxplus \psi_1, j_2 \boxplus \psi_2) \cdot e^{i\theta\widehat{\mathfrak{S}}_I^T(j_1 \boxplus \psi_1, j_2 \boxplus \psi_2)} \\
 &= \min\{\mathfrak{R}_I^T((j_1 \boxplus \psi_1)) \cdot e^{i\theta\widehat{\mathfrak{S}}_I^T((j_1 \boxplus \psi_1))}, \mathfrak{R}_I^T((j_2 \boxplus \psi_2)) \cdot e^{i\theta\widehat{\mathfrak{S}}_I^T((j_2 \boxplus \psi_2))}\} \\
 &\geq \min\{\min\{\mathfrak{R}_I^T(j_1) \cdot e^{i\theta\widehat{\mathfrak{S}}_I^T(j_1)}, \mathfrak{R}_I^T(\psi_1) \cdot e^{i\theta\widehat{\mathfrak{S}}_I^T(\psi_1)}\}, \\
 &\quad \min\{\mathfrak{R}_I^T(j_2) \cdot e^{i\theta\widehat{\mathfrak{S}}_I^T(j_2)}, \mathfrak{R}_I^T(\psi_2) \cdot e^{i\theta\widehat{\mathfrak{S}}_I^T(\psi_2)}\}\} \\
 &= \min\{\min\{\mathfrak{R}_I^T(j_1) \cdot e^{i\theta\widehat{\mathfrak{S}}_I^T(j_1)}, \mathfrak{R}_I^T(j_2) \cdot e^{i\theta\widehat{\mathfrak{S}}_I^T(j_2)}\}, \\
 &\quad \min\{\mathfrak{R}_I^T(\psi_1) \cdot e^{i\theta\widehat{\mathfrak{S}}_I^T(\psi_1)}, \mathfrak{R}_I^T(\psi_2) \cdot e^{i\theta\widehat{\mathfrak{S}}_I^T(\psi_2)}\}\} \\
 &= \min\{\mathfrak{R}_I^T((j_1, j_2)) \cdot e^{i\theta\widehat{\mathfrak{S}}_I^T((j_1, j_2))}, \mathfrak{R}_I^T((\psi_1, \psi_2)) \cdot e^{i\theta\widehat{\mathfrak{S}}_I^T((\psi_1, \psi_2))}\} \\
 &= \min\{\mathfrak{R}_I^T(j) \cdot e^{i\theta\widehat{\mathfrak{S}}_I^T(j)}, \mathfrak{R}_I^T(\psi) \cdot e^{i\theta\widehat{\mathfrak{S}}_I^T(\psi)}\}
 \end{aligned}$$

$$\text{Also } \mathfrak{R}_I^T((j \boxplus \psi)) \cdot e^{i\theta\widehat{\mathfrak{S}}_I^T((j \boxplus \psi))} \geq \min\{\mathfrak{R}_I^T(j) \cdot e^{i\theta\widehat{\mathfrak{S}}_I^T(j)}, \mathfrak{R}_I^T(\psi) \cdot e^{i\theta\widehat{\mathfrak{S}}_I^T(\psi)}\},$$

$$\mathfrak{R}_I^T((j \boxplus \psi)) \cdot e^{i\theta\widehat{\mathfrak{S}}_I^T((j \boxplus \psi))} \geq \min\{\mathfrak{R}_I^T(j) \cdot e^{i\theta\widehat{\mathfrak{S}}_I^T(j)}, \mathfrak{R}_I^T(\psi) \cdot e^{i\theta\widehat{\mathfrak{S}}_I^T(\psi)}\}.$$

Now,  $\Re_{\Upsilon}^I(j \boxplus_1 \psi) \cdot e^{i\theta \Im_{\Upsilon}^I(j \boxplus_1 \psi)}$

$$\begin{aligned}
 &= \Re_{\Upsilon}^I[(((j_1, j_2)) \boxplus_1 (\psi_1, \psi_2))] \cdot e^{i\theta \Im_{\Upsilon}^I[(((j_1, j_2)) \boxplus_1 (\psi_1, \psi_2))]} \\
 &= \Re_{\Upsilon}^I((j_1 \boxplus_1 \psi_1, j_2 \boxplus_1 \psi_2)) \cdot e^{i\theta \Im_{\Upsilon}^I((j_1 \boxplus_1 \psi_1, j_2 \boxplus_1 \psi_2))} \\
 &= \frac{\Re_{\Upsilon}^I((j_1 \boxplus_1 \psi_1)) \cdot e^{i\theta \Im_{\Upsilon}^I((j_1 \boxplus_1 \psi_1))} + \Re_{\Upsilon}^I((j_2 \boxplus_1 \psi_2)) \cdot e^{i\theta \Im_{\Upsilon}^I((j_2 \boxplus_1 \psi_2))}}{2} \\
 &\geq \frac{1}{2} \left[ \frac{\Re_{\Upsilon}^I(j_1) \cdot e^{i\theta \Im_{\Upsilon}^I(j_1)} + \Re_{\Upsilon}^I(\psi_1) \cdot e^{i\theta \Im_{\Upsilon}^I(\psi_1)}}{2} + \frac{\Re_{\Upsilon}^I(j_2) \cdot e^{i\theta \Im_{\Upsilon}^I(j_2)} + \Re_{\Upsilon}^I(\psi_2) \cdot e^{i\theta \Im_{\Upsilon}^I(\psi_2)}}{2} \right] \\
 &= \frac{1}{2} \left[ \frac{\Re_{\Upsilon}^I(j_1) \cdot e^{i\theta \Im_{\Upsilon}^I(j_1)} + \Re_{\Upsilon}^I(j_2) \cdot e^{i\theta \Im_{\Upsilon}^I(j_2)}}{2} + \frac{\Re_{\Upsilon}^I(\psi_1) \cdot e^{i\theta \Im_{\Upsilon}^I(\psi_1)} + \Re_{\Upsilon}^I(\psi_2) \cdot e^{i\theta \Im_{\Upsilon}^I(\psi_2)}}{2} \right] \\
 &= \frac{\Re_{\Upsilon}^I((j_1, j_2)) \cdot e^{i\theta \Im_{\Upsilon}^I((j_1, j_2))} + \Re_{\Upsilon}^I((\psi_1, \psi_2)) \cdot e^{i\theta \Im_{\Upsilon}^I((\psi_1, \psi_2))}}{2} \\
 &= \frac{\Re_{\Upsilon}^I(j) \cdot e^{i\theta \Im_{\Upsilon}^I(j)} + \Re_{\Upsilon}^I(\psi) \cdot e^{i\theta \Im_{\Upsilon}^I(\psi)}}{2}
 \end{aligned}$$

Also  $\Re_{\Upsilon}^I((j \boxplus_2 \psi)) \cdot e^{i\theta \Im_{\Upsilon}^I((j \boxplus_2 \psi))} \geq \frac{\Re_{\Upsilon}^I(j) \cdot e^{i\theta \Im_{\Upsilon}^I(j)} + \Re_{\Upsilon}^I(\psi) \cdot e^{i\theta \Im_{\Upsilon}^I(\psi)}}{2}$

and  $\Re_{\Upsilon}^I((j \boxplus_3 \psi)) \cdot e^{i\theta \Im_{\Upsilon}^I((j \boxplus_3 \psi))} \geq \frac{\Re_{\Upsilon}^I(j) \cdot e^{i\theta \Im_{\Upsilon}^I(j)} + \Re_{\Upsilon}^I(\psi) \cdot e^{i\theta \Im_{\Upsilon}^I(\psi)}}{2}$ .

Similarly,  $\Re_{\Upsilon}^F((j \boxplus_1 \psi)) \cdot e^{i\theta \Im_{\Upsilon}^F((j \boxplus_1 \psi))} \leq \max\{\Re_{\Upsilon}^F(j) \cdot e^{i\theta \Im_{\Upsilon}^F(j)}, \Re_{\Upsilon}^F(\psi) \cdot e^{i\theta \Im_{\Upsilon}^F(\psi)}\}$ ,

$\Re_{\Upsilon}^F((j \boxplus_2 \psi)) \cdot e^{i\theta \Im_{\Upsilon}^F((j \boxplus_2 \psi))} \leq \max\{\Re_{\Upsilon}^F(j) \cdot e^{i\theta \Im_{\Upsilon}^F(j)}, \Re_{\Upsilon}^F(\psi) \cdot e^{i\theta \Im_{\Upsilon}^F(\psi)}\}$  and

$\Re_{\Upsilon}^F((j \boxplus_3 \psi)) \cdot e^{i\theta \Im_{\Upsilon}^F((j \boxplus_3 \psi))} \leq \max\{\Re_{\Upsilon}^F(j) \cdot e^{i\theta \Im_{\Upsilon}^F(j)}, \Re_{\Upsilon}^F(\psi) \cdot e^{i\theta \Im_{\Upsilon}^F(\psi)}\}$ .

Therefore,  $\Upsilon$  is a ComCNSBS of  $S \times S$ .

Conversely, Assume that  $\Upsilon$  is a ComCNSBS of  $S \times S$ .

Let  $j = ((j_1, j_2))$ ,  $\psi = ((\psi_1, \psi_2)) \in S \times S$ . Now,

$$\begin{aligned}
 &\min\{\widehat{\Re}_{\Upsilon}^T((j_1 \boxplus_1 \psi_1)) \cdot e^{i\theta \Im_{\Upsilon}^T((j_1 \boxplus_1 \psi_1))}, \widehat{\Re}_{\Upsilon}^T((j_2 \boxplus_1 \psi_2)) \cdot e^{i\theta \Im_{\Upsilon}^T((j_2 \boxplus_1 \psi_2))}\} \\
 &= \widehat{\Re}_{\Upsilon}^T(j_1 \boxplus_1 \psi_1, j_2 \boxplus_1 \psi_2) \cdot e^{i\theta \Im_{\Upsilon}^T(j_1 \boxplus_1 \psi_1, j_2 \boxplus_1 \psi_2)} \\
 &= \widehat{\Re}_{\Upsilon}^T[(((j_1, j_2)) \boxplus_1 ((\psi_1, \psi_2)))] \cdot e^{i\theta \Im_{\Upsilon}^T[(((j_1, j_2)) \boxplus_1 ((\psi_1, \psi_2)))]} \\
 &= \widehat{\Re}_{\Upsilon}^T(j \boxplus_1 \psi) \cdot e^{i\theta \Im_{\Upsilon}^T(j \boxplus_1 \psi)} \\
 &\geq \min\{\widehat{\Re}_{\Upsilon}^T(j) \cdot e^{i\theta \Im_{\Upsilon}^T(j)}, \widehat{\Re}_{\Upsilon}^T(\psi) \cdot e^{i\theta \Im_{\Upsilon}^T(\psi)}\} \\
 &= \min\{\widehat{\Re}_{\Upsilon}^T((j_1, j_2)) \cdot e^{i\theta \Im_{\Upsilon}^T((j_1, j_2))}, \widehat{\Re}_{\Upsilon}^T((\psi_1, \psi_2)) \cdot e^{i\theta \Im_{\Upsilon}^T((\psi_1, \psi_2))}\} \\
 &= \min\{\min\{\widehat{\Re}_{\Upsilon}^T(j_1) \cdot e^{i\theta \Im_{\Upsilon}^T(j_1)}, \widehat{\Re}_{\Upsilon}^T(j_2) \cdot e^{i\theta \Im_{\Upsilon}^T(j_2)}\}, \min\{\widehat{\Re}_{\Upsilon}^T(\psi_1) \cdot e^{i\theta \Im_{\Upsilon}^T(\psi_1)}, \widehat{\Re}_{\Upsilon}^T(\psi_2) \cdot e^{i\theta \Im_{\Upsilon}^T(\psi_2)}\}\}
 \end{aligned}$$

If  $\widehat{\Re}_{\Upsilon}^T((j_1 \boxplus_1 \psi_1)) \cdot e^{i\theta \Im_{\Upsilon}^T((j_1 \boxplus_1 \psi_1))} \leq \widehat{\Re}_{\Upsilon}^T((j_2 \boxplus_1 \psi_2)) \cdot e^{i\theta \Im_{\Upsilon}^T((j_2 \boxplus_1 \psi_2))}$ , then  $\widehat{\Re}_{\Upsilon}^T(j_1) \cdot e^{i\theta \Im_{\Upsilon}^T(j_1)} \leq \widehat{\Re}_{\Upsilon}^T(j_2) \cdot e^{i\theta \Im_{\Upsilon}^T(j_2)}$  and  $\widehat{\Re}_{\Upsilon}^T(\psi_1) \cdot e^{i\theta \Im_{\Upsilon}^T(\psi_1)} \leq \widehat{\Re}_{\Upsilon}^T(\psi_2) \cdot e^{i\theta \Im_{\Upsilon}^T(\psi_2)}$ . We get  $\widehat{\Re}_{\Upsilon}^T((j_1 \boxplus_1 \psi_1)) \cdot e^{i\theta \Im_{\Upsilon}^T((j_1 \boxplus_1 \psi_1))} \geq$

$\min\{\widehat{\Re}_{\Upsilon}^T(j_1) \cdot e^{i\theta \Im_{\Upsilon}^T(j_1)}, \widehat{\Re}_{\Upsilon}^T(\psi_1) \cdot e^{i\theta \Im_{\Upsilon}^T(\psi_1)}\}$  for all  $j_1, \psi_1 \in S$ , and

$\min\{\widehat{\Re}_{\Upsilon}^T((j_1 \boxplus_2 \psi_1)) \cdot e^{i\theta \Im_{\Upsilon}^T((j_1 \boxplus_2 \psi_1))}, \widehat{\Re}_{\Upsilon}^T((j_2 \boxplus_2 \psi_2)) \cdot e^{i\theta \Im_{\Upsilon}^T((j_2 \boxplus_2 \psi_2))}\} \geq \min\{\min\{\widehat{\Re}_{\Upsilon}^T(j_1) \cdot e^{i\theta \Im_{\Upsilon}^T(j_1)}, \widehat{\Re}_{\Upsilon}^T(j_2) \cdot e^{i\theta \Im_{\Upsilon}^T(j_2)}\}, \min\{\widehat{\Re}_{\Upsilon}^T(\psi_1) \cdot e^{i\theta \Im_{\Upsilon}^T(\psi_1)}, \widehat{\Re}_{\Upsilon}^T(\psi_2) \cdot e^{i\theta \Im_{\Upsilon}^T(\psi_2)}\}\}$

If  $\widehat{\Re}_{\Upsilon}^T((j_1 \boxplus_2 \psi_1)) \cdot e^{i\theta \Im_{\Upsilon}^T((j_1 \boxplus_2 \psi_1))} \leq \widehat{\Re}_{\Upsilon}^T((j_2 \boxplus_2 \psi_2)) \cdot e^{i\theta \Im_{\Upsilon}^T((j_2 \boxplus_2 \psi_2))}$ , then  $\widehat{\Re}_{\Upsilon}^T((j_1 \boxplus_2 \psi_1)) \cdot e^{i\theta \Im_{\Upsilon}^T((j_1 \boxplus_2 \psi_1))} \geq$

$\min\{\widehat{\Re}_{\Upsilon}^T(j_1) \cdot e^{i\theta \Im_{\Upsilon}^T(j_1)}, \widehat{\Re}_{\Upsilon}^T(\psi_1) \cdot e^{i\theta \Im_{\Upsilon}^T(\psi_1)}\}$ .

$\min\{\widehat{\Re}_{\Upsilon}^T((j_1 \boxplus_3 \psi_1)) \cdot e^{i\theta \Im_{\Upsilon}^T((j_1 \boxplus_3 \psi_1))}, \widehat{\Re}_{\Upsilon}^T((j_2 \boxplus_3 \psi_2)) \cdot e^{i\theta \Im_{\Upsilon}^T((j_2 \boxplus_3 \psi_2))}\} \geq \min\{\min\{\widehat{\Re}_{\Upsilon}^T(j_1) \cdot e^{i\theta \Im_{\Upsilon}^T(j_1)}, \widehat{\Re}_{\Upsilon}^T(j_2) \cdot e^{i\theta \Im_{\Upsilon}^T(j_2)}\}, \min\{\widehat{\Re}_{\Upsilon}^T(\psi_1) \cdot e^{i\theta \Im_{\Upsilon}^T(\psi_1)}, \widehat{\Re}_{\Upsilon}^T(\psi_2) \cdot e^{i\theta \Im_{\Upsilon}^T(\psi_2)}\}\}$

If  $\widehat{\Re}_{\Upsilon}^T((j_1 \boxplus_3 \psi_1)) \cdot e^{i\theta \Im_{\Upsilon}^T((j_1 \boxplus_3 \psi_1))} \leq \widehat{\Re}_{\Upsilon}^T((j_2 \boxplus_3 \psi_2)) \cdot e^{i\theta \Im_{\Upsilon}^T((j_2 \boxplus_3 \psi_2))}$ , then  $\widehat{\Re}_{\Upsilon}^T((j_1 \boxplus_3 \psi_1)) \cdot e^{i\theta \Im_{\Upsilon}^T((j_1 \boxplus_3 \psi_1))} \geq$

$\min\{\widehat{\Re}_{\Upsilon}^T(j_1) \cdot e^{i\theta \Im_{\Upsilon}^T(j_1)}, \widehat{\Re}_{\Upsilon}^T(\psi_1) \cdot e^{i\theta \Im_{\Upsilon}^T(\psi_1)}\}$ .

Now,

$$\begin{aligned}
& \frac{1}{2} \left[ \widehat{\mathcal{R}}_{\Gamma}^I((j_1 \boxplus_1 \psi_1)) \cdot e^{i\theta \widehat{\mathfrak{Z}}_{\Gamma}^I((j_1 \boxplus_1 \psi_1))} + \widehat{\mathcal{R}}_{\Gamma}^I((j_2 \boxplus_1 \psi_2)) \cdot e^{i\theta \widehat{\mathfrak{Z}}_{\Gamma}^I((j_2 \boxplus_1 \psi_2))} \right] \\
&= \widehat{\mathcal{R}}_{\Gamma}^I(j_1 \boxplus_1 \psi_1, j_2 \boxplus_1 \psi_2) \cdot e^{i\theta \widehat{\mathfrak{Z}}_{\Gamma}^I(j_1 \boxplus_1 \psi_1, j_2 \boxplus_1 \psi_2)} \\
&= \widehat{\mathcal{R}}_{\Gamma}^I([(j_1, j_2)] \boxplus_1 ((\psi_1, \psi_2))) \cdot e^{i\theta \widehat{\mathfrak{Z}}_{\Gamma}^I([(j_1, j_2)] \boxplus_1 ((\psi_1, \psi_2)))} \\
&= \widehat{\mathcal{R}}_{\Gamma}^I((j \boxplus_1 \psi)) \cdot e^{i\theta \widehat{\mathfrak{Z}}_{\Gamma}^I((j \boxplus_1 \psi))} \\
&\geq \frac{\widehat{\mathcal{R}}_{\Gamma}^I(j) \cdot e^{i\theta \widehat{\mathfrak{Z}}_{\Gamma}^I(j)} + \widehat{\mathcal{R}}_{\Gamma}^I(\psi) \cdot e^{i\theta \widehat{\mathfrak{Z}}_{\Gamma}^I(\psi)}}{2} \\
&= \frac{\widehat{\mathcal{R}}_{\Gamma}^I((j_1, j_2)) \cdot e^{i\theta \widehat{\mathfrak{Z}}_{\Gamma}^I((j_1, j_2))} + \widehat{\mathcal{R}}_{\Gamma}^I((\psi_1, \psi_2)) \cdot e^{i\theta \widehat{\mathfrak{Z}}_{\Gamma}^I((\psi_1, \psi_2))}}{2} \\
&= \frac{1}{2} \left[ \frac{\widehat{\mathcal{R}}_{\Gamma}^I(j_1) \cdot e^{i\theta \widehat{\mathfrak{Z}}_{\Gamma}^I(j_1)} + \widehat{\mathcal{R}}_{\Gamma}^I(j_2) \cdot e^{i\theta \widehat{\mathfrak{Z}}_{\Gamma}^I(j_2)}}{2} + \frac{\widehat{\mathcal{R}}_{\Gamma}^I(\psi_1) \cdot e^{i\theta \widehat{\mathfrak{Z}}_{\Gamma}^I(\psi_1)} + \widehat{\mathcal{R}}_{\Gamma}^I(\psi_2) \cdot e^{i\theta \widehat{\mathfrak{Z}}_{\Gamma}^I(\psi_2)}}{2} \right]
\end{aligned}$$

If  $\widehat{\mathfrak{R}}_T^I((j_1 \boxplus_1 \psi_1)) \cdot e^{i\theta\widehat{\mathfrak{S}}_T^I((j_1 \boxplus_1 \psi_1))} \leq \widehat{\mathfrak{R}}_T^I((j_2 \boxplus_1 \psi_2)) \cdot e^{i\theta\widehat{\mathfrak{S}}_T^I((j_2 \boxplus_1 \psi_2))}$ , then  $\widehat{\mathfrak{R}}_T^I(j_1) \cdot e^{i\theta\widehat{\mathfrak{S}}_T^I(j_1)} \leq \widehat{\mathfrak{R}}_T^I(j_2) \cdot e^{i\theta\widehat{\mathfrak{S}}_T^I(j_2)}$  and  $\widehat{\mathfrak{R}}_T^I(\psi_1) \cdot e^{i\theta\widehat{\mathfrak{S}}_T^I(\psi_1)} < \widehat{\mathfrak{R}}_T^I(\psi_2) \cdot e^{i\theta\widehat{\mathfrak{S}}_T^I(\psi_2)}$ .

$$\text{We get } \widehat{\mathfrak{R}}_{\Gamma}^I((j_1 \boxplus_1 \psi_1)) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Gamma}^T((j_1 \boxplus_1 \psi_1))} \geq \frac{\widehat{\mathfrak{R}}_{\Gamma}^I(j_1) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Gamma}^T(j_1)} + \widehat{\mathfrak{R}}_{\Gamma}^I(\psi_1) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Gamma}^T(\psi_1)}}{2}.$$

Similarly,  $\widehat{\mathfrak{R}}_{\Gamma}((j_1 \boxplus_2 \psi_1)) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Gamma}((j_1 \boxplus_2 \psi_1))} \geq \frac{\widehat{\mathfrak{R}}_{\Gamma}(j_1) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Gamma}(j_1)} + \widehat{\mathfrak{R}}_{\Gamma}(\psi_1) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Gamma}(\psi_1)}}{2}$

$$\text{and } \widehat{\mathfrak{R}}_{\Gamma}^I((j_1 \boxplus_3 \psi_1)) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Gamma}^I((j_1 \boxplus_3 \psi_1))} \geq \frac{\widehat{\mathfrak{R}}_{\Gamma}^I(j_1) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Gamma}^I(j_1)} + \widehat{\mathfrak{R}}_{\Gamma}^I(\psi_1) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Gamma}^I(\psi_1)}}{2}.$$

Similarly to prove that

$$\max\{\widehat{\mathfrak{R}}_r^F((j_1 \boxplus_1 \psi_1)) \cdot e^{i\theta \widehat{\mathfrak{Z}}_r^F((j_1 \boxplus_1 \psi_1))}, \widehat{\mathfrak{R}}_r^F((j_2 \boxplus_1 \psi_2)) \cdot e^{i\theta \widehat{\mathfrak{Z}}_r^F((j_2 \boxplus_1 \psi_2))}\} \leq \max\{\max\{\widehat{\mathfrak{R}}_r^F(j_1) \cdot e^{i\theta \widehat{\mathfrak{Z}}_r^F(j_1)}, \widehat{\mathfrak{R}}_r^F(j_2) \cdot e^{i\theta \widehat{\mathfrak{Z}}_r^F(j_2)}\}, \max\{\widehat{\mathfrak{R}}_r^F(\psi_1) \cdot e^{i\theta \widehat{\mathfrak{Z}}_r^F(\psi_1)}, \widehat{\mathfrak{R}}_r^F(\psi_2) \cdot e^{i\theta \widehat{\mathfrak{Z}}_r^F(\psi_2)}\}\}$$

If  $\widehat{\mathfrak{R}}_F^F((j_1 \boxplus_1 \psi_1)) \cdot e^{i\theta \widehat{\mathfrak{S}}_F^F((j_1 \boxplus_1 \psi_1))} \geq \widehat{\mathfrak{R}}_F^F((j_2 \boxplus_1 \psi_2)) \cdot e^{i\theta \widehat{\mathfrak{S}}_F^F((j_2 \boxplus_1 \psi_2))}$ , then  $\widehat{\mathfrak{R}}_F^F(j_1) \cdot e^{i\theta \widehat{\mathfrak{S}}_F^F(j_1)} \geq \widehat{\mathfrak{R}}_F^F(j_2) \cdot e^{i\theta \widehat{\mathfrak{S}}_F^F(j_2)}$  and  $\widehat{\mathfrak{R}}_F^F(\psi_1) \cdot e^{i\theta \widehat{\mathfrak{S}}_F^F(\psi_1)} \geq \widehat{\mathfrak{R}}_F^F(\psi_2) \cdot e^{i\theta \widehat{\mathfrak{S}}_F^F(\psi_2)}$ .

We get  $\widehat{\mathfrak{R}}_{\Gamma}^F((j_1 \boxplus_1 \psi_1)) \cdot e^{i\widehat{\mathfrak{S}}_{\Gamma}^F((j_1 \boxplus_1 \psi_1))} \leq \max\{\widehat{\mathfrak{R}}_{\Gamma}^F(j_1) \cdot e^{i\widehat{\mathfrak{S}}_{\Gamma}^F(j_1)}, \widehat{\mathfrak{R}}_{\Gamma}^F(\psi_1) \cdot e^{i\widehat{\mathfrak{S}}_{\Gamma}^F(\psi_1)}\}.$

$$\begin{aligned} & \max\{\widehat{\mathfrak{R}}_F^F((j_1 \boxplus_2 \psi_1)) \cdot e^{i\theta\mathfrak{Z}_F^F(\widehat{(j_1 \boxplus_2 \psi_1)})}, \widehat{\mathfrak{R}}_F^F((j_2 \boxplus_2 \psi_2)) \cdot e^{i\theta\mathfrak{Z}_F^F(\widehat{(j_2 \boxplus_2 \psi_2)})}\} \\ & \leq \max\{\max\{\widehat{\mathfrak{R}}_F^F(j_1) \cdot e^{i\theta\mathfrak{Z}_F^F(j_1)}, \widehat{\mathfrak{R}}_F^F(j_2) \cdot e^{i\theta\mathfrak{Z}_F^F(j_2)}\}, \max\{\widehat{\mathfrak{R}}_F^F(\psi_1) \cdot e^{i\theta\mathfrak{Z}_F^F(\psi_1)}, \widehat{\mathfrak{R}}_F^F(\psi_2) \cdot e^{i\theta\mathfrak{Z}_F^F(\psi_2)}\}\} \end{aligned}$$

If  $\widehat{\mathfrak{R}}_F^F((j_1 \boxplus_2 \psi_1)) \cdot e^{i\theta \Im_F^F((\widehat{j_1 \boxplus_2 \psi_1}))} \geq \widehat{\mathfrak{R}}_F^F((j_2 \boxplus_2 \psi_2)) \cdot e^{i\theta \Im_F^F((\widehat{j_2 \boxplus_2 \psi_2}))}$ , then  $\widehat{\mathfrak{R}}_F^F((j_1 \boxplus_2 \psi_1)) \cdot e^{i\theta \Im_F^F((\widehat{j_1 \boxplus_2 \psi_1}))} \leq \max\{\widehat{\mathfrak{R}}_F^F(j_1) \cdot e^{i\theta \Im_F^F(j_1)}, \widehat{\mathfrak{R}}_F^F(\psi_1) \cdot e^{i\theta \Im_F^F(\psi_1)}\}$ .

$$\max\{\widehat{\mathfrak{R}}_F^F((j_1 \boxplus_3 \psi_1)) \cdot e^{i\theta \widehat{\mathfrak{S}}_F^F((j_1 \boxplus_3 \psi_1))}, \widehat{\mathfrak{R}}_F^F((j_2 \boxplus_3 \psi_2)) \cdot e^{i\theta \widehat{\mathfrak{S}}_F^F((j_2 \boxplus_3 \psi_2))}\} \leq \max\{\max\{\widehat{\mathfrak{R}}_F^F(j_1) \cdot e^{i\theta \widehat{\mathfrak{S}}_F^F(j_1)}, \widehat{\mathfrak{R}}_F^F(j_2) \cdot e^{i\theta \widehat{\mathfrak{S}}_F^F(j_2)}\}, \max\{\widehat{\mathfrak{R}}_F^F(\psi_1) \cdot e^{i\theta \widehat{\mathfrak{S}}_F^F(\psi_1)}, \widehat{\mathfrak{R}}_F^F(\psi_2) \cdot e^{i\theta \widehat{\mathfrak{S}}_F^F(\psi_2)}\}\}$$

If  $\widehat{\mathfrak{R}}_F^F((j_1 \boxplus_3 \psi_1)) \cdot e^{i\theta \Im_F^T((j_1 \boxplus_3 \psi_1))} \geq \widehat{\mathfrak{R}}_F^F((j_2 \boxplus_3 \psi_2)) \cdot e^{i\theta \Im_F^T((j_2 \boxplus_3 \psi_2))}$ , then  $\widehat{\mathfrak{R}}_F^F((j_1 \boxplus_3 \psi_1)) \cdot e^{i\theta \Im_F^T((j_1 \boxplus_3 \psi_1))} \leq \max\{\widehat{\mathfrak{R}}_F^F(j_1) \cdot e^{i\theta \Im_F^T(j_1)}, \widehat{\mathfrak{R}}_F^F(\psi_1) \cdot e^{i\theta \Im_F^T(\psi_1)}\}$ .

Let  $j = ((j_1, j_2)), \psi = ((\psi_1, \psi_2)) \in S \times S$ . Now,

$$\begin{aligned}
& \min\{\mathfrak{R}_T^T((j_1 \boxplus_1 \psi_1)) \cdot e^{i\theta\mathfrak{Z}_T^T((j_1 \boxplus_1 \psi_1))}, \mathfrak{R}_T^T((j_2 \boxplus_1 \psi_2)) \cdot e^{i\theta\mathfrak{Z}_T^T((j_2 \boxplus_1 \psi_2))}\} \\
&= \mathfrak{R}_T^T(j_1 \boxplus_1 \psi_1, j_2 \boxplus_1 \psi_2) \cdot e^{i\theta\mathfrak{Z}_T^T(j_1 \boxplus_1 \psi_1, j_2 \boxplus_1 \psi_2)} \\
&= \mathfrak{R}_T^T([(j_1, j_2)) \boxplus_1 ((\psi_1, \psi_2))]) \cdot e^{i\theta\mathfrak{Z}_T^T([(j_1, j_2)) \boxplus_1 ((\psi_1, \psi_2))])} \\
&= \mathfrak{R}_T^T(j \boxplus_1 \psi) \cdot e^{i\theta\mathfrak{Z}_T^T(j \boxplus_1 \psi)} \\
&\geq \min\{\mathfrak{R}_T^T(j) \cdot e^{i\theta\mathfrak{Z}_T^T(j)}, \mathfrak{R}_T^T(\psi) \cdot e^{i\theta\mathfrak{Z}_T^T(\psi)}\} \\
&= \min\{\mathfrak{R}_T^T((j_1, j_2))\} \cdot e^{i\theta\mathfrak{Z}_T^T((j_1, j_2))}, \mathfrak{R}_T^T((\psi_1, \psi_2)) \cdot e^{i\theta\mathfrak{Z}_T^T((\psi_1, \psi_2))}\} \\
&= \min\{\min\{\mathfrak{R}_T^T(j_1) \cdot e^{i\theta\mathfrak{Z}_T^T(j_1)}, \mathfrak{R}_T^T(j_2) \cdot e^{i\theta\mathfrak{Z}_T^T(j_2)}\}, \min\{\mathfrak{R}_T^T(\psi_1) \cdot e^{i\theta\mathfrak{Z}_T^T(\psi_1)}, \mathfrak{R}_T^T(\psi_2) \cdot e^{i\theta\mathfrak{Z}_T^T(\psi_2)}\}\}
\end{aligned}$$

If  $\mathfrak{R}_\Gamma^T((j_1 \boxplus_1 \psi_1)) \cdot e^{i\theta\mathfrak{S}_\Gamma^T((j_1 \boxplus_1 \psi_1))} \leq \mathfrak{R}_\Gamma^T((j_2 \boxplus_1 \psi_2)) \cdot e^{i\theta\mathfrak{S}_\Gamma^T((j_2 \boxplus_1 \psi_2))}$ , then  $\mathfrak{R}_\Gamma^T(j_1) \cdot e^{i\theta\mathfrak{S}_\Gamma^T(j_1)} \leq \mathfrak{R}_\Gamma^T(j_2) \cdot e^{i\theta\mathfrak{S}_\Gamma^T(j_2)}$  and  $\mathfrak{R}_\Gamma^T(\psi_1) \cdot e^{i\theta\mathfrak{S}_\Gamma^T(\psi_1)} \leq \mathfrak{R}_\Gamma^T(\psi_2) \cdot e^{i\theta\mathfrak{S}_\Gamma^T(\psi_2)}$ . We get  $\mathfrak{R}_\Gamma^T((j_1 \boxplus_1 \psi_1)) \cdot e^{i\theta\mathfrak{S}_\Gamma^T((j_1 \boxplus_1 \psi_1))} \geq \min\{\mathfrak{R}_\Gamma^T(j_1) \cdot e^{i\theta\mathfrak{S}_\Gamma^T(j_1)}, \mathfrak{R}_\Gamma^T(\psi_1) \cdot e^{i\theta\mathfrak{S}_\Gamma^T(\psi_1)}\}$  for all  $j_1, \psi_1 \in S$ , and  $\min\{\mathfrak{R}_\Gamma^T((j_1 \boxplus_2 \psi_1)) \cdot e^{i\theta\mathfrak{S}_\Gamma^T((j_1 \boxplus_2 \psi_1))}, \mathfrak{R}_\Gamma^T((j_2 \boxplus_2 \psi_2)) \cdot e^{i\theta\mathfrak{S}_\Gamma^T((j_2 \boxplus_2 \psi_2))}\} \geq \min\{\min\{\mathfrak{R}_\Gamma^T(j_1) \cdot e^{i\theta\mathfrak{S}_\Gamma^T(j_1)}, \mathfrak{R}_\Gamma^T(j_2) \cdot e^{i\theta\mathfrak{S}_\Gamma^T(j_2)}\}, \min\{\mathfrak{R}_\Gamma^T(\psi_1) \cdot e^{i\theta\mathfrak{S}_\Gamma^T(\psi_1)}, \mathfrak{R}_\Gamma^T(\psi_2) \cdot e^{i\theta\mathfrak{S}_\Gamma^T(\psi_2)}\}\}$ . If  $\mathfrak{R}_\Gamma^T((j_1 \boxplus_2 \psi_1)) \cdot e^{i\theta\mathfrak{S}_\Gamma^T((j_1 \boxplus_2 \psi_1))} \leq \mathfrak{R}_\Gamma^T((j_2 \boxplus_2 \psi_2)) \cdot e^{i\theta\mathfrak{S}_\Gamma^T((j_2 \boxplus_2 \psi_2))}$ , then  $\mathfrak{R}_\Gamma^T((j_1 \boxplus_2 \psi_1)) \cdot e^{i\theta\mathfrak{S}_\Gamma^T((j_1 \boxplus_2 \psi_1))} \geq \min\{\mathfrak{R}_\Gamma^T(j_1) \cdot e^{i\theta\mathfrak{S}_\Gamma^T(j_1)}, \mathfrak{R}_\Gamma^T(\psi_1) \cdot e^{i\theta\mathfrak{S}_\Gamma^T(\psi_1)}\}$ .  $\min\{\mathfrak{R}_\Gamma^T((j_1 \boxplus_3 \psi_1)) \cdot e^{i\theta\mathfrak{S}_\Gamma^T((j_1 \boxplus_3 \psi_1))}, \mathfrak{R}_\Gamma^T((j_2 \boxplus_3 \psi_2)) \cdot e^{i\theta\mathfrak{S}_\Gamma^T((j_2 \boxplus_3 \psi_2))}\} \geq \min\{\min\{\mathfrak{R}_\Gamma^T(j_1) \cdot e^{i\theta\mathfrak{S}_\Gamma^T(j_1)}, \mathfrak{R}_\Gamma^T(j_2) \cdot e^{i\theta\mathfrak{S}_\Gamma^T(j_2)}\}, \min\{\mathfrak{R}_\Gamma^T(\psi_1) \cdot e^{i\theta\mathfrak{S}_\Gamma^T(\psi_1)}, \mathfrak{R}_\Gamma^T(\psi_2) \cdot e^{i\theta\mathfrak{S}_\Gamma^T(\psi_2)}\}\}$ . If  $\mathfrak{R}_\Gamma^T((j_1 \boxplus_3 \psi_1)) \cdot e^{i\theta\mathfrak{S}_\Gamma^T((j_1 \boxplus_3 \psi_1))} \leq \mathfrak{R}_\Gamma^T((j_2 \boxplus_3 \psi_2)) \cdot e^{i\theta\mathfrak{S}_\Gamma^T((j_2 \boxplus_3 \psi_2))}$ , then  $\mathfrak{R}_\Gamma^T((j_1 \boxplus_3 \psi_1)) \cdot e^{i\theta\mathfrak{S}_\Gamma^T((j_1 \boxplus_3 \psi_1))} \geq \min\{\mathfrak{R}_\Gamma^T(j_1) \cdot e^{i\theta\mathfrak{S}_\Gamma^T(j_1)}, \mathfrak{R}_\Gamma^T(\psi_1) \cdot e^{i\theta\mathfrak{S}_\Gamma^T(\psi_1)}\}$ . Now,

$$\begin{aligned} & \frac{1}{2} \left[ \mathfrak{R}_\Gamma^I((j_1 \boxplus_1 \psi_1)) \cdot e^{i\theta\mathfrak{S}_\Gamma^I((j_1 \boxplus_1 \psi_1))} + \mathfrak{R}_\Gamma^I((j_2 \boxplus_1 \psi_2)) \cdot e^{i\theta\mathfrak{S}_\Gamma^I((j_2 \boxplus_1 \psi_2))} \right] \\ &= \mathfrak{R}_\Gamma^I(j_1 \boxplus_1 \psi_1, j_2 \boxplus_1 \psi_2) \cdot e^{i\theta\mathfrak{S}_\Gamma^I(j_1 \boxplus_1 \psi_1, j_2 \boxplus_1 \psi_2)} \\ &= \mathfrak{R}_\Gamma^I([(j_1, j_2)] \boxplus_1 [(\psi_1, \psi_2)]) \cdot e^{i\theta\mathfrak{S}_\Gamma^I([(j_1, j_2)] \boxplus_1 [(\psi_1, \psi_2)])} \\ &= \mathfrak{R}_\Gamma^I(j \boxplus_1 \psi) \cdot e^{i\theta\mathfrak{S}_\Gamma^I(j \boxplus_1 \psi)} \\ &\geq \frac{\mathfrak{R}_\Gamma^I(j) \cdot e^{i\theta\mathfrak{S}_\Gamma^I(j)} + \mathfrak{R}_\Gamma^I(\psi) \cdot e^{i\theta\mathfrak{S}_\Gamma^I(\psi)}}{2} \\ &= \frac{\mathfrak{R}_\Gamma^I((j_1, j_2)) \cdot e^{i\theta\mathfrak{S}_\Gamma^I((j_1, j_2))} + \mathfrak{R}_\Gamma^I((\psi_1, \psi_2)) \cdot e^{i\theta\mathfrak{S}_\Gamma^I((\psi_1, \psi_2))}}{2} \\ &= \frac{1}{2} \left[ \frac{\mathfrak{R}_\Gamma^I(j_1) \cdot e^{i\theta\mathfrak{S}_\Gamma^I(j_1)} + \mathfrak{R}_\Gamma^I(j_2) \cdot e^{i\theta\mathfrak{S}_\Gamma^I(j_2)}}{2} + \frac{\mathfrak{R}_\Gamma^I(\psi_1) \cdot e^{i\theta\mathfrak{S}_\Gamma^I(\psi_1)} + \mathfrak{R}_\Gamma^I(\psi_2) \cdot e^{i\theta\mathfrak{S}_\Gamma^I(\psi_2)}}{2} \right] \end{aligned}$$

If  $\mathfrak{R}_\Gamma^I((j_1 \boxplus_1 \psi_1)) \cdot e^{i\theta\mathfrak{S}_\Gamma^I((j_1 \boxplus_1 \psi_1))} \leq \mathfrak{R}_\Gamma^I((j_2 \boxplus_1 \psi_2)) \cdot e^{i\theta\mathfrak{S}_\Gamma^I((j_2 \boxplus_1 \psi_2))}$ , then  $\mathfrak{R}_\Gamma^I(j_1) \cdot e^{i\theta\mathfrak{S}_\Gamma^I(j_1)} \leq \mathfrak{R}_\Gamma^I(j_2) \cdot e^{i\theta\mathfrak{S}_\Gamma^I(j_2)}$  and  $\mathfrak{R}_\Gamma^I(\psi_1) \cdot e^{i\theta\mathfrak{S}_\Gamma^I(\psi_1)} \leq \mathfrak{R}_\Gamma^I(\psi_2) \cdot e^{i\theta\mathfrak{S}_\Gamma^I(\psi_2)}$ .

We get  $\mathfrak{R}_\Gamma^I((j_1 \boxplus_1 \psi_1)) \cdot e^{i\theta\mathfrak{S}_\Gamma^I((j_1 \boxplus_1 \psi_1))} \geq \frac{\mathfrak{R}_\Gamma^I(j_1) \cdot e^{i\theta\mathfrak{S}_\Gamma^I(j_1)} + \mathfrak{R}_\Gamma^I(\psi_1) \cdot e^{i\theta\mathfrak{S}_\Gamma^I(\psi_1)}}{2}$ .

Similarly,  $\mathfrak{R}_\Gamma^I((j_1 \boxplus_2 \psi_1)) \cdot e^{i\theta\mathfrak{S}_\Gamma^I((j_1 \boxplus_2 \psi_1))} \geq \frac{\mathfrak{R}_\Gamma^I(j_1) \cdot e^{i\theta\mathfrak{S}_\Gamma^I(j_1)} + \mathfrak{R}_\Gamma^I(\psi_1) \cdot e^{i\theta\mathfrak{S}_\Gamma^I(\psi_1)}}{2}$

and  $\mathfrak{R}_\Gamma^I((j_1 \boxplus_3 \psi_1)) \cdot e^{i\theta\mathfrak{S}_\Gamma^I((j_1 \boxplus_3 \psi_1))} \geq \frac{\mathfrak{R}_\Gamma^I(j_1) \cdot e^{i\theta\mathfrak{S}_\Gamma^I(j_1)} + \mathfrak{R}_\Gamma^I(\psi_1) \cdot e^{i\theta\mathfrak{S}_\Gamma^I(\psi_1)}}{2}$ .

Similarly to prove that

$\max\{\mathfrak{R}_\Gamma^F((j_1 \boxplus_1 \psi_1)) \cdot e^{i\theta\mathfrak{S}_\Gamma^F((j_1 \boxplus_1 \psi_1))}, \mathfrak{R}_\Gamma^F((j_2 \boxplus_1 \psi_2)) \cdot e^{i\theta\mathfrak{S}_\Gamma^F((j_2 \boxplus_1 \psi_2))}\} \leq \max\{\max\{\mathfrak{R}_\Gamma^F(j_1) \cdot e^{i\theta\mathfrak{S}_\Gamma^F(j_1)}, \mathfrak{R}_\Gamma^F(j_2) \cdot e^{i\theta\mathfrak{S}_\Gamma^F(j_2)}\}, \max\{\mathfrak{R}_\Gamma^F(\psi_1) \cdot e^{i\theta\mathfrak{S}_\Gamma^F(\psi_1)}, \mathfrak{R}_\Gamma^F(\psi_2) \cdot e^{i\theta\mathfrak{S}_\Gamma^F(\psi_2)}\}\}$ . If  $\mathfrak{R}_\Gamma^F((j_1 \boxplus_1 \psi_1)) \cdot e^{i\theta\mathfrak{S}_\Gamma^F((j_1 \boxplus_1 \psi_1))} \geq \mathfrak{R}_\Gamma^F((j_2 \boxplus_1 \psi_2)) \cdot e^{i\theta\mathfrak{S}_\Gamma^F((j_2 \boxplus_1 \psi_2))}$ , then  $\mathfrak{R}_\Gamma^F(j_1) \cdot e^{i\theta\mathfrak{S}_\Gamma^F(j_1)} \geq \mathfrak{R}_\Gamma^F(j_2) \cdot e^{i\theta\mathfrak{S}_\Gamma^F(j_2)}$  and  $\mathfrak{R}_\Gamma^F(\psi_1) \cdot e^{i\theta\mathfrak{S}_\Gamma^F(\psi_1)} \geq \mathfrak{R}_\Gamma^F(\psi_2) \cdot e^{i\theta\mathfrak{S}_\Gamma^F(\psi_2)}$ .

We get  $\mathfrak{R}_\Gamma^F((j_1 \boxplus_1 \psi_1)) \cdot e^{i\theta\mathfrak{S}_\Gamma^F((j_1 \boxplus_1 \psi_1))} \leq \max\{\mathfrak{R}_\Gamma^F(j_1) \cdot e^{i\theta\mathfrak{S}_\Gamma^F(j_1)}, \mathfrak{R}_\Gamma^F(\psi_1) \cdot e^{i\theta\mathfrak{S}_\Gamma^F(\psi_1)}\}$ .

$\max\{\mathfrak{R}_\Gamma^F((j_1 \boxplus_2 \psi_1)) \cdot e^{i\theta\mathfrak{S}_\Gamma^F((j_1 \boxplus_2 \psi_1))}, \mathfrak{R}_\Gamma^F((j_2 \boxplus_2 \psi_2)) \cdot e^{i\theta\mathfrak{S}_\Gamma^F((j_2 \boxplus_2 \psi_2))}\} \leq \max\{\max\{\mathfrak{R}_\Gamma^F(j_1) \cdot e^{i\theta\mathfrak{S}_\Gamma^F(j_1)}, \mathfrak{R}_\Gamma^F(j_2) \cdot e^{i\theta\mathfrak{S}_\Gamma^F(j_2)}\}, \max\{\mathfrak{R}_\Gamma^F(\psi_1) \cdot e^{i\theta\mathfrak{S}_\Gamma^F(\psi_1)}, \mathfrak{R}_\Gamma^F(\psi_2) \cdot e^{i\theta\mathfrak{S}_\Gamma^F(\psi_2)}\}\}$

If  $\mathfrak{R}_\Gamma^F((j_1 \boxplus_2 \psi_1)) \cdot e^{i\theta\mathfrak{S}_\Gamma^F((j_1 \boxplus_2 \psi_1))} \geq \mathfrak{R}_\Gamma^F((j_2 \boxplus_2 \psi_2)) \cdot e^{i\theta\mathfrak{S}_\Gamma^F((j_2 \boxplus_2 \psi_2))}$ , then  $\mathfrak{R}_\Gamma^F((j_1 \boxplus_2 \psi_1)) \cdot e^{i\theta\mathfrak{S}_\Gamma^F((j_1 \boxplus_2 \psi_1))} \leq \max\{\mathfrak{R}_\Gamma^F(j_1) \cdot e^{i\theta\mathfrak{S}_\Gamma^F(j_1)}, \mathfrak{R}_\Gamma^F(\psi_1) \cdot e^{i\theta\mathfrak{S}_\Gamma^F(\psi_1)}\}$ .

$\max\{\mathfrak{R}_\Gamma^F((j_1 \boxplus_3 \psi_1)) \cdot e^{i\theta\mathfrak{S}_\Gamma^F((j_1 \boxplus_3 \psi_1))}, \mathfrak{R}_\Gamma^F((j_2 \boxplus_3 \psi_2)) \cdot e^{i\theta\mathfrak{S}_\Gamma^F((j_2 \boxplus_3 \psi_2))}\} \leq \max\{\max\{\mathfrak{R}_\Gamma^F(j_1) \cdot e^{i\theta\mathfrak{S}_\Gamma^F(j_1)}, \mathfrak{R}_\Gamma^F(j_2) \cdot e^{i\theta\mathfrak{S}_\Gamma^F(j_2)}\}, \max\{\mathfrak{R}_\Gamma^F(\psi_1) \cdot e^{i\theta\mathfrak{S}_\Gamma^F(\psi_1)}, \mathfrak{R}_\Gamma^F(\psi_2) \cdot e^{i\theta\mathfrak{S}_\Gamma^F(\psi_2)}\}\}$

If  $\mathfrak{R}_\Gamma^F((j_1 \boxplus_3 \psi_1)) \cdot e^{i\theta\mathfrak{S}_\Gamma^F((j_1 \boxplus_3 \psi_1))} \geq \mathfrak{R}_\Gamma^F((j_2 \boxplus_3 \psi_2)) \cdot e^{i\theta\mathfrak{S}_\Gamma^F((j_2 \boxplus_3 \psi_2))}$ , then  $\mathfrak{R}_\Gamma^F((j_1 \boxplus_3 \psi_1)) \cdot e^{i\theta\mathfrak{S}_\Gamma^F((j_1 \boxplus_3 \psi_1))} \leq \max\{\mathfrak{R}_\Gamma^F(j_1) \cdot e^{i\theta\mathfrak{S}_\Gamma^F(j_1)}, \mathfrak{R}_\Gamma^F(\psi_1) \cdot e^{i\theta\mathfrak{S}_\Gamma^F(\psi_1)}\}$ .

So,  $\Gamma$  is a ComCNSBS of  $S$ .

**Theorem 3.11.** Suppose  $\Gamma$  is a subset of  $S$ . Then  $R = (\widehat{\mathfrak{R}_\Gamma^T} \cdot e^{i\theta\widehat{\mathfrak{S}_\Gamma^T}}, \widehat{\mathfrak{R}_\Gamma^I} \cdot e^{i\theta\widehat{\mathfrak{S}_\Gamma^I}}, \widehat{\mathfrak{R}_\Gamma^F} \cdot e^{i\theta\widehat{\mathfrak{S}_\Gamma^F}}, \mathfrak{R}_\Gamma^T \cdot e^{i\theta\mathfrak{S}_\Gamma^T}, \mathfrak{R}_\Gamma^I \cdot e^{i\theta\mathfrak{S}_\Gamma^I}, \mathfrak{R}_\Gamma^F \cdot e^{i\theta\mathfrak{S}_\Gamma^F})$  is a ComCNSBS of  $S$  if and only if  $\mathfrak{R}^{(\varphi, \varkappa)}$  is a SBS, where  $\varphi, \varkappa \in D[0, 1]$ .

**Proof.** Assume that  $S$  has a ComCNSBS in  $\widehat{\mathfrak{R}}$ . For all  $\varphi, \varkappa \in D[0, 1]$  and  $j_1, j_2 \in \widehat{\mathfrak{R}^{(\varphi, \varkappa)}}$ . Now,  $\widehat{\mathfrak{R}_\Gamma^T}(j_1) \cdot$

**Definition 3.12.** Let  $(S_1, \mathbb{U}_1, \mathbb{U}_2, \mathbb{U}_3)$  and  $(S_2, \mathbb{M}_1, \mathbb{M}_2, \mathbb{M}_3)$  be bisemirings and  $F : S_1 \rightarrow S_2$  and  $\Gamma$  be the ComCNSBS of  $S_1$ ,  $\Upsilon$  be the ComCNSBS of  $F(S_1) = S_2$ . If  $\mathfrak{R}_\Gamma \cdot e^{i\theta\mathfrak{Z}_\Gamma} = [\widehat{\mathfrak{R}}_\Gamma^T \cdot e^{i\theta\widehat{\mathfrak{Z}}_\Gamma^T}, \widehat{\mathfrak{R}}_\Gamma^I \cdot e^{i\theta\widehat{\mathfrak{Z}}_\Gamma^I}, \widehat{\mathfrak{R}}_\Gamma^F \cdot e^{i\theta\widehat{\mathfrak{Z}}_\Gamma^F}, \mathfrak{R}_\Gamma \cdot e^{i\theta\mathfrak{Z}_\Gamma}, \mathfrak{R}_\Gamma^T \cdot e^{i\theta\mathfrak{Z}_\Gamma^T}, \mathfrak{R}_\Gamma^I \cdot e^{i\theta\mathfrak{Z}_\Gamma^I}, \mathfrak{R}_\Gamma^F \cdot e^{i\theta\mathfrak{Z}_\Gamma^F}]$  is a ComCNS in  $S_1$ , then  $\mathfrak{R}_\Upsilon$  is a ComCNS in  $S_2$ , described as

$$\begin{aligned}\widehat{\mathfrak{R}}_{\Gamma}^T(\psi) \cdot e^{i\theta\widehat{\mathfrak{Z}}_{\Gamma}^T}(\psi) &= \begin{cases} \sup \widehat{\mathfrak{R}}_{\Gamma}^T(j) \cdot e^{i\theta\widehat{\mathfrak{Z}}_{\Gamma}^T}(j) & \text{if } j \in F^{-1}\psi \\ 0 & \text{otherwise} \end{cases} \\ \widehat{\mathfrak{R}}_{\Gamma}^I(\psi) \cdot e^{i\theta\widehat{\mathfrak{Z}}_{\Gamma}^I}(\psi) &= \begin{cases} \sup \widehat{\mathfrak{R}}_{\Gamma}^I(j) \cdot e^{i\theta\widehat{\mathfrak{Z}}_{\Gamma}^I}(j) & \text{if } j \in F^{-1}\psi \\ 0 & \text{otherwise} \end{cases} \\ \widehat{\mathfrak{R}}_{\Gamma}^F(\psi) \cdot e^{i\theta\widehat{\mathfrak{Z}}_{\Gamma}^F}(\psi) &= \begin{cases} \inf \widehat{\mathfrak{R}}_{\Gamma}^F(j) \cdot e^{i\theta\widehat{\mathfrak{Z}}_{\Gamma}^F}(j) & \text{if } j \in F^{-1}\psi \\ 1 & \text{otherwise} \end{cases}\end{aligned}$$

$$\begin{aligned}\mathfrak{R}_\Gamma^T(\psi) \cdot e^{i\theta\mathfrak{S}_\Gamma^T}(\psi) &= \begin{cases} \sup \mathfrak{R}_\Gamma^T(j) \cdot e^{i\theta\mathfrak{S}_\Gamma^T}(j) & \text{if } j \in F^{-1}\psi \\ 0 & \text{otherwise} \end{cases} \\ \mathfrak{R}_\Gamma^I(\psi) \cdot e^{i\theta\mathfrak{S}_\Gamma^I}(\psi) &= \begin{cases} \sup \mathfrak{R}_\Gamma^I(j) \cdot e^{i\theta\mathfrak{S}_\Gamma^I}(j) & \text{if } j \in F^{-1}\psi \\ 0 & \text{otherwise} \end{cases} \\ \mathfrak{R}_\Gamma^F(\psi) \cdot e^{i\theta\mathfrak{S}_\Gamma^F}(\psi) &= \begin{cases} \inf \mathfrak{R}_\Gamma^F(j) \cdot e^{i\theta\mathfrak{S}_\Gamma^F}(j) & \text{if } j \in F^{-1}\psi \\ 1 & \text{otherwise} \end{cases}\end{aligned}$$

for each  $j \in S_1$  and  $\psi \in S_2$  is said to be the image of  $R_\Gamma$  under  $F$ .

Similarly, If  $\mathfrak{R}_\Gamma \cdot e^{i\theta\mathfrak{S}_\Gamma} = [\widehat{\mathfrak{R}_\Gamma^T} \cdot e^{i\theta\widehat{\mathfrak{S}_\Gamma^T}}, \widehat{\mathfrak{R}_\Gamma^I} \cdot e^{i\theta\widehat{\mathfrak{S}_\Gamma^I}}, \widehat{\mathfrak{R}_\Gamma^F} \cdot e^{i\theta\widehat{\mathfrak{S}_\Gamma^F}}]$ ,  $\mathfrak{R}_\Gamma \cdot e^{i\theta\mathfrak{S}_\Gamma}$ ,  $\mathfrak{R}_\Gamma^T \cdot e^{i\theta\mathfrak{S}_\Gamma^T}$ ,  $\mathfrak{R}_\Gamma^I \cdot e^{i\theta\mathfrak{S}_\Gamma^I}$ ,  $\mathfrak{R}_\Gamma^F \cdot e^{i\theta\mathfrak{S}_\Gamma^F}$  is a ComCNS in  $S_2$ , then ComCNS  $\mathfrak{R}_\Gamma = F \circ \mathfrak{R}_\Gamma$  in  $S_1$  i.e., the ComCNS defined by  $\mathfrak{R}_\Gamma(j) \cdot e^{i\theta\mathfrak{S}_\Gamma(j)}$ ,  $\mathfrak{R}_\Gamma(F(j)) \cdot e^{i\theta\mathfrak{S}_\Gamma(F(j))}$ ,  $\mathfrak{R}_\Gamma = F \circ \mathfrak{R}_\Gamma$  in  $S_1$  [i.e., the ComCNS defined by  $\mathfrak{R}_\Gamma(j) \cdot e^{i\theta\mathfrak{S}_\Gamma(j)} = \mathfrak{R}_\Gamma(F(j)) \cdot e^{i\theta\mathfrak{S}_\Gamma(F(j))}$ ] is described the preimage of  $\mathfrak{R}_\Gamma$  under  $F$ .

**Theorem 3.13.** A ComCNSBS is the homomorphic image of every ComCNSBS.

**Proof.** Let  $F : S_1 \rightarrow S_2$  be any homomorphism. Now,  $F((j \uplus_1 \psi)) = F(j) \mathfrak{m}_1 F(\psi)$ ,  $F((j \uplus_2 \psi)) = F(j) \mathfrak{m}_2 F(\psi)$  and  $F((j \uplus_3 \psi)) = F(j) \mathfrak{m}_3 F(\psi)$  for all  $j, \psi \in S_1$ . Let  $\widehat{\Upsilon} = F(\Gamma)$ ,  $\Gamma$  is any ComCNSBS of  $S_1$ . Let  $F(j), F(\psi) \in S_2$ . Let  $j \in uF^{-1}(F(j))$  and  $\psi \in F^{-1}(F(\psi))$  be such that  $\widehat{\mathfrak{R}_\Gamma^T}(j) \cdot e^{i\theta\widehat{\mathfrak{S}_\Gamma^T}(j)} = \sup_{j \in F^{-1}(F(j))} \widehat{\mathfrak{R}_\Gamma^T}(j) \cdot e^{i\theta\widehat{\mathfrak{S}_\Gamma^T}(j)}$  and  $\widehat{\mathfrak{R}_\Gamma^T}(\psi) \cdot e^{i\theta\widehat{\mathfrak{S}_\Gamma^T}(\psi)} = \sup_{j \in F^{-1}(F(\psi))} \widehat{\mathfrak{R}_\Gamma^T}(j) \cdot e^{i\theta\widehat{\mathfrak{S}_\Gamma^T}(j)}$ . Now,

$$\begin{aligned}\widehat{\mathfrak{R}_\Gamma^T}((F(j) \mathfrak{m}_1 F(\psi))) \cdot e^{i\theta\widehat{\mathfrak{S}_\Gamma^T}((F(j) \mathfrak{m}_1 F(\psi)))} &= \sup_{(j') \in F^{-1}(F(j) \mathfrak{m}_1 F(\psi))} \widehat{\mathfrak{R}_\Gamma^T}(j') \cdot e^{i\theta\widehat{\mathfrak{S}_\Gamma^T}(j')} \\ &= \sup_{(j') \in F^{-1}(F((j \uplus_1 \psi)))} \widehat{\mathfrak{R}_\Gamma^T}(j') \cdot e^{i\theta\widehat{\mathfrak{S}_\Gamma^T}(j')} \\ &= \widehat{\mathfrak{R}_\Gamma^T}((j \uplus_1 \psi)) \cdot e^{i\theta\widehat{\mathfrak{S}_\Gamma^T}((j \uplus_1 \psi))} \\ &\geq \min\{\widehat{\mathfrak{R}_\Gamma^T}(j) \cdot e^{i\theta\widehat{\mathfrak{S}_\Gamma^T}(j)}, \widehat{\mathfrak{R}_\Gamma^T}(\psi) \cdot e^{i\theta\widehat{\mathfrak{S}_\Gamma^T}(\psi)}\} \\ &= \min\{\widehat{\mathfrak{R}_\Gamma^T}F(j) \cdot e^{i\theta\widehat{\mathfrak{S}_\Gamma^T}F(j)}, \widehat{\mathfrak{R}_\Gamma^T}F(\psi) \cdot e^{i\theta\widehat{\mathfrak{S}_\Gamma^T}F(\psi)}\}.\end{aligned}$$

Thus,  $\widehat{\mathfrak{R}_\Gamma^T}((F(j) \mathfrak{m}_1 F(\psi))) \cdot e^{i\theta\widehat{\mathfrak{S}_\Gamma^T}((F(j) \mathfrak{m}_1 F(\psi)))} \geq \min\{\widehat{\mathfrak{R}_\Gamma^T}F(j) \cdot e^{i\theta\widehat{\mathfrak{S}_\Gamma^T}F(j)}, \widehat{\mathfrak{R}_\Gamma^T}F(\psi) \cdot e^{i\theta\widehat{\mathfrak{S}_\Gamma^T}F(\psi)}\}$ .

Similarly,  $\widehat{\mathfrak{R}_\Gamma^T}((F(j) \mathfrak{m}_2 F(\psi))) \cdot e^{i\theta\widehat{\mathfrak{S}_\Gamma^T}((F(j) \mathfrak{m}_2 F(\psi)))} \geq \min\{\widehat{\mathfrak{R}_\Gamma^T}F(j) \cdot e^{i\theta\widehat{\mathfrak{S}_\Gamma^T}F(j)}, \widehat{\mathfrak{R}_\Gamma^T}F(\psi) \cdot e^{i\theta\widehat{\mathfrak{S}_\Gamma^T}F(\psi)}\}$  and  $\widehat{\mathfrak{R}_\Gamma^T}((F(j) \mathfrak{m}_3 F(\psi))) \cdot e^{i\theta\widehat{\mathfrak{S}_\Gamma^T}((F(j) \mathfrak{m}_3 F(\psi)))} \geq \min\{\widehat{\mathfrak{R}_\Gamma^T}F(j) \cdot e^{i\theta\widehat{\mathfrak{S}_\Gamma^T}F(j)}, \widehat{\mathfrak{R}_\Gamma^T}F(\psi) \cdot e^{i\theta\widehat{\mathfrak{S}_\Gamma^T}F(\psi)}\}$ .

Let  $j \in F^{-1}(F(j))$  and  $\psi \in F^{-1}(F(\psi))$  be such that  $\widehat{\mathfrak{R}_\Gamma^I}(j) \cdot e^{i\theta\widehat{\mathfrak{S}_\Gamma^I}(j)} = \sup_{j \in F^{-1}(F(j))} \widehat{\mathfrak{R}_\Gamma^I}(j) \cdot e^{i\theta\widehat{\mathfrak{S}_\Gamma^I}(j)}$  and

$\widehat{\mathfrak{R}_\Gamma^I}(\psi) \cdot e^{i\theta\widehat{\mathfrak{S}_\Gamma^I}(\psi)} = \sup_{j \in F^{-1}(F(\psi))} \widehat{\mathfrak{R}_\Gamma^I}(j) \cdot e^{i\theta\widehat{\mathfrak{S}_\Gamma^I}(j)}$ . Now,

$$\begin{aligned}\widehat{\mathfrak{R}_\Gamma^I}((F(j) \mathfrak{m}_1 F(\psi))) \cdot e^{i\theta\widehat{\mathfrak{S}_\Gamma^I}((F(j) \mathfrak{m}_1 F(\psi)))} &= \sup_{(j') \in F^{-1}(F(j) \mathfrak{m}_1 F(\psi))} \widehat{\mathfrak{R}_\Gamma^I}(j') \cdot e^{i\theta\widehat{\mathfrak{S}_\Gamma^I}(j')} \\ &= \sup_{(j') \in F^{-1}(F((j \uplus_1 \psi)))} \widehat{\mathfrak{R}_\Gamma^I}(j') \cdot e^{i\theta\widehat{\mathfrak{S}_\Gamma^I}(j')} \\ &= \widehat{\mathfrak{R}_\Gamma^I}((j \uplus_1 \psi)) \cdot e^{i\theta\widehat{\mathfrak{S}_\Gamma^I}((j \uplus_1 \psi))} \\ &\geq \frac{\widehat{\mathfrak{R}_\Gamma^I}(j) \cdot e^{i\theta\widehat{\mathfrak{S}_\Gamma^I}(j)} + \widehat{\mathfrak{R}_\Gamma^I}(\psi) \cdot e^{i\theta\widehat{\mathfrak{S}_\Gamma^I}(\psi)}}{2} \\ &= \frac{\widehat{\mathfrak{R}_\Gamma^I}F(j) \cdot e^{i\theta\widehat{\mathfrak{S}_\Gamma^I}F(j)} + \widehat{\mathfrak{R}_\Gamma^I}F(\psi) \cdot e^{i\theta\widehat{\mathfrak{S}_\Gamma^I}F(\psi)}}{2}.\end{aligned}$$

Thus,  $\widehat{\mathfrak{R}_\Gamma^I}((F(j) \mathfrak{m}_1 F(\psi))) \cdot e^{i\theta\widehat{\mathfrak{S}_\Gamma^I}((F(j) \mathfrak{m}_1 F(\psi)))} \geq \frac{\widehat{\mathfrak{R}_\Gamma^I}F(j) \cdot e^{i\theta\widehat{\mathfrak{S}_\Gamma^I}F(j)} + \widehat{\mathfrak{R}_\Gamma^I}F(\psi) \cdot e^{i\theta\widehat{\mathfrak{S}_\Gamma^I}F(\psi)}}{2}$ .

Similarly,  $\widehat{\mathfrak{R}_\Gamma^I}((F(j) \mathfrak{m}_2 F(\psi))) \cdot e^{i\theta\widehat{\mathfrak{S}_\Gamma^I}((F(j) \mathfrak{m}_2 F(\psi)))} \geq \frac{\widehat{\mathfrak{R}_\Gamma^I}F(j) \cdot e^{i\theta\widehat{\mathfrak{S}_\Gamma^I}F(j)} + \widehat{\mathfrak{R}_\Gamma^I}F(\psi) \cdot e^{i\theta\widehat{\mathfrak{S}_\Gamma^I}F(\psi)}}{2}$  and

$$\widehat{\mathfrak{R}}_{\Upsilon}^I((F(j) \mathfrak{M}_3 F(\psi))) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Upsilon}^I((F(j) \mathfrak{M}_3 F(\psi)))} \geq \frac{\widehat{\mathfrak{R}}_{\Upsilon}^I F(j) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Upsilon}^I F(j)} + \widehat{\mathfrak{R}}_{\Upsilon}^I F(\psi) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Upsilon}^I F(\psi)}}{2}.$$

Let  $F(j), F(\psi) \in S_2$ . Let  $j \in F^{-1}(F(j))$  and  $\psi \in F^{-1}(F(\psi))$  be such that  $\widehat{\mathfrak{R}}_{\Upsilon}^F(j) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Upsilon}^F(j)} = \inf_{j \in F^{-1}(F(j))} \widehat{\mathfrak{R}}_{\Upsilon}^F(j) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Upsilon}^F(j)}$  and  $\widehat{\mathfrak{R}}_{\Upsilon}^F(\psi) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Upsilon}^F(\psi)} = \inf_{\psi \in F^{-1}(F(\psi))} \widehat{\mathfrak{R}}_{\Upsilon}^F(\psi) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Upsilon}^F(\psi)}$ . Now,

$$\begin{aligned} \widehat{\mathfrak{R}}_{\Upsilon}^F((F(j) \mathfrak{M}_1 F(\psi))) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Upsilon}^F((F(j) \mathfrak{M}_1 F(\psi)))} &= \inf_{(j') \in F^{-1}(F(j) \mathfrak{M}_1 F(\psi))} \widehat{\mathfrak{R}}_{\Upsilon}^F(j') \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Upsilon}^F(j')} \\ &= \inf_{(j') \in F^{-1}(F((j \mathfrak{U}_1 \psi)))} \widehat{\mathfrak{R}}_{\Upsilon}^F(j') \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Upsilon}^F(j')} \\ &= \widehat{\mathfrak{R}}_{\Upsilon}^F((j \mathfrak{U}_1 \psi)) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Upsilon}^F((j \mathfrak{U}_1 \psi))} \\ &\leq \max\{\widehat{\mathfrak{R}}_{\Upsilon}^F(j) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Upsilon}^F(j)}, \widehat{\mathfrak{R}}_{\Upsilon}^F(\psi) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Upsilon}^F(\psi)}\} \\ &= \max\{\widehat{\mathfrak{R}}_{\Upsilon}^F F(j) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Upsilon}^F F(j)}, \widehat{\mathfrak{R}}_{\Upsilon}^F F(\psi) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Upsilon}^F F(\psi)}\}. \end{aligned}$$

Thus,  $\widehat{\mathfrak{R}}_{\Upsilon}^F((F(j) \mathfrak{M}_1 F(\psi))) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Upsilon}^F((F(j) \mathfrak{M}_1 F(\psi)))} \leq \max\{\widehat{\mathfrak{R}}_{\Upsilon}^F F(j) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Upsilon}^F F(j)}, \widehat{\mathfrak{R}}_{\Upsilon}^F F(\psi) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Upsilon}^F F(\psi)}\}$ .

Similarly,  $\widehat{\mathfrak{R}}_{\Upsilon}^F((F(j) \mathfrak{M}_2 F(\psi))) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Upsilon}^F((F(j) \mathfrak{M}_2 F(\psi)))} \leq \max\{\widehat{\mathfrak{R}}_{\Upsilon}^F F(j) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Upsilon}^F F(j)}, \widehat{\mathfrak{R}}_{\Upsilon}^F F(\psi) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Upsilon}^F F(\psi)}\}$  and  $\widehat{\mathfrak{R}}_{\Upsilon}^F((F(j) \mathfrak{M}_3 F(\psi))) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Upsilon}^F((F(j) \mathfrak{M}_3 F(\psi)))} \leq \max\{\widehat{\mathfrak{R}}_{\Upsilon}^F F(j) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Upsilon}^F F(j)}, \widehat{\mathfrak{R}}_{\Upsilon}^F F(\psi) \cdot e^{i\theta \widehat{\mathfrak{S}}_{\Upsilon}^F F(\psi)}\}$ .

Let  $F : S_1 \rightarrow S_2$  be any homomorphism. Now,  $F((j \mathfrak{U}_1 \psi)) = F(j) \mathfrak{M}_1 F(\psi)$ ,  $F((j \mathfrak{U}_2 \psi)) = F(j) \mathfrak{M}_2 F(\psi)$  and  $F((j \mathfrak{U}_3 \psi)) = F(j) \mathfrak{M}_3 F(\psi)$  for all  $j, \psi \in S_1$ . Let  $\Upsilon = F(\Gamma)$ ,  $\Gamma$  is any ComCNSBS of  $S_1$ . Let  $F(j), F(\psi) \in S_2$ . Let  $j \in F^{-1}(F(j))$  and  $\psi \in F^{-1}(F(\psi))$  be such that  $\mathfrak{R}_{\Gamma}^T(j) \cdot e^{i\theta \mathfrak{S}_{\Gamma}^T(j)} = \sup_{j \in F^{-1}(F(j))} \mathfrak{R}_{\Gamma}^T(j) \cdot e^{i\theta \mathfrak{S}_{\Gamma}^T(j)}$  and

$\mathfrak{R}_{\Gamma}^T(\psi) \cdot e^{i\theta \mathfrak{S}_{\Gamma}^T(\psi)} = \sup_{\psi \in F^{-1}(F(\psi))} \mathfrak{R}_{\Gamma}^T(\psi) \cdot e^{i\theta \mathfrak{S}_{\Gamma}^T(\psi)}$ . Now,

$$\begin{aligned} \mathfrak{R}_{\Upsilon}^T((F(j) \mathfrak{M}_1 F(\psi))) \cdot e^{i\theta \mathfrak{S}_{\Upsilon}^T((F(j) \mathfrak{M}_1 F(\psi)))} &= \sup_{(j') \in F^{-1}(F(j) \mathfrak{M}_1 F(\psi))} \mathfrak{R}_{\Upsilon}^T(j') \cdot e^{i\theta \mathfrak{S}_{\Upsilon}^T(j')} \\ &= \sup_{(j') \in F^{-1}(F((j \mathfrak{U}_1 \psi)))} \mathfrak{R}_{\Upsilon}^T(j') \cdot e^{i\theta \mathfrak{S}_{\Upsilon}^T(j')} \\ &= \mathfrak{R}_{\Upsilon}^T((j \mathfrak{U}_1 \psi)) \cdot e^{i\theta \mathfrak{S}_{\Upsilon}^T((j \mathfrak{U}_1 \psi))} \\ &\geq \min\{\mathfrak{R}_{\Gamma}^T(j) \cdot e^{i\theta \mathfrak{S}_{\Gamma}^T(j)}, \mathfrak{R}_{\Gamma}^T(\psi) \cdot e^{i\theta \mathfrak{S}_{\Gamma}^T(\psi)}\} \\ &= \min\{\mathfrak{R}_{\Upsilon}^T F(j) \cdot e^{i\theta \mathfrak{S}_{\Upsilon}^T F(j)}, \mathfrak{R}_{\Upsilon}^T F(\psi) \cdot e^{i\theta \mathfrak{S}_{\Upsilon}^T F(\psi)}\}. \end{aligned}$$

Thus,  $\mathfrak{R}_{\Upsilon}^T((F(j) \mathfrak{M}_1 F(\psi))) \cdot e^{i\theta \mathfrak{S}_{\Upsilon}^T((F(j) \mathfrak{M}_1 F(\psi)))} \geq \min\{\mathfrak{R}_{\Upsilon}^T F(j) \cdot e^{i\theta \mathfrak{S}_{\Upsilon}^T F(j)}, \mathfrak{R}_{\Upsilon}^T F(\psi) \cdot e^{i\theta \mathfrak{S}_{\Upsilon}^T F(\psi)}\}$ .

Similarly,  $\mathfrak{R}_{\Upsilon}^T((F(j) \mathfrak{M}_2 F(\psi))) \cdot e^{i\theta \mathfrak{S}_{\Upsilon}^T((F(j) \mathfrak{M}_2 F(\psi)))} \geq \min\{\mathfrak{R}_{\Upsilon}^T F(j) \cdot e^{i\theta \mathfrak{S}_{\Upsilon}^T F(j)}, \mathfrak{R}_{\Upsilon}^T F(\psi) \cdot e^{i\theta \mathfrak{S}_{\Upsilon}^T F(\psi)}\}$  and  $\mathfrak{R}_{\Upsilon}^T((F(j) \mathfrak{M}_3 F(\psi))) \cdot e^{i\theta \mathfrak{S}_{\Upsilon}^T((F(j) \mathfrak{M}_3 F(\psi)))} \geq \min\{\mathfrak{R}_{\Upsilon}^T F(j) \cdot e^{i\theta \mathfrak{S}_{\Upsilon}^T F(j)}, \mathfrak{R}_{\Upsilon}^T F(\psi) \cdot e^{i\theta \mathfrak{S}_{\Upsilon}^T F(\psi)}\}$ .

Let  $j \in F^{-1}(F(j))$  and  $\psi \in F^{-1}(F(\psi))$  be such that  $\mathfrak{R}_{\Gamma}^I(j) \cdot e^{i\theta \mathfrak{S}_{\Gamma}^I(j)} = \sup_{j \in F^{-1}(F(j))} \mathfrak{R}_{\Gamma}^I(j) \cdot e^{i\theta \mathfrak{S}_{\Gamma}^I(j)}$  and

$\mathfrak{R}_{\Gamma}^I(\psi) \cdot e^{i\theta \mathfrak{S}_{\Gamma}^I(\psi)} = \sup_{\psi \in F^{-1}(F(\psi))} \mathfrak{R}_{\Gamma}^I(\psi) \cdot e^{i\theta \mathfrak{S}_{\Gamma}^I(\psi)}$ . Now,

$$\begin{aligned} \mathfrak{R}_{\Upsilon}^I((F(j) \mathfrak{M}_1 F(\psi))) \cdot e^{i\theta \mathfrak{S}_{\Upsilon}^I((F(j) \mathfrak{M}_1 F(\psi)))} &= \sup_{(j') \in F^{-1}(F(j) \mathfrak{M}_1 F(\psi))} \mathfrak{R}_{\Upsilon}^I(j') \cdot e^{i\theta \mathfrak{S}_{\Upsilon}^I(j')} \\ &= \sup_{(j') \in F^{-1}(F((j \mathfrak{U}_1 \psi)))} \mathfrak{R}_{\Upsilon}^I(j') \cdot e^{i\theta \mathfrak{S}_{\Upsilon}^I(j')} \\ &= \mathfrak{R}_{\Upsilon}^I((j \mathfrak{U}_1 \psi)) \cdot e^{i\theta \mathfrak{S}_{\Upsilon}^I((j \mathfrak{U}_1 \psi))} \\ &\geq \frac{\mathfrak{R}_{\Gamma}^I(j) \cdot e^{i\theta \mathfrak{S}_{\Gamma}^I(j)} + \mathfrak{R}_{\Gamma}^I(\psi) \cdot e^{i\theta \mathfrak{S}_{\Gamma}^I(\psi)}}{2} \\ &= \frac{\mathfrak{R}_{\Upsilon}^I F(j) \cdot e^{i\theta \mathfrak{S}_{\Upsilon}^I F(j)} + \mathfrak{R}_{\Upsilon}^I F(\psi) \cdot e^{i\theta \mathfrak{S}_{\Upsilon}^I F(\psi)}}{2}. \end{aligned}$$

Thus,  $\mathfrak{R}_{\Upsilon}^I((F(j) \mathfrak{M}_1 F(\psi))) \cdot e^{i\theta \mathfrak{S}_{\Upsilon}^I((F(j) \mathfrak{M}_1 F(\psi)))} \geq \frac{\mathfrak{R}_{\Upsilon}^I F(j) \cdot e^{i\theta \mathfrak{S}_{\Upsilon}^I F(j)} + \mathfrak{R}_{\Upsilon}^I F(\psi) \cdot e^{i\theta \mathfrak{S}_{\Upsilon}^I F(\psi)}}{2}$ .

Similarly,  $\mathfrak{R}_{\Upsilon}^I((F(j) \mathfrak{M}_2 F(\psi))) \cdot e^{i\theta \mathfrak{S}_{\Upsilon}^I((F(j) \mathfrak{M}_2 F(\psi)))} \geq \frac{\mathfrak{R}_{\Upsilon}^I F(j) \cdot e^{i\theta \mathfrak{S}_{\Upsilon}^I F(j)} + \mathfrak{R}_{\Upsilon}^I F(\psi) \cdot e^{i\theta \mathfrak{S}_{\Upsilon}^I F(\psi)}}{2}$  and



$\mathfrak{R}_Y^I((F(j) \cap_3 F(\psi))) \cdot e^{i\theta \mathfrak{S}_Y^I((F(j) \cap_3 F(\psi)))} \geq \frac{\mathfrak{R}_Y^I(F(j)) \cdot e^{i\theta \mathfrak{S}_Y^I(F(j))} + \mathfrak{R}_Y^I(F(\psi)) \cdot e^{i\theta \mathfrak{S}_Y^I(F(\psi))}}{2}$ .  
 Let  $F(j), F(\psi) \in S_2$ . Let  $j \in F^{-1}(F(j))$  and  $\psi \in F^{-1}(F(\psi))$  be such that  $\mathfrak{R}_Y^F(j) \cdot e^{i\theta \mathfrak{S}_Y^F(j)} = \inf_{j \in F^{-1}(F(j))} \mathfrak{R}_Y^F(j) \cdot e^{i\theta \mathfrak{S}_Y^F(j)}$  and  $\mathfrak{R}_Y^F(\psi) \cdot e^{i\theta \mathfrak{S}_Y^F(\psi)} = \inf_{\psi \in F^{-1}(F(\psi))} \mathfrak{R}_Y^F(\psi) \cdot e^{i\theta \mathfrak{S}_Y^F(\psi)}$ . Now,

$$\begin{aligned} \mathfrak{R}_Y^F((F(j) \cap_1 F(\psi))) \cdot e^{i\theta \mathfrak{S}_Y^F((F(j) \cap_1 F(\psi)))} &= \inf_{(j') \in F^{-1}(F(j) \cap_1 F(\psi))} \mathfrak{R}_Y^F(j') \cdot e^{i\theta \mathfrak{S}_Y^F(j')} \\ &= \inf_{(j') \in F^{-1}(F((j \cup_1 \psi)))} \mathfrak{R}_Y^F(j') \cdot e^{i\theta \mathfrak{S}_Y^F(j')} \\ &= \mathfrak{R}_Y^F((j \cup_1 \psi)) \cdot e^{i\theta \mathfrak{S}_Y^F((j \cup_1 \psi))} \\ &\leq \max\{\mathfrak{R}_Y^F(j) \cdot e^{i\theta \mathfrak{S}_Y^F(j)}, \mathfrak{R}_Y^F(\psi) \cdot e^{i\theta \mathfrak{S}_Y^F(\psi)}\} \\ &= \max\{\mathfrak{R}_Y^F F(j) \cdot e^{i\theta \mathfrak{S}_Y^F F(j)}, \mathfrak{R}_Y^F F(\psi) \cdot e^{i\theta \mathfrak{S}_Y^F F(\psi)}\}. \end{aligned}$$

Thus,  $\mathfrak{R}_Y^F((F(j) \cap_1 F(\psi))) \cdot e^{i\theta \mathfrak{S}_Y^F((F(j) \cap_1 F(\psi)))} \leq \max\{\mathfrak{R}_Y^F F(j) \cdot e^{i\theta \mathfrak{S}_Y^F F(j)}, \mathfrak{R}_Y^F F(\psi) \cdot e^{i\theta \mathfrak{S}_Y^F F(\psi)}\}$ .  
 Similarly,  $\mathfrak{R}_Y^F((F(j) \cap_2 F(\psi))) \cdot e^{i\theta \mathfrak{S}_Y^F((F(j) \cap_2 F(\psi)))} \leq \max\{\mathfrak{R}_Y^F F(j) \cdot e^{i\theta \mathfrak{S}_Y^F F(j)}, \mathfrak{R}_Y^F F(\psi) \cdot e^{i\theta \mathfrak{S}_Y^F F(\psi)}\}$  and  $\mathfrak{R}_Y^F((F(j) \cap_3 F(\psi))) \cdot e^{i\theta \mathfrak{S}_Y^F((F(j) \cap_3 F(\psi)))} \leq \max\{\mathfrak{R}_Y^F F(j) \cdot e^{i\theta \mathfrak{S}_Y^F F(j)}, \mathfrak{R}_Y^F F(\psi) \cdot e^{i\theta \mathfrak{S}_Y^F F(\psi)}\}$ . Thus,  $Y$  is a ComCNSBS of  $S_2$ .

**Theorem 3.14.** A ComCNSBS is the homomorphic preimage of every ComCNSBS.

**Proof.** Let  $F : S_1 \rightarrow S_2$  be a homomorphism. Now,  $F((j \cup_1 \psi)) = F(j) \cap_1 F(\psi)$ ,  $F((j \cup_2 \psi)) = F(j) \cap_2 F(\psi)$  and  $F((j \cup_3 \psi)) = F(j) \cap_3 F(\psi)$  for all  $j, \psi \in S_1$ . Let  $\hat{Y} = F(Y)$ ,  $\hat{Y}$  is a ComCNSBS of  $S_2$ . Let  $j, \psi \in S_1$ . Now,  $\widehat{\mathfrak{R}}_Y^T((j \cup_1 \psi)) \cdot e^{i\theta \widehat{\mathfrak{S}}_Y^T((j \cup_1 \psi))} = \widehat{\mathfrak{R}}_Y^T(F((j \cup_1 \psi))) \cdot e^{i\theta \widehat{\mathfrak{S}}_Y^T(F((j \cup_1 \psi)))} = \widehat{\mathfrak{R}}_Y^T((F(j) \cap_1 F(\psi))) \cdot e^{i\theta \widehat{\mathfrak{S}}_Y^T((F(j) \cap_1 F(\psi)))} \geq \min\{\widehat{\mathfrak{R}}_Y^T F(j) \cdot e^{i\theta \widehat{\mathfrak{S}}_Y^T F(j)}, \widehat{\mathfrak{R}}_Y^T F(\psi) \cdot e^{i\theta \widehat{\mathfrak{S}}_Y^T F(\psi)}\} = \min\{\widehat{\mathfrak{R}}_Y^T(j) \cdot e^{i\theta \widehat{\mathfrak{S}}_Y^T(j)}, \widehat{\mathfrak{R}}_Y^T(\psi) \cdot e^{i\theta \widehat{\mathfrak{S}}_Y^T(\psi)}\}$ . Thus,  $\widehat{\mathfrak{R}}_Y^T((j \cup_1 \psi)) \cdot e^{i\theta \widehat{\mathfrak{S}}_Y^T((j \cup_1 \psi))} \geq \min\{\widehat{\mathfrak{R}}_Y^T(j) \cdot e^{i\theta \widehat{\mathfrak{S}}_Y^T(j)}, \widehat{\mathfrak{R}}_Y^T(\psi) \cdot e^{i\theta \widehat{\mathfrak{S}}_Y^T(\psi)}\}$ .  
 Now,  $\widehat{\mathfrak{R}}_Y^T((j \cup_1 \psi)) \cdot e^{i\theta \widehat{\mathfrak{S}}_Y^T((j \cup_1 \psi))} = \widehat{\mathfrak{R}}_Y^T(F((j \cup_1 \psi))) \cdot e^{i\theta \widehat{\mathfrak{S}}_Y^T(F((j \cup_1 \psi)))} = \widehat{\mathfrak{R}}_Y^T((F(j) \cap_1 F(\psi))) \cdot e^{i\theta \widehat{\mathfrak{S}}_Y^T((F(j) \cap_1 F(\psi)))} \geq \frac{\widehat{\mathfrak{R}}_Y^T F(j) \cdot e^{i\theta \widehat{\mathfrak{S}}_Y^T F(j)} + \widehat{\mathfrak{R}}_Y^T F(\psi) \cdot e^{i\theta \widehat{\mathfrak{S}}_Y^T F(\psi)}}{2} = \frac{\widehat{\mathfrak{R}}_Y^T(j) \cdot e^{i\theta \widehat{\mathfrak{S}}_Y^T(j)} + \widehat{\mathfrak{R}}_Y^T(\psi) \cdot e^{i\theta \widehat{\mathfrak{S}}_Y^T(\psi)}}{2}$ . Thus,  $\widehat{\mathfrak{R}}_Y^T((j \cup_1 \psi)) \cdot e^{i\theta \widehat{\mathfrak{S}}_Y^T((j \cup_1 \psi))} \geq \frac{\widehat{\mathfrak{R}}_Y^T(j) \cdot e^{i\theta \widehat{\mathfrak{S}}_Y^T(j)} + \widehat{\mathfrak{R}}_Y^T(\psi) \cdot e^{i\theta \widehat{\mathfrak{S}}_Y^T(\psi)}}{2}$ . Now,  $\widehat{\mathfrak{R}}_Y^F((j \cup_1 \psi)) \cdot e^{i\theta \widehat{\mathfrak{S}}_Y^F((j \cup_1 \psi))} = \widehat{\mathfrak{R}}_Y^F(F((j \cup_1 \psi))) \cdot e^{i\theta \widehat{\mathfrak{S}}_Y^F(F((j \cup_1 \psi)))} = \widehat{\mathfrak{R}}_Y^F((F(j) \cap_1 F(\psi))) \cdot e^{i\theta \widehat{\mathfrak{S}}_Y^F((F(j) \cap_1 F(\psi)))} \leq \max\{\widehat{\mathfrak{R}}_Y^F F(j) \cdot e^{i\theta \widehat{\mathfrak{S}}_Y^F F(j)}, \widehat{\mathfrak{R}}_Y^F F(\psi) \cdot e^{i\theta \widehat{\mathfrak{S}}_Y^F F(\psi)}\} = \max\{\widehat{\mathfrak{R}}_Y^F(j) \cdot e^{i\theta \widehat{\mathfrak{S}}_Y^F(j)}, \widehat{\mathfrak{R}}_Y^F(\psi) \cdot e^{i\theta \widehat{\mathfrak{S}}_Y^F(\psi)}\}$ . Thus,  $\widehat{\mathfrak{R}}_Y^F((j \cup_1 \psi)) \cdot e^{i\theta \widehat{\mathfrak{S}}_Y^F((j \cup_1 \psi))} \leq \max\{\widehat{\mathfrak{R}}_Y^F(j) \cdot e^{i\theta \widehat{\mathfrak{S}}_Y^F(j)}, \widehat{\mathfrak{R}}_Y^F(\psi) \cdot e^{i\theta \widehat{\mathfrak{S}}_Y^F(\psi)}\}$ .

Let  $F : S_1 \rightarrow S_2$  be a homomorphism. Now,  $F((j \cup_1 \psi)) = F(j) \cap_1 F(\psi)$ ,  $F((j \cup_2 \psi)) = F(j) \cap_2 F(\psi)$  and  $F((j \cup_3 \psi)) = F(j) \cap_3 F(\psi)$  for all  $j, \psi \in S_1$ . Let  $Y = F(Y)$ ,  $Y$  is a ComCNSBS of  $S_2$ . Let  $j, \psi \in S_1$ . Now,  $\mathfrak{R}_Y^T((j \cup_1 \psi)) \cdot e^{i\theta \mathfrak{S}_Y^T((j \cup_1 \psi))} = \mathfrak{R}_Y^T(F((j \cup_1 \psi))) \cdot e^{i\theta \mathfrak{S}_Y^T(F((j \cup_1 \psi)))} = \mathfrak{R}_Y^T((F(j) \cap_1 F(\psi))) \cdot e^{i\theta \mathfrak{S}_Y^T((F(j) \cap_1 F(\psi)))} \geq \min\{\mathfrak{R}_Y^T F(j) \cdot e^{i\theta \mathfrak{S}_Y^T F(j)}, \mathfrak{R}_Y^T F(\psi) \cdot e^{i\theta \mathfrak{S}_Y^T F(\psi)}\} = \min\{\mathfrak{R}_Y^T(j) \cdot e^{i\theta \mathfrak{S}_Y^T(j)}, \mathfrak{R}_Y^T(\psi) \cdot e^{i\theta \mathfrak{S}_Y^T(\psi)}\}$ . Thus,  $\mathfrak{R}_Y^T((j \cup_1 \psi)) \cdot e^{i\theta \mathfrak{S}_Y^T((j \cup_1 \psi))} \geq \min\{\mathfrak{R}_Y^T(j) \cdot e^{i\theta \mathfrak{S}_Y^T(j)}, \mathfrak{R}_Y^T(\psi) \cdot e^{i\theta \mathfrak{S}_Y^T(\psi)}\}$ .  
 Now,  $\mathfrak{R}_Y^T((j \cup_1 \psi)) \cdot e^{i\theta \mathfrak{S}_Y^T((j \cup_1 \psi))} = \mathfrak{R}_Y^T(F((j \cup_1 \psi))) \cdot e^{i\theta \mathfrak{S}_Y^T(F((j \cup_1 \psi)))} = \mathfrak{R}_Y^T((F(j) \cap_1 F(\psi))) \cdot e^{i\theta \mathfrak{S}_Y^T((F(j) \cap_1 F(\psi)))} \geq \frac{\mathfrak{R}_Y^T F(j) \cdot e^{i\theta \mathfrak{S}_Y^T F(j)} + \mathfrak{R}_Y^T F(\psi) \cdot e^{i\theta \mathfrak{S}_Y^T F(\psi)}}{2} = \frac{\mathfrak{R}_Y^T(j) \cdot e^{i\theta \mathfrak{S}_Y^T(j)} + \mathfrak{R}_Y^T(\psi) \cdot e^{i\theta \mathfrak{S}_Y^T(\psi)}}{2}$ . Thus,  $\mathfrak{R}_Y^T((j \cup_1 \psi)) \cdot e^{i\theta \mathfrak{S}_Y^T((j \cup_1 \psi))} \geq \frac{\mathfrak{R}_Y^T(j) \cdot e^{i\theta \mathfrak{S}_Y^T(j)} + \mathfrak{R}_Y^T(\psi) \cdot e^{i\theta \mathfrak{S}_Y^T(\psi)}}{2}$ . Now,  $\mathfrak{R}_Y^F((j \cup_1 \psi)) \cdot e^{i\theta \mathfrak{S}_Y^F((j \cup_1 \psi))} = \mathfrak{R}_Y^F(F((j \cup_1 \psi))) \cdot e^{i\theta \mathfrak{S}_Y^F(F((j \cup_1 \psi)))} = \mathfrak{R}_Y^F((F(j) \cap_1 F(\psi))) \cdot e^{i\theta \mathfrak{S}_Y^F((F(j) \cap_1 F(\psi)))} \leq \max\{\mathfrak{R}_Y^F F(j) \cdot e^{i\theta \mathfrak{S}_Y^F F(j)}, \mathfrak{R}_Y^F F(\psi) \cdot e^{i\theta \mathfrak{S}_Y^F F(\psi)}\} = \max\{\mathfrak{R}_Y^F(j) \cdot e^{i\theta \mathfrak{S}_Y^F(j)}, \mathfrak{R}_Y^F(\psi) \cdot e^{i\theta \mathfrak{S}_Y^F(\psi)}\}$ . Thus,  $\mathfrak{R}_Y^F((j \cup_1 \psi)) \cdot e^{i\theta \mathfrak{S}_Y^F((j \cup_1 \psi))} \leq \max\{\mathfrak{R}_Y^F(j) \cdot e^{i\theta \mathfrak{S}_Y^F(j)}, \mathfrak{R}_Y^F(\psi) \cdot e^{i\theta \mathfrak{S}_Y^F(\psi)}\}$ .

**Theorem 3.15.** If  $F : S_1 \rightarrow S_2$  is a homomorphism, then  $F(\widehat{\Gamma}_{(\varphi, \varkappa)})$  is a SBS of ComCNSBS  $\hat{Y}$  of  $S_2$ .

**Proof.** The mapping  $F : S_1 \rightarrow S_2$  be a homomorphism. Now,  $F((j \cup_1 \psi)) = F(j) \cap_1 F(\psi)$ ,  $F((j \cup_2 \psi)) = F(j) \cap_2 F(\psi)$  and  $F((j \cup_3 \psi)) = F(j) \cap_3 F(\psi)$  for all  $j, \psi \in S_1$ . Let  $\hat{Y} = F(Y)$ ,  $\hat{Y}$  is a ComCNSBS of  $S_2$ . By Theorem 3.13,  $\hat{\Gamma}_{(\varphi, \varkappa)}$  be any SBS of  $\Gamma$ . Suppose that  $j, \psi \in \widehat{\Gamma}_{(\varphi, \varkappa)}$ . Then  $j \cup_1 \psi, j \cup_2 \psi$  and  $j \cup_3 \psi \in \widehat{\Gamma}_{(\varphi, \varkappa)}$ . Now,  $\widehat{\mathfrak{R}}_Y^T(F(j)) \cdot e^{i\theta \widehat{\mathfrak{S}}_Y^T(F(j))} = \widehat{\mathfrak{R}}_Y^T(j) \cdot e^{i\theta \widehat{\mathfrak{S}}_Y^T(j)} \geq \varnothing$ ,  $\widehat{\mathfrak{R}}_Y^T(F(\psi)) \cdot e^{i\theta \widehat{\mathfrak{S}}_Y^T(F(\psi))} = \widehat{\mathfrak{R}}_Y^T(\psi) \cdot e^{i\theta \widehat{\mathfrak{S}}_Y^T(\psi)} \geq \varnothing$ . Thus,  $\widehat{\mathfrak{R}}_Y^T((F(j) \cap_1 F(\psi))) \cdot e^{i\theta \widehat{\mathfrak{S}}_Y^T((F(j) \cap_1 F(\psi)))} \geq \widehat{\mathfrak{R}}_Y^T((j \cup_1 \psi)) \cdot e^{i\theta \widehat{\mathfrak{S}}_Y^T((j \cup_1 \psi))} \geq \varnothing$ .

Now,  $\widehat{\mathfrak{R}}_{\Upsilon}^I(F(j)) \cdot e^{i\theta\widehat{\mathfrak{S}}_{\Upsilon}^I(F(j))} = \widehat{\mathfrak{R}}_{\Upsilon}^I(j) \cdot e^{i\theta\widehat{\mathfrak{S}}_{\Upsilon}^I(j)} \geq \wp$ ,  $\widehat{\mathfrak{R}}_{\Upsilon}^I(F(\psi)) \cdot e^{i\theta\widehat{\mathfrak{S}}_{\Upsilon}^I(F(\psi))} = \widehat{\mathfrak{R}}_{\Upsilon}^I(\psi) \cdot e^{i\theta\widehat{\mathfrak{S}}_{\Upsilon}^I(\psi)} \geq \wp$ . Thus,  $\widehat{\mathfrak{R}}_{\Upsilon}^I((F(j) \mathfrak{M}_1 F(\psi))) \cdot e^{i\theta\widehat{\mathfrak{S}}_{\Upsilon}^I((F(j) \mathfrak{M}_1 F(\psi)))} \geq \widehat{\mathfrak{R}}_{\Upsilon}^I((j \mathfrak{U}_1 \psi)) \cdot e^{i\theta\widehat{\mathfrak{S}}_{\Upsilon}^I((j \mathfrak{U}_1 \psi))} \geq \wp$ . Now,  $\widehat{\mathfrak{R}}_{\Upsilon}^F(F(j)) \cdot e^{i\theta\widehat{\mathfrak{S}}_{\Upsilon}^F(F(j))} = \widehat{\mathfrak{R}}_{\Upsilon}^F(j) \cdot e^{i\theta\widehat{\mathfrak{S}}_{\Upsilon}^F(j)} \leq \varkappa$ ,  $\widehat{\mathfrak{R}}_{\Upsilon}^F(F(\psi)) \cdot e^{i\theta\widehat{\mathfrak{S}}_{\Upsilon}^F(F(\psi))} = \widehat{\mathfrak{R}}_{\Upsilon}^F(\psi) \cdot e^{i\theta\widehat{\mathfrak{S}}_{\Upsilon}^F(\psi)} \leq \varkappa$ . Thus,  $\widehat{\mathfrak{R}}_{\Upsilon}^F((F(j) \mathfrak{M}_1 F(\psi))) \cdot e^{i\theta\widehat{\mathfrak{S}}_{\Upsilon}^F((F(j) \mathfrak{M}_1 F(\psi)))} \leq \widehat{\mathfrak{R}}_{\Upsilon}^F((j \mathfrak{U}_1 \psi)) \cdot e^{i\theta\widehat{\mathfrak{S}}_{\Upsilon}^F((j \mathfrak{U}_1 \psi))} \leq \varkappa$ , for all  $F(j), F(\psi) \in S_2$ . Similarly other operations,  $F(\widehat{\Gamma}_{(\wp, \varkappa)})$  is a SBS of ComCNSBS  $\widehat{\Upsilon}$  of  $S_2$ .

Let  $F : S_1 \rightarrow S_2$  be a homomorphism. Now,  $F((j \mathfrak{U}_1 \psi)) = F(j) \mathfrak{M}_1 F(\psi)$ ,  $F((j \mathfrak{U}_2 \psi)) = F(j) \mathfrak{M}_2 F(\psi)$  and  $F((j \mathfrak{U}_3 \psi)) = F(j) \mathfrak{M}_3 F(\psi)$  for all  $j, \psi \in S_1$ . Let  $\Upsilon = F(\Gamma)$ ,  $\Gamma$  is a ComCNSBS of  $S_1$ . By Theorem 3.13,  $\Upsilon$  is a ComCNSBS of  $S_2$ . Let  $\Gamma_{(\wp, \varkappa)}$  be any SBS of  $\Gamma$ . Suppose that  $j, \psi \in \Gamma_{(\wp, \varkappa)}$ . Then  $j \mathfrak{U}_1 \psi, j \mathfrak{U}_2 \psi$  and  $j \mathfrak{U}_3 \psi \in \Gamma_{(\wp, \varkappa)}$ . Now,  $\mathfrak{R}_{\Upsilon}^T(F(j)) \cdot e^{i\theta\mathfrak{S}_{\Upsilon}^T(F(j))} = \mathfrak{R}_{\Upsilon}^T(j) \cdot e^{i\theta\mathfrak{S}_{\Upsilon}^T(j)} \geq \wp$ ,  $\mathfrak{R}_{\Upsilon}^T(F(\psi)) \cdot e^{i\theta\mathfrak{S}_{\Upsilon}^T(F(\psi))} = \mathfrak{R}_{\Upsilon}^T(\psi) \cdot e^{i\theta\mathfrak{S}_{\Upsilon}^T(\psi)} \geq \wp$ . Thus,  $\mathfrak{R}_{\Upsilon}^T((F(j) \mathfrak{M}_1 F(\psi))) \cdot e^{i\theta\mathfrak{S}_{\Upsilon}^T((F(j) \mathfrak{M}_1 F(\psi)))} \geq \mathfrak{R}_{\Upsilon}^T((j \mathfrak{U}_1 \psi)) \cdot e^{i\theta\mathfrak{S}_{\Upsilon}^T((j \mathfrak{U}_1 \psi))} \geq \wp$ . Now,  $\mathfrak{R}_{\Upsilon}^I(F(j)) \cdot e^{i\theta\mathfrak{S}_{\Upsilon}^I(F(j))} = \mathfrak{R}_{\Upsilon}^I(j) \cdot e^{i\theta\mathfrak{S}_{\Upsilon}^I(j)} \geq \wp$ ,  $\mathfrak{R}_{\Upsilon}^I(F(\psi)) \cdot e^{i\theta\mathfrak{S}_{\Upsilon}^I(F(\psi))} = \mathfrak{R}_{\Upsilon}^I(\psi) \cdot e^{i\theta\mathfrak{S}_{\Upsilon}^I(\psi)} \geq \wp$ . Thus,  $\mathfrak{R}_{\Upsilon}^I((F(j) \mathfrak{M}_1 F(\psi))) \cdot e^{i\theta\mathfrak{S}_{\Upsilon}^I((F(j) \mathfrak{M}_1 F(\psi)))} \geq \mathfrak{R}_{\Upsilon}^I((j \mathfrak{U}_1 \psi)) \cdot e^{i\theta\mathfrak{S}_{\Upsilon}^I((j \mathfrak{U}_1 \psi))} \geq \wp$ . Now,  $\mathfrak{R}_{\Upsilon}^F(F(j)) \cdot e^{i\theta\mathfrak{S}_{\Upsilon}^F(F(j))} = \mathfrak{R}_{\Upsilon}^F(j) \cdot e^{i\theta\mathfrak{S}_{\Upsilon}^F(j)} \leq \varkappa$ ,  $\mathfrak{R}_{\Upsilon}^F(F(\psi)) \cdot e^{i\theta\mathfrak{S}_{\Upsilon}^F(F(\psi))} = \mathfrak{R}_{\Upsilon}^F(\psi) \cdot e^{i\theta\mathfrak{S}_{\Upsilon}^F(\psi)} \leq \varkappa$ . Thus,  $\mathfrak{R}_{\Upsilon}^F((F(j) \mathfrak{M}_1 F(\psi))) \cdot e^{i\theta\mathfrak{S}_{\Upsilon}^F((F(j) \mathfrak{M}_1 F(\psi)))} \leq \mathfrak{R}_{\Upsilon}^F((j \mathfrak{U}_1 \psi)) \cdot e^{i\theta\mathfrak{S}_{\Upsilon}^F((j \mathfrak{U}_1 \psi))} \leq \varkappa$ , for all  $F(j), F(\psi) \in S_2$ . Similarly other operations,  $F(\Gamma_{(\wp, \varkappa)})$  is a SBS of ComCNSBS  $\Upsilon$  of  $S_2$ .

**Theorem 3.16.** If  $F : S_1 \rightarrow S_2$  is any homomorphism, then  $\widehat{\Gamma}_{(\wp, \varkappa)}$  is a SBS of ComCNSBS  $\Gamma$  of  $S_1$ .

**Proof.** Let  $F : S_1 \rightarrow S_2$  be any homomorphism. We have  $F((j \mathfrak{U}_1 \psi)) = F(j) \mathfrak{M}_1 F(\psi)$ ,  $F((j \mathfrak{U}_2 \psi)) = F(j) \mathfrak{M}_2 F(\psi)$  and  $F((j \mathfrak{U}_3 \psi)) = F(j) \mathfrak{M}_3 F(\psi)$  for all  $j, \psi \in S_1$ . Let  $\widehat{\Upsilon} = F(\widehat{\Gamma})$ ,  $\widehat{\Upsilon}$  is a ComCNSBS of  $S_2$ . By Theorem 3.14,  $\Gamma$  is a ComCNSBS of  $S_1$ . Let  $F(\widehat{\Gamma}_{(\wp, \varkappa)})$  be a SBS of  $\widehat{\Upsilon}$ . Suppose that  $F(j), F(\psi) \in F(\widehat{\Gamma}_{(\wp, \varkappa)})$ . Now,  $F((j \mathfrak{U}_1 \psi)), F((j \mathfrak{U}_2 \psi))$  and  $F((j \mathfrak{U}_3 \psi)) \in F(\widehat{\Gamma}_{(\wp, \varkappa)})$ . Now,  $\widehat{\mathfrak{R}}_{\Upsilon}^T(j) \cdot e^{i\theta\widehat{\mathfrak{S}}_{\Upsilon}^T(j)} = \widehat{\mathfrak{R}}_{\Upsilon}^T(F(j)) \cdot e^{i\theta\widehat{\mathfrak{S}}_{\Upsilon}^T(F(j))} \geq \wp$ ,  $\widehat{\mathfrak{R}}_{\Upsilon}^T(\psi) \cdot e^{i\theta\widehat{\mathfrak{S}}_{\Upsilon}^T(\psi)} = \widehat{\mathfrak{R}}_{\Upsilon}^T(F(\psi)) \cdot e^{i\theta\widehat{\mathfrak{S}}_{\Upsilon}^T(F(\psi))} \geq \wp$ . Thus,  $\widehat{\mathfrak{R}}_{\Upsilon}^T((j \mathfrak{U}_1 \psi)) \cdot e^{i\theta\widehat{\mathfrak{S}}_{\Upsilon}^T((j \mathfrak{U}_1 \psi))} \geq \min\{\widehat{\mathfrak{R}}_{\Upsilon}^T(j) \cdot e^{i\theta\widehat{\mathfrak{S}}_{\Upsilon}^T(j)}, \widehat{\mathfrak{R}}_{\Upsilon}^T(\psi) \cdot e^{i\theta\widehat{\mathfrak{S}}_{\Upsilon}^T(\psi)}\} \geq \wp$ . Now,  $\widehat{\mathfrak{R}}_{\Upsilon}^I(j) \cdot e^{i\theta\widehat{\mathfrak{S}}_{\Upsilon}^I(j)} = \widehat{\mathfrak{R}}_{\Upsilon}^I(F(j)) \cdot e^{i\theta\widehat{\mathfrak{S}}_{\Upsilon}^I(F(j))} \geq \wp$ ,  $\widehat{\mathfrak{R}}_{\Upsilon}^I(\psi) \cdot e^{i\theta\widehat{\mathfrak{S}}_{\Upsilon}^I(\psi)} = \widehat{\mathfrak{R}}_{\Upsilon}^I(F(\psi)) \cdot e^{i\theta\widehat{\mathfrak{S}}_{\Upsilon}^I(F(\psi))} \geq \wp$ . Thus,  $\widehat{\mathfrak{R}}_{\Upsilon}^I((j \mathfrak{U}_1 \psi)) \cdot e^{i\theta\widehat{\mathfrak{S}}_{\Upsilon}^I((j \mathfrak{U}_1 \psi))} \geq \min\{\widehat{\mathfrak{R}}_{\Upsilon}^I(j) \cdot e^{i\theta\widehat{\mathfrak{S}}_{\Upsilon}^I(j)}, \widehat{\mathfrak{R}}_{\Upsilon}^I(\psi) \cdot e^{i\theta\widehat{\mathfrak{S}}_{\Upsilon}^I(\psi)}\} \geq \wp$ . Now,  $\widehat{\mathfrak{R}}_{\Upsilon}^F(j) \cdot e^{i\theta\widehat{\mathfrak{S}}_{\Upsilon}^F(j)} = \widehat{\mathfrak{R}}_{\Upsilon}^F(F(j)) \cdot e^{i\theta\widehat{\mathfrak{S}}_{\Upsilon}^F(F(j))} \leq \varkappa$ ,  $\widehat{\mathfrak{R}}_{\Upsilon}^F(\psi) \cdot e^{i\theta\widehat{\mathfrak{S}}_{\Upsilon}^F(\psi)} = \widehat{\mathfrak{R}}_{\Upsilon}^F(F(\psi)) \cdot e^{i\theta\widehat{\mathfrak{S}}_{\Upsilon}^F(F(\psi))} \leq \varkappa$ . Thus,  $\widehat{\mathfrak{R}}_{\Upsilon}^F((j \mathfrak{U}_1 \psi)) \cdot e^{i\theta\widehat{\mathfrak{S}}_{\Upsilon}^F((j \mathfrak{U}_1 \psi))} = \widehat{\mathfrak{R}}_{\Upsilon}^F((F(j) \mathfrak{M}_1 F(\psi))) \cdot e^{i\theta\widehat{\mathfrak{S}}_{\Upsilon}^F((F(j) \mathfrak{M}_1 F(\psi)))} \leq \max\{\widehat{\mathfrak{R}}_{\Upsilon}^F(j) \cdot e^{i\theta\widehat{\mathfrak{S}}_{\Upsilon}^F(j)}, \widehat{\mathfrak{R}}_{\Upsilon}^F(\psi) \cdot e^{i\theta\widehat{\mathfrak{S}}_{\Upsilon}^F(\psi)}\} \leq \varkappa$ , for all  $j, \psi \in S_1$ . Similarly other operations,  $\widehat{\Gamma}_{(\wp, \varkappa)}$  is a SBS of ComCNSBS  $\Gamma$  of  $S_1$ .

Let  $F : S_1 \rightarrow S_2$  be any homomorphism. We have  $F((j \mathfrak{U}_1 \psi)) = F(j) \mathfrak{M}_1 F(\psi)$ ,  $F((j \mathfrak{U}_2 \psi)) = F(j) \mathfrak{M}_2 F(\psi)$  and  $F((j \mathfrak{U}_3 \psi)) = F(j) \mathfrak{M}_3 F(\psi)$  for all  $j, \psi \in S_1$ . Let  $\Upsilon = F(\Gamma)$ ,  $\Upsilon$  is a ComCNSBS of  $S_2$ . By Theorem 3.14,  $\Gamma$  is a ComCNSBS of  $S_1$ . Let  $F(\Gamma_{(\wp, \varkappa)})$  be a SBS of  $\Upsilon$ . Suppose that  $F(j), F(\psi) \in F(\Gamma_{(\wp, \varkappa)})$ . Now,  $F((j \mathfrak{U}_1 \psi)), F((j \mathfrak{U}_2 \psi))$  and  $F((j \mathfrak{U}_3 \psi)) \in F(\Gamma_{(\wp, \varkappa)})$ . Now,  $\mathfrak{R}_{\Upsilon}^T(j) \cdot e^{i\theta\mathfrak{S}_{\Upsilon}^T(j)} = \mathfrak{R}_{\Upsilon}^T(F(j)) \cdot e^{i\theta\mathfrak{S}_{\Upsilon}^T(F(j))} \geq \wp$ ,  $\mathfrak{R}_{\Upsilon}^T(\psi) \cdot e^{i\theta\mathfrak{S}_{\Upsilon}^T(\psi)} = \mathfrak{R}_{\Upsilon}^T(F(\psi)) \cdot e^{i\theta\mathfrak{S}_{\Upsilon}^T(F(\psi))} \geq \wp$ . Thus,  $\mathfrak{R}_{\Upsilon}^T((j \mathfrak{U}_1 \psi)) \cdot e^{i\theta\mathfrak{S}_{\Upsilon}^T((j \mathfrak{U}_1 \psi))} \geq \min\{\mathfrak{R}_{\Upsilon}^T(j) \cdot e^{i\theta\mathfrak{S}_{\Upsilon}^T(j)}, \mathfrak{R}_{\Upsilon}^T(\psi) \cdot e^{i\theta\mathfrak{S}_{\Upsilon}^T(\psi)}\} \geq \wp$ . Now,  $\mathfrak{R}_{\Upsilon}^I(j) \cdot e^{i\theta\mathfrak{S}_{\Upsilon}^I(j)} = \mathfrak{R}_{\Upsilon}^I(F(j)) \cdot e^{i\theta\mathfrak{S}_{\Upsilon}^I(F(j))} \geq \wp$ ,  $\mathfrak{R}_{\Upsilon}^I(\psi) \cdot e^{i\theta\mathfrak{S}_{\Upsilon}^I(\psi)} = \mathfrak{R}_{\Upsilon}^I(F(\psi)) \cdot e^{i\theta\mathfrak{S}_{\Upsilon}^I(F(\psi))} \geq \wp$ . Thus,  $\mathfrak{R}_{\Upsilon}^I((j \mathfrak{U}_1 \psi)) \cdot e^{i\theta\mathfrak{S}_{\Upsilon}^I((j \mathfrak{U}_1 \psi))} \geq \frac{\mathfrak{R}_{\Upsilon}^I(j) \cdot e^{i\theta\mathfrak{S}_{\Upsilon}^I(j)} + \mathfrak{R}_{\Upsilon}^I(\psi) \cdot e^{i\theta\mathfrak{S}_{\Upsilon}^I(\psi)}}{2} \geq \wp$ . Now,  $\mathfrak{R}_{\Upsilon}^F(j) \cdot e^{i\theta\mathfrak{S}_{\Upsilon}^F(j)} = \mathfrak{R}_{\Upsilon}^F(F(j)) \cdot e^{i\theta\mathfrak{S}_{\Upsilon}^F(F(j))} \leq \varkappa$ ,  $\mathfrak{R}_{\Upsilon}^F(\psi) \cdot e^{i\theta\mathfrak{S}_{\Upsilon}^F(\psi)} = \mathfrak{R}_{\Upsilon}^F(F(\psi)) \cdot e^{i\theta\mathfrak{S}_{\Upsilon}^F(F(\psi))} \leq \varkappa$ . Thus,  $\mathfrak{R}_{\Upsilon}^F((j \mathfrak{U}_1 \psi)) \cdot e^{i\theta\mathfrak{S}_{\Upsilon}^F((j \mathfrak{U}_1 \psi))} = \mathfrak{R}_{\Upsilon}^F((F(j) \mathfrak{M}_1 F(\psi))) \cdot e^{i\theta\mathfrak{S}_{\Upsilon}^F((F(j) \mathfrak{M}_1 F(\psi)))} \leq \max\{\mathfrak{R}_{\Upsilon}^F(j) \cdot e^{i\theta\mathfrak{S}_{\Upsilon}^F(j)}, \mathfrak{R}_{\Upsilon}^F(\psi) \cdot e^{i\theta\mathfrak{S}_{\Upsilon}^F(\psi)}\} \leq \varkappa$ , for all  $j, \psi \in S_1$ . Similarly other operations,  $\Gamma_{(\wp, \varkappa)}$  is a SBS of ComCNSBS  $\Gamma$  of  $S_1$ .

#### 4 Conclusion and future direction

A novel kind of ComCNSBS is presented in this work. The concept of three grades is approached in a novel way by the complex cubic neutrosophic SBS, which describes three grades in terms of a complex number. A complicated form of three grades signifies a paradigm change since it converts three grades into a two dimensional parameter. Neutrosophic SBS with complicated interval values was defined. The concepts of

ComCNNSBS and ComCNSBS level sets were established. Applying the cubic set and anti cubic set to bisemiring is our aim. Additionally, a study of the characteristics of different conversions is done. In addition to data analysis, we are trying to handle images and signals, thus it makes sense to use F-transforms to expand the use of new fuzzy structures. Thus, we ought to consider using cubic SBS, soft set ComCNSBS, and soft set ComCNNSBS in the future.

## References

- [1] L. A. Zadeh, Fuzzy sets, *Information and Control*, 8, (1965), 338-353.
- [2] K. Atanassov, Intuitionistic fuzzy sets, *Fuzzy Sets and Systems*, 20(1), (1986) 87-96.
- [3] R. R. Yager, Pythagorean membership grades in multi criteria decision-making, *IEEE Trans. Fuzzy Systems*, 22, (2014), 958-965.
- [4] S. Ashraf, S. Abdullah, T. Mahmood, F. Ghani and T. Mahmood, Spherical fuzzy sets and their applications in multi-attribute decision making problems, *Journal of Intelligent and Fuzzy Systems*, 36, (2019), 2829-284.
- [5] B.C. Cuong and V. Kreinovich, Picture fuzzy sets a new concept for computational intelligence problems, in *Proceedings of 2013 Third World Congress on Information and Communication Technologies (WICT 2013)*, IEEE, (2013), 1-6.
- [6] M Palanikumar, K Arulmozhi, Novel possibility Pythagorean interval valued fuzzy soft set method for a decision making, *TWMS J. App. and Eng. Math.*, 13(1), (2023), 327–340.
- [7] Ali, O. Mashhadani, S. Alhakam, I. M., S. "A New Paradigm for Decision Making under Uncertainty in Signature Forensics Applications based on Neutrosophic Rule Engine," *Journal of International Journal of Neutrosophic Science*, vol. 24, no. 2, pp. 268-282, 2024.
- [8] Hasan, Z. "Deep Learning for Super Resolution and Applications," *Journal of Galoitica: Journal of Mathematical Structures and Applications*, vol. 8, no. 2, pp. 34-42, 2023.
- [9] F. Smarandache, *A unifying field in logics Neutrosophy Neutrosophic Probability, Set and Logic*, Rehoboth American Research Press (1999).
- [10] Daniel Ramot, Ron Milo, Menahem Friedman, and Abraham Kandel, Complex fuzzy set, *IEEE Transactions on Fuzzy System*, 10(2), 2002.
- [11] S.J Golan, *Semirings and their Applications*, Kluwer Academic Publishers, London, 1999.
- [12] Faward Hussian, Raja Muhammad Hashism, Ajab Khan, Muhammad Naeem, Generalization of bisemirings, *International Journal of Computer Science and Information Security*, 14(9), (2016), 275-289.
- [13] K. M. Lee, Bipolar-valued fuzzy sets and their operations, *Proc. Int. Conf. Intelligent Technologies Bangkok, Thailand*, (2000) 307-312.
- [14] J. Ahsan, K. Saifullah, and F. Khan, Fuzzy semirings, *Fuzzy Sets and systems*, 60, (1993), 309-320.
- [15] Javed Ahsan, John N. Mordeson, and Muhammad Shabir, *Fuzzy Semirings with Applications to Automata Theory*, Springer Heidelberg New York Dordrecht, London, 2012.
- [16] M.K Sen, S. Ghosh An introduction to bisemirings, *Southeast Asian Bulletin of Mathematics*, 28(3), (2001), 547-559.
- [17] Ozcek, M. "A Review on the Structure of Fuzzy Regular Proper Mappings in Fuzzy Topological Spaces and Their Properties," *Journal of Pure Mathematics for Theoretical Computer Science*, vol. 3, no. 2, pp.60-71, 2023.
- [18] Palanikumar M, Selvi G, Ganeshsree Selvachandran and Tan S.L, New approach to bisemiring theory via the bipolar-valued neutrosophic normal sets, *Neutrosophic Sets and Systems*, 55, 427-450, 2023.