



## On The Weak Fuzzy Complex Differential Equations and Some Types of the 1st Order 1st degree WFC-ODEs

Hussein Edduweh<sup>1</sup>, Ahmed Salem Heilat<sup>2,\*</sup>, Lama Razouk<sup>3</sup>, Sara A. Khalil<sup>4</sup>, Ahmed Atallah Alsarairah<sup>5</sup>, Abdallah Al-Husban<sup>6,7</sup>

<sup>1</sup>Department of Mathematics, The University of Texas at Arlington, Arlington, TX 76019-0407, USA

<sup>2</sup>Department of Mathematics, Faculty of Science and Information Technology, Jadara University, P.O. Box 733, Irbid 21110, Jordan

<sup>3</sup>Department of Mathematics, Faculty of Sciences, Tishreen University, Latakia, Syria

<sup>4</sup>Mathematics Department, Faculty of Science, Applied Science Private University (ASU) Amman, Jordan

<sup>5</sup>The university of Jordan–Aqaba Department of computer information systems, Jordan

<sup>6</sup>Department of Mathematics, Faculty of Science and Technology, Irbid National University, P.O. Box: 2600 Irbid, Jordan

<sup>7</sup>Jadara Research Center, Jadara University, Irbid 21110, Jordan

Emails: [Husseinsaid.edduweh@mavs.uta.edu](mailto:Husseinsaid.edduweh@mavs.uta.edu); [ahmed\\_heilat@yahoo.com](mailto:ahmed_heilat@yahoo.com); [Lamarazouk94@gmail.com](mailto:Lamarazouk94@gmail.com); [s\\_khalil@asu.edu.jo](mailto:s_khalil@asu.edu.jo); [a.alsarairah@ju.edu.jo](mailto:a.alsarairah@ju.edu.jo); [dralhoban@inu.edu.jo](mailto:dralhoban@inu.edu.jo)

### Abstract

The objective of this paper is to introduce the concept of Weak Fuzzy Complex differential equations. We have defined the general solution of the  $n$ -th order Weak Fuzzy Complex ordinary differential equation. That we have used a special isomorphism transformation function to write the WFC-ODE as two Real ODEs and solved them with respect to their own variables. Then, by the inverse of the transformation function, we have got the general solution in  $F(J)$  as a structure of two general solutions in  $R$ . Therefore, we have shown some types of first-order first-degree separable, exact, and linear WFC-ODEs. Also, we have found their general solutions with examples to demonstrate them.

**Keywords:** Weak Fuzzy Complex (WFC) Numbers; Weak Fuzzy Complex Functions; Differential Equations (DE)

### 1. Introduction

Real number extensions have always been the focus of attention of researchers from all over the world. In the year 2023, Weak Fuzzy Complex numbers appeared for the first time in [9], where they defined them as a new generalization of classical real numbers by applying fuzzy operators [3,5] studied WFC vector spaces and matrices. The necessary and sufficient conditions for Weak Fuzzy Complex Pythagoras triples and Weak Fuzzy Complex Pythagoras quadruples have been handled in [2,7], as applications of Weak Fuzzy Complex numbers in Diophantine equations. Also, [1] studied WFC linear Diophantine equations in two WFC variables. In addition, a special transformation function that has an important role in working with variables from a Real number set instead of a Weak Fuzzy Complex set was defined in [10], where the foundation of functions in WFC variables was studied. Actually, one of the important results of using this type of numbers, modeling the solutions of some vectorial equations defined by norms in 3-dimensional Euclidean space  $A$ -Curves [4]. However, Python introduced Weak Fuzzy Complex numbers in [11].

On the other side, we know that DEs are important for modeling time dependent processes in many disciplines, e.g. in engineering, physics, chemistry, and economics. For instance, in classical mechanics, if the position, velocity, acceleration, and various forces acting on a body are given, Newton's second law of motion allows us to express these variables as a differential equation and, by solving it, to compute the position of the body as a function of time. According to the important role of differential equations in a lot of fields, we study them in WFC variables.

In section 2 of this paper, we mention the main concepts. Then, section 3 shows the definition of the WFC differential equations and section 4 focuses on some types of first-order WFC-ODEs with some related examples.

## 2. Preliminaries

The foundational concepts that are important to get ODEs in WFC variables are presented in this section.

**Definition 1 ([9])** The set of Weak Fuzzy Complex numbers was defined as follows, where ‘ $J$ ’ is the Weak Fuzzy Complex operator ( $J \notin R$ ):

$$F_J = \{x_0 + x_1 J; x_0, x_1 \in R, J^2 = t \in ]0, 1[ \}$$

Properties of the Weak Fuzzy Complex numbers: ([9])

Let  $X = x_0 + x_1 J, Y = y_0 + y_1 J \in F_J$ , where  $x_0, x_1, y_0, y_1 \in R$

- ◆ Addition:  $X + Y = (x_0 + y_0) + (x_1 + y_1)J$ .
- ◆ Multiplication  $X \times Y = (x_0 y_0 + x_1 y_1 t) + (x_0 y_1 + x_1 y_0)J$ .
- ◆ The conjugate of  $X$  is:  $\bar{X} = x_0 + x_1 J$

**Definition 2 ([9])** Let  $\varphi$  be the transformation function from  $F_J$  to  $R \times R$ , which we define as follows:

$$\varphi: F_J \mapsto R \times R.$$

$$\varphi(x_0 + x_1 J) = (x_0 + x_1(-\sqrt{t}), x_0 + x_1(+\sqrt{t})) = (x_0 - x_1 \sqrt{t}, x_0 + x_1 \sqrt{t});$$

where  $J^2 = t \in ]0, 1[ \Rightarrow J = \pm\sqrt{t}$  and  $x_0, x_1, y_0, y_1 \in R$  (This map is an isomorphism).

**Definition 3** Let  $\varphi: F_J \mapsto R \times R$  such that:  $\varphi(X) = (a, b)$ , the inverse function of  $\varphi$  is defined as follows:

$$\varphi^{-1}: R \times R \mapsto F_J$$

$$\varphi^{-1}(a, b) = \frac{1}{2}[a + b] + \frac{1}{2\sqrt{t}}J[b - a].$$

**Definition 4** On the ring of Weak Fuzzy Complex numbers  $F_J$ , we define the Weak Fuzzy Complex function in one variable as follows:

$$f: F_J \mapsto F_J,$$

$$f = f(X),$$

$$X = x_0 + x_1 J; x_0, x_1 \in R$$

**Remark 1** Let  $f: F_J \mapsto F_J$  be a Weak Fuzzy Complex function in one variable  $X = x_0 + x_1 J; J^2 = t$ . Hence,  $f$  can be written by using two classical functions in two variables as follows:

$$g: R^2 \mapsto R, \text{ s.t. } g = g(x_0, x_1)$$

$$h: R^2 \mapsto R, \text{ s.t. } h = h(x_0, x_1)$$

where  $f(X) = g(x_0, x_1) + h(x_0, x_1)J$ .

**Definition 5** Let  $\varphi: F_J \mapsto R \times R, f: F_J \mapsto F_J$  such that:

$$\varphi(x_0 + x_1 J) = (x_0 - x_1 \sqrt{t}, x_0 + x_1 \sqrt{t}); J^2 = t \in ]0, 1[,$$

$$f = f(X), X = x_0 + x_1 J; x_0, x_1 \in R.$$

the canonical formula as follows:

$$f(X) = \varphi^{-1} \circ \varphi(f(X)) = g(x_0, x_1) + Jh(x_0, x_1).$$

**Remark 2** For  $f: F_J \mapsto F_J$  s.t.  $f = f(X)$ , then

$$\varphi(f(X)) = (f_1(x_0 - x_1 \sqrt{t}), f_2(x_0 + x_1 \sqrt{t})); f_1, f_2: R \mapsto R.$$

On the other hand, we have:

$$g(x_0, x_1) = \frac{1}{2}[f_1(x_0 - x_1 \sqrt{t}) + f_2(x_0 + x_1 \sqrt{t})],$$

$$h(x_0, x_1) = \frac{1}{2\sqrt{t}} J[f_2(x_0 + x_1 \sqrt{t}) - f_1(x_0 - x_1 \sqrt{t})].$$

**Definition 6** Let  $X = x_0 + x_1 J, Y = y_0 + y_1 J \in F_J$ , we say that  $X \leq Y$ , if and only if:

$$\begin{cases} x_0 - x_1 \sqrt{t} \leq y_0 - y_1 \sqrt{t} \\ x_0 + x_1 \sqrt{t} \leq y_0 + y_1 \sqrt{t} \end{cases}$$

**Definition 7** Let  $A = a_0 + a_1 J, B = b_0 + b_1 J \in F_J$ , we define the interval  $[A, B]$  if and only if  $A \leq B$ , according to the definition of the partial order relation ( $\leq$ ).

• If  $A \not\leq B$ , then  $[A, B] = \phi$ .

• We can understand  $[A, B]$  as follows:

$$[A, B] = \{C = c_0 + c_1 J \in F_J; A \leq C \leq B\}.$$

**Definition 8** Let  $X \in F_J$ , we say that  $X = x_0 + x_1 J$  is not an invertible WFC element if one of its  $\varphi(X)$  components at least is zero s.t.

$$x_0 - x_1 \sqrt{t} = 0$$

$$\text{or } x_0 + x_1 \sqrt{t} = 0$$

and that depends on the value of  $\sqrt{t}$ .

**Definition 9** Let  $f$  be a Weak Fuzzy Complex function in one variable defined on  $F_J$ , where

$$\varphi(f(X)) = (f_1(x_0 - x_1 \sqrt{t}), f_2(x_0 + x_1 \sqrt{t})) \in \mathbf{R} \times \mathbf{R};$$

then we say that

1)  $f$  is not invertible if at least one of its  $\varphi(f(X))$  components is zero s.t.

$$f_1(x_0 - x_1 \sqrt{t}) = 0$$

$$\text{or } f_2(x_0 + x_1 \sqrt{t}) = 0$$

s.t.  $f$  is invertible on  $F_J \setminus \{X; \varphi(f(X)) = (f_1, f_2) \in \mathbf{R} \times \mathbf{R} \setminus \{(0,0), (0,k), (k,0); k \in \mathbf{R}\}\}$

$$= F_J \setminus \{X; f(X) \in F_J \setminus \{(\frac{1}{2} \mp \frac{1}{2\sqrt{t}} J)k, 0; k \in \mathbf{R}\}\}$$

$$= F_J \setminus \{X; f(X) \notin \{(\frac{1}{2} \mp \frac{1}{2\sqrt{t}} J)k, 0; k \in \mathbf{R}\}\}$$

where  $\varphi^{-1}(0, k) = \varphi^{-1}(0,1)k = (\frac{1}{2} + \frac{1}{2\sqrt{t}} J)k$  and  $\varphi^{-1}(k, 0) = \varphi^{-1}(1,0)k = (\frac{1}{2} - \frac{1}{2\sqrt{t}} J)k$ .

2) a singular point of the WFC function is a point where the function is not defined or not invertible.

**Definition 10** Let  $f: F_J \mapsto F_J$  be a Weak Fuzzy Complex function in one variable, where

$$\varphi(f(X)) = (f_1(x_0 - x_1 \sqrt{t}), f_2(x_0 + x_1 \sqrt{t})); f_1, f_2: \mathbf{R} \mapsto \mathbf{R}, \text{ then we say:}$$

1)  $f$  is continuous on  $F_J$  if and only if  $f_1, f_2$  are continuous on  $\mathbf{R}$ .

2)  $f$  is differentiable on  $F_J$  if and only if  $f_1, f_2$  are differentiable on  $\mathbf{R}$ , with respect to their own variables.

3)  $f$  is integrable on  $F_J$  if and only if  $f_1, f_2$  are integrable on  $\mathbf{R}$ .

**Definition 11** Let  $f: F_J \mapsto F_J$  be a differentiable/integrable function on  $F_J$ . We define

$$1) f'(X) = \varphi^{-1}(f'_1(x_0 - x_1 \sqrt{t}), f'_2(x_0 + x_1 \sqrt{t}))$$

$$2) \int f(X).dX = \varphi^{-1}(\int f_1.d(x_0 - x_1 \sqrt{t}), \int f_2.d(x_0 + x_1 \sqrt{t})).$$

**Definition 12** Let  $f: F_J \mapsto F_J$  be a differentiable Weak Fuzzy Complex function on  $F_J$ . We can derive  $f$   $n$ -time as follows:

$$f^{(n)}(X) = \varphi^{-1}(f_1^{(n)}(x_0 - x_1\sqrt{t}), f_2^{(n)}(x_0 + x_1\sqrt{t})); n \in \mathbf{N}.$$

### 3. The Differential Equations in $F_J$

The derivative represents a rate of change quantity concerning the change in another quantity. We know that the differential equation is an equation that contains at least one derivative of the dependent variable with respect to the other variables (i.e., the independent variable). If the independent variable is a single, then we call the related equation an ‘‘Ordinary Differential Equation’’ (ODE) which arises most commonly in the study of dynamical systems and electrical networks. ODE is much easier to be treated than ‘‘Partial Differential Equation’’ (PDE), whose unknown function depends on two or more independent variables. In this article, we will be interested in studying ODEs on (WFC-ODEs) [6, 8, 12, 13,14,15,16,17,18,19,20,21,22,23-36].

**Definition 13** Let  $Y$  be a dependent variable, and  $Y = f(X)$  represents an unknown weak fuzzy complex function where the independent variable  $X = x_0 + x_1J$  is a weak fuzzy complex variable. We define *the weak fuzzy complex differential equation of  $n$ -th order* (the highest derivative appearing in the differential equation) as the following:

$$\mathcal{F}(X, Y, Y', Y'', \dots, Y^{(n)}) = 0 \quad (1)$$

where  $Y^{(n)} = \frac{d^n Y}{dX^n}$  and  $Y = y_0 + y_1J = \varphi^{-1}(y_0 - y_1\sqrt{t}, y_0 + y_1\sqrt{t}) = \varphi^{-1}(f_1(x_0 - x_1\sqrt{t}), f_2(x_0 + x_1\sqrt{t}))$ .

**Remark 3** We can consider the differential operator is as follows:

$$\frac{d^n}{dX^n} = \varphi^{-1} \left( \frac{d^n}{d(x_0 - x_1\sqrt{t})^n}, \frac{d^n}{d(x_0 + x_1\sqrt{t})^n} \right) \text{ where } X = \varphi^{-1}(x_0 - x_1\sqrt{t}, x_0 + x_1\sqrt{t}).$$

**Remark 4** The WFC-ODE with the WFC independent variable  $X = x_0 + x_1J$  is equivalent to two ODEs, the first one is with the independent real variable  $x_0 - x_1\sqrt{t}$  and the second ODE is with the independent real variable  $x_0 + x_1\sqrt{t}$ .

**Proof.**

$$\mathcal{F}(X, Y, Y', Y'', \dots, Y^{(n)}) = 0$$

$$\mathcal{F}(x_0 + x_1J, y_0 + y_1J, y'_0 + y'_1J, y''_0 + y''_1J, \dots, y_0^{(n)} + y_1^{(n)}J) = 0$$

By using the function  $\varphi$ , we get

$$\varphi^{-1}(\mathcal{F}_1(x_0 - x_1\sqrt{t}, y_0 - y_1\sqrt{t}, y'_0 - y'_1\sqrt{t}, y''_0 - y''_1\sqrt{t}, \dots, y_0^{(n)} - y_1^{(n)}\sqrt{t}), \mathcal{F}_2(x_0 + x_1\sqrt{t}, y_0 + y_1\sqrt{t}, y'_0 + y'_1\sqrt{t}, y''_0 + y''_1\sqrt{t}, \dots, y_0^{(n)} + y_1^{(n)}\sqrt{t})) = 0$$

$$\Rightarrow \begin{cases} \mathcal{F}_1(x_0 - x_1\sqrt{t}, y_0 - y_1\sqrt{t}, y'_0 - y'_1\sqrt{t}, y''_0 - y''_1\sqrt{t}, \dots, y_1^{(n)} - y_2^{(n)}\sqrt{t}) = 0 & (2-1) \\ \mathcal{F}_2(x_0 + x_1\sqrt{t}, y_0 + y_1\sqrt{t}, y'_0 + y'_1\sqrt{t}, y''_0 + y''_1\sqrt{t}, \dots, y_1^{(n)} + y_2^{(n)}\sqrt{t}) = 0 & (2-2) \end{cases} \quad (2)$$

$$\text{where } \mathcal{F} = \varphi^{-1}(\mathcal{F}_1, \mathcal{F}_2) = \frac{1}{2}(\mathcal{F}_1 + \mathcal{F}_2) + \frac{1}{2\sqrt{t}}J(\mathcal{F}_2 - \mathcal{F}_1).$$

**Definition 14** The functions  $y_0 - y_1\sqrt{t} = f_1(x_0 - x_1\sqrt{t})$  and  $y_0 + y_1\sqrt{t} = f_2(x_0 + x_1\sqrt{t})$  are called *general solutions* to (2-1) and (2-2), respectively, on intervals where  $f_1$  and  $f_2$  are defined (domain of solutions) on  $I_1 \subseteq \mathbf{R}$  and  $I_2 \subseteq \mathbf{R}$  if each of them is an  $n$ -times differentiable in the related interval with respect to their own variables  $x_0 - x_1\sqrt{t}, x_0 + x_1\sqrt{t}$ , respectively.

**Remark 5.** Each general solution of (2-1) and (2-2) contains in general  $n$  real arbitrary constants of integrations respectively. Definition 15. A function

$$Y = f(X) = \varphi^{-1}(f_1(x_0 - x_1\sqrt{t}), f_2(x_0 + x_1\sqrt{t})) = \frac{1}{2}(f_1 + f_2) + \frac{1}{2\sqrt{t}}J(f_2 - f_1)$$

is called *the general solution* of (1) on  $I = \varphi^{-1}(I_1 \times I_2) \subseteq F_J$  if  $f_1$  and  $f_2$  are the general solutions for (2-1) and (2-2), respectively. Such that (1) is satisfied identically when  $f(X)$  and its derivatives are substituted into  $\mathcal{F}$ :

$$\mathcal{F}(X, f(X), f'(X), f''(X), \dots, f^{(n)}(X)) = 0 \quad (3)$$

where

$$\mathcal{F}(X, Y, Y', Y'', \dots, Y^{(n)}) = \mathcal{F}(X, f(X), f'(X), f''(X), \dots, f^{(n)}(X))$$

$$= \varphi^{-1}\varphi(\mathcal{F}(X, f(X), f'(X), f''(X), \dots, f^{(n)}(X)))$$

$$\begin{aligned}
 &= \varphi^{-1}\mathcal{F}(\varphi(X, f(X), f'(X), f''(X), \dots, f^{(n)}(X))) \\
 &= \varphi^{-1}\mathcal{F}((x_0 - x_1\sqrt{t}, x_0 + x_1\sqrt{t}), \\
 &(f_1(x_0 - x_1\sqrt{t}), f_2(x_0 + x_1\sqrt{t})), \\
 &(f'_1(x_0 - x_1\sqrt{t}), f'_2(x_0 + x_1\sqrt{t})), \\
 &(f''_1(x_0 - x_1\sqrt{t}), f''_2(x_0 + x_1\sqrt{t})), \dots, \\
 &(f_1^{(n)}(x_0 - x_1\sqrt{t}), f_2^{(n)}(x_0 + x_1\sqrt{t}))) \\
 &= \varphi^{-1}(\mathcal{F}_1[(x_0 - x_1\sqrt{t}), f_1(x_0 - x_1\sqrt{t}), f'_1(x_0 - x_1\sqrt{t}), f''_1(x_0 - x_1\sqrt{t}), \dots, f_1^{(n)}(x_0 - x_1\sqrt{t})], \\
 &\mathcal{F}_2[(x_0 + x_1\sqrt{t}), f_2(x_0 + x_1\sqrt{t}), f'_2(x_0 + x_1\sqrt{t}), f''_2(x_0 + x_1\sqrt{t}), \dots, f_2^{(n)}(x_0 + x_1\sqrt{t})]).
 \end{aligned}$$

So that, (3) is equivalent to:

$$\begin{cases}
 \mathcal{F}_1[(x_0 - x_1\sqrt{t}), f_1(x_0 - x_1\sqrt{t}), f'_1(x_0 - x_1\sqrt{t}), f''_1(x_0 - x_1\sqrt{t}), \dots, f_1^{(n)}(x_0 - x_1\sqrt{t})] = 0 & (4-1) \\
 \mathcal{F}_2[(x_0 + x_1\sqrt{t}), f_2(x_0 + x_1\sqrt{t}), f'_2(x_0 + x_1\sqrt{t}), f''_2(x_0 + x_1\sqrt{t}), \dots, f_2^{(n)}(x_0 + x_1\sqrt{t})] = 0 & (4-2)
 \end{cases} \quad (4)$$

where:  $\mathcal{F} = \varphi^{-1}(\mathcal{F}_1, \mathcal{F}_2)$ ,  $x_0 - x_1\sqrt{t} \in I_1$ ,  $x_0 + x_1\sqrt{t} \in I_2$

❖ In other words,  $Y = f(X) = \varphi^{-1}(f_1(x_0 - x_1\sqrt{t}), f_2(x_0 + x_1\sqrt{t}))$  is the general solution of (1) on  $I \subseteq F_j$  if  $Y$  is n-times differentiable WFC function on  $I$  and satisfies the equation on  $I$ .

#### 4. First-order and First-degree ODEs in $F_j$

**Definition 16.** The weak fuzzy complex differential equation of the first order is written as follows:

$$\mathcal{F}(X, Y, Y') = 0 \tag{5}$$

$$\Leftrightarrow \begin{cases} \mathcal{F}_1(x_0 - x_1\sqrt{t}, y_0 - y_1\sqrt{t}, y'_0 - y'_1\sqrt{t}) = 0 & (6-1) \\ \mathcal{F}_2(x_0 + x_1\sqrt{t}, y_0 + y_1\sqrt{t}, y'_0 + y'_1\sqrt{t}) = 0 & (6-2) \end{cases} \tag{6}$$

where  $y_0 - y_1\sqrt{t} = f_1(x_0 - x_1\sqrt{t})$ ,  $y_0 + y_1\sqrt{t} = f_2(x_0 + x_1\sqrt{t})$

$$Y' = \frac{dY}{dX} = f'(X) = \varphi^{-1}(f'_1(x_0 - x_1\sqrt{t}), f'_2(x_0 + x_1\sqrt{t}))$$

$$\frac{d}{dX} = \varphi^{-1}\left(\frac{d}{d(x_0 - x_1\sqrt{t})}, \frac{d}{d(x_0 + x_1\sqrt{t})}\right)$$

and  $\mathcal{F} = \varphi^{-1}(\mathcal{F}_1, \mathcal{F}_2)$

**Definition 17.** Let  $y_0 - y_1\sqrt{t} = f_1(x_0 - x_1\sqrt{t})$  and  $y_0 + y_1\sqrt{t} = f_2(x_0 + x_1\sqrt{t})$  are the general solutions to (6-1) on  $I_1 \subseteq \mathbf{R}$  and (6-2) on  $I_2 \subseteq \mathbf{R}$  with real arbitrary constants  $C_0, C_1$ , respectively, then

$$Y = f(X) = \varphi^{-1}(f_1(x_0 - x_1\sqrt{t}), f_2(x_0 + x_1\sqrt{t})),$$

is the general solution of (5) on  $I = \varphi^{-1}(I_1 \times I_2) \subseteq F_j$  with a WFC constant  $C = \varphi^{-1}(C_0, C_1)$  if

$\mathcal{F}(X, f(X), f'(X)) \equiv 0$  holds for all  $X$ ,

where  $= \varphi^{-1}(x_0 - x_1\sqrt{t}, x_0 + x_1\sqrt{t})$ ,  $x_0 - x_1\sqrt{t} \in I_1$ ,  $x_0 + x_1\sqrt{t} \in I_2$  and

$$\begin{cases} \mathcal{F}_1[(x_0 - x_1\sqrt{t}), f_1(x_0 - x_1\sqrt{t}), f'_1(x_0 - x_1\sqrt{t})] = 0 \\ \mathcal{F}_2[(x_0 + x_1\sqrt{t}), f_2(x_0 + x_1\sqrt{t}), f'_2(x_0 + x_1\sqrt{t})] = 0. \end{cases}$$

Therefore, the integrals of (6-1) and (6-2) with respect to  $x_0 - x_1\sqrt{t}$ ,  $x_0 + x_1\sqrt{t}$  respectively, are defined by the following forms:

$$\begin{cases} \int \mathcal{F}_1(x_0 - x_1\sqrt{t}, y_0 - y_1\sqrt{t}, y'_0 - y'_1\sqrt{t})d(x_0 - x_1\sqrt{t}) = C_0 \\ \int \mathcal{F}_2(x_0 + x_1\sqrt{t}, y_0 + y_1\sqrt{t}, y'_0 + y'_1\sqrt{t})d(x_0 + x_1\sqrt{t}) = C_1. \end{cases}$$

Then, the integral can be written as follows:

$$\int \mathcal{F}(X, Y, Y') dX = \varphi^{-1} \left( \int \mathcal{F}_1 d(x_0 - x_1 \sqrt{t}), \int \mathcal{F}_2 d(x_0 + x_1 \sqrt{t}) \right) \\ = \frac{1}{2} \left( \int \mathcal{F}_1 d(x_0 - x_1 \sqrt{t}) + \int \mathcal{F}_2 d(x_0 + x_1 \sqrt{t}) \right) + \frac{1}{2\sqrt{t}} J \left( \int \mathcal{F}_2 d(x_0 + x_1 \sqrt{t}) - \int \mathcal{F}_1 d(x_0 - x_1 \sqrt{t}) \right)$$

Which means that the solution of (5) equivalents to

$$\varphi^{-1} \left( \int \mathcal{F}_1 d(x_0 - x_1 \sqrt{t}), \int \mathcal{F}_2 d(x_0 + x_1 \sqrt{t}) \right) = \varphi^{-1}(C_0, C_1).$$

❖ In other words,  $Y = f(X) = \varphi^{-1}(f_1(x_0 - x_1 \sqrt{t}), f_2(x_0 + x_1 \sqrt{t}))$  is the general solution of (4) on  $I \subseteq F_J$  if  $Y$  is a differentiable WFC function on  $I$ .

**Definition 18.** We consider the explicit WFC first order differential equation

$$Y' = \mathcal{F}(X, Y) \quad (7)$$

where  $Y = y_0 + y_1 J = \varphi^{-1}(y_0 - y_1 \sqrt{t}, y_0 + y_1 \sqrt{t}) = \varphi^{-1}(f_1(x_0 - x_1 \sqrt{t}), f_2(x_0 + x_1 \sqrt{t}))$ ,

$I = \varphi^{-1}(I_1 \times I_2) \subseteq F_J, = \varphi^{-1}(x_0 - x_1 \sqrt{t}, x_0 + x_1 \sqrt{t}), x_0 - x_1 \sqrt{t} \in I_1 \subseteq \mathbf{R}, x_0 + x_1 \sqrt{t} \in I_2 \subseteq \mathbf{R}$

**Definition 19.** A function  $(X): I \rightarrow F_J$ , is called the general solution to the differential equation (7) on  $I \subseteq F_J$  if  $f_1$  and  $f_2$  are differentiable in  $I_1$  and  $I_2$  with respect to their own variables  $x_0 - x_1 \sqrt{t}$ ,  $x_0 + x_1 \sqrt{t}$  respectively, i.e., if

$$\begin{cases} (y_0 - y_1 \sqrt{t})' = \mathcal{F}_1(x_0 - x_1 \sqrt{t}, y_0 - y_1 \sqrt{t}) \text{ for all } x_0 - x_1 \sqrt{t} \in I_1 \\ (y_0 + y_1 \sqrt{t})' = \mathcal{F}_2(x_0 + x_1 \sqrt{t}, y_0 + y_1 \sqrt{t}) \text{ for all } x_0 + x_1 \sqrt{t} \in I_2 \end{cases}$$

$$\Rightarrow \begin{cases} y_0 - y_1 \sqrt{t} = \int \mathcal{F}_1 d(x_0 - x_1 \sqrt{t}) \\ y_0 + y_1 \sqrt{t} = \int \mathcal{F}_2 d(x_0 + x_1 \sqrt{t}) \end{cases}$$

$$\Rightarrow Y = \varphi^{-1}(y_0 - y_1 \sqrt{t}, y_0 + y_1 \sqrt{t}) = \varphi^{-1} \left( \int \mathcal{F}_1 d(x_0 - x_1 \sqrt{t}), \int \mathcal{F}_2 d(x_0 + x_1 \sqrt{t}) \right) \in I$$

$$Y = \frac{1}{2} \left( \int \mathcal{F}_1 d(x_0 - x_1 \sqrt{t}) + \int \mathcal{F}_2 d(x_0 + x_1 \sqrt{t}) \right) + \frac{1}{2\sqrt{t}} J \left( \int \mathcal{F}_2 d(x_0 + x_1 \sqrt{t}) - \int \mathcal{F}_1 d(x_0 - x_1 \sqrt{t}) \right)$$

$I = \varphi^{-1}(I_1 \times I_2) \subseteq F_J, X = \varphi^{-1}(x_0 - x_1 \sqrt{t}, x_0 + x_1 \sqrt{t}), x_0 - x_1 \sqrt{t} \in I_1 \subseteq \mathbf{R}, x_0 + x_1 \sqrt{t} \in I_2 \subseteq \mathbf{R}$ .

**General Algorithm** to solve 1<sup>st</sup> order 1<sup>st</sup> degree WFC-ODEs:

**Step1:** Use  $\varphi$  to write the WFC-DE as two Real DEs

**Step2:** Solve each one of Real DEs with respect to their own variables

**Step3:** Write the WFC general solution as a structure of the two real solutions

**Famous Types of First order and First degree WFC-ODEs:**

#### 4.1. Separable type

$$4.1.1. Y' = h(X)g(Y) \quad (I)$$

**Step1:**

This ODE can be written as:

$$\begin{cases} (y_0 - y_1 \sqrt{t})' = h_1(x_0 - x_1 \sqrt{t})g_1(y_0 - y_1 \sqrt{t}) & (8-1) \\ (y_0 + y_1 \sqrt{t})' = h_2(x_0 + x_1 \sqrt{t})g_2(y_0 + y_1 \sqrt{t}) & (8-2) \end{cases} \quad (8)$$

Dividing (8-1) and (8-2) by  $g_1$  and  $g_2$ , respectively, where  $g_1(y_0 - y_1 \sqrt{t}) \neq 0$  and  $g_2(y_0 + y_1 \sqrt{t}) \neq 0$

$$\Rightarrow \begin{cases} \frac{1}{g_1(y_0 - y_1\sqrt{t})} d(y_0 - y_1\sqrt{t}) = h_1(x_0 - x_1\sqrt{t})d(x_0 - x_1\sqrt{t}) & (9-1) \\ \frac{1}{g_2(y_0 + y_1\sqrt{t})} d(y_0 + y_1\sqrt{t}) = h_2(x_0 + x_1\sqrt{t})d(x_0 + x_1\sqrt{t}) & (9-2) \end{cases} \quad (9)$$

It means  $\frac{1}{g(Y)} dY = h(X)dX$  (where  $g$  is invertible).

(9-1), (9-2) are called *separable* ODEs where  $h_1, h_2, g_1$  and  $g_2$  are real functions.

### Step2:

Integrating both sides with respect to  $x_0 - x_1\sqrt{t}$  and  $x_0 + x_1\sqrt{t}$  respectively, we get

$$\Rightarrow \begin{cases} \int \frac{1}{g_1(y_0 - y_1\sqrt{t})} \frac{d(y_0 - y_1\sqrt{t})}{d(x_0 - x_1\sqrt{t})} d(x_0 - x_1\sqrt{t}) = \int h_1(x_0 - x_1\sqrt{t})d(x_0 - x_1\sqrt{t}) \\ \int \frac{1}{g_2(y_0 + y_1\sqrt{t})} \frac{d(y_0 + y_1\sqrt{t})}{d(x_0 + x_1\sqrt{t})} d(x_0 + x_1\sqrt{t}) = \int h_2(x_0 + x_1\sqrt{t})d(x_0 + x_1\sqrt{t}) \end{cases}$$

$$\Leftrightarrow \int \frac{1}{g(Y)} \frac{dY}{dX} dX = \int h(X)dX.$$

Assuming that these integrals maybe evaluated, we get the general solutions to (10-1), (10-2) respectively:

$$\begin{cases} G_1(y_0 - y_1\sqrt{t}) = H_1(x_0 - x_1\sqrt{t}) + C_0 & (11-1) \\ G_2(y_0 + y_1\sqrt{t}) = H_2(x_0 + x_1\sqrt{t}) + C_1 & (11-2) \end{cases} \quad (11)$$

where  $C_0, C_1 \in \mathbf{R}$  are arbitrary constants.

### Step3:

$$\varphi^{-1}(G_1, G_2) = \varphi^{-1}(H_1, H_2) + \varphi^{-1}(C_0, C_1); C = \varphi^{-1}(C_0, C_1)$$

$$\Rightarrow G(Y) = H(X) + C.$$

### Example 1.

$$Y' = e^Y \sin(X) \Leftrightarrow dY = e^Y \sin(X)dX$$

$$\begin{aligned} \varphi^{-1}(d(y_0 - y_1\sqrt{t}), d(y_0 + y_1\sqrt{t})) \\ = \varphi^{-1}(e^{y_0 - y_1\sqrt{t}} \sin(x_0 - x_1\sqrt{t})d(x_0 - x_1\sqrt{t}), e^{y_0 + y_1\sqrt{t}} \sin(x_0 + x_1\sqrt{t})d(x_0 + x_1\sqrt{t})) \\ \Rightarrow \begin{cases} \frac{1}{e^{y_0 - y_1\sqrt{t}}} d(y_0 - y_1\sqrt{t}) = \sin(x_0 - x_1\sqrt{t})d(x_0 - x_1\sqrt{t}) \\ \frac{1}{e^{y_0 + y_1\sqrt{t}}} d(y_0 + y_1\sqrt{t}) = \sin(x_0 + x_1\sqrt{t})d(x_0 + x_1\sqrt{t}) \end{cases} \end{aligned}$$

by integrating both sides of the equation with respect to  $x_0 - x_1\sqrt{t}$  and  $x_0 + x_1\sqrt{t}$ , respectively:

$$\begin{aligned} \begin{cases} \int \frac{1}{e^{y_0 - y_1\sqrt{t}}} d(y_0 - y_1\sqrt{t}) = \int \sin(x_0 - x_1\sqrt{t})d(x_0 - x_1\sqrt{t}) \\ \int \frac{1}{e^{y_0 + y_1\sqrt{t}}} d(y_0 + y_1\sqrt{t}) = \int \sin(x_0 + x_1\sqrt{t})d(x_0 + x_1\sqrt{t}) \end{cases} \\ \Rightarrow \begin{cases} -e^{-(y_0 - y_1\sqrt{t})} = -\cos(x_0 - x_1\sqrt{t}) - C_0 \\ -e^{-(y_0 + y_1\sqrt{t})} = -\cos(x_0 + x_1\sqrt{t}) - C_1 \end{cases} \\ \Rightarrow \begin{cases} y_0 - y_1\sqrt{t} = -\ln[\cos(x_0 - x_1\sqrt{t}) + C_0] \text{ (where } \cos(x_0 - x_1\sqrt{t}) + C_0 > 0) \\ y_0 + y_1\sqrt{t} = -\ln[\cos(x_0 + x_1\sqrt{t}) + C_1] \text{ (where } \cos(x_0 + x_1\sqrt{t}) + C_1 > 0) \end{cases} \\ \xRightarrow{\varphi^{-1}} Y = \frac{1}{2}(y_0 - y_1\sqrt{t} + y_0 + y_1\sqrt{t}) + \frac{1}{2\sqrt{t}}J(y_0 + y_1\sqrt{t} - y_0 + y_1\sqrt{t}) \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{2} (-\ln[\cos(x_0 - x_1\sqrt{t}) + C_0] - \ln[\cos(x_0 + x_1\sqrt{t}) + C_1]) + \frac{1}{2\sqrt{t}} J(-\ln[\cos(x_0 + x_1\sqrt{t}) + C_1] + \ln[\cos(x_0 - x_1\sqrt{t}) + C_0]) \\
&= \varphi^{-1}(-\ln[\cos(x_0 - x_1\sqrt{t}) + C_0], -\ln[\cos(x_0 + x_1\sqrt{t}) + C_1]) \\
&= -\ln(\varphi^{-1}[\cos(x_0 - x_1\sqrt{t}) + C_0, \cos(x_0 + x_1\sqrt{t}) + C_1]) \\
&= -\ln(\varphi^{-1}(\cos(x_0 - x_1\sqrt{t}), \cos(x_0 + x_1\sqrt{t}))) + \varphi^{-1}(C_0, C_1)
\end{aligned}$$

Thus, the general solution to the given WFC-ODE is

$$Y = -\ln[\cos(X) + C] \text{ where } \cos(X) + C > 0.$$

$$4.1.2. Y' = \mathcal{F}(X) \quad (II)$$

$$\Rightarrow dY = \mathcal{F}(X)dX$$

**Step1:**

$$(II) \Leftrightarrow \begin{cases} d(y_0 - y_1\sqrt{t}) = \mathcal{F}_1(x_0 - x_1\sqrt{t})d(x_0 - x_1\sqrt{t}) & (12-1) \\ d(y_0 + y_1\sqrt{t}) = \mathcal{F}_2(x_0 + x_1\sqrt{t})d(x_0 + x_1\sqrt{t}) & (12-2) \end{cases} \quad (12)$$

**Step2:**

Solving the WFC-DE (II) is equivalent to solve two Real separable DEs (12-1), (12-2):

Suppose the function  $\mathcal{F}_1, \mathcal{F}_2$  is continuous functions in intervals  $I_1, I_2$ , respectively. Integrating both sides with respect to  $x_0 - x_1\sqrt{t}$  and  $x_0 + x_1\sqrt{t}$  respectively, we get

$$\begin{aligned}
\int d(y_0 - y_1\sqrt{t}) &= \int \mathcal{F}_1(x_0 - x_1\sqrt{t})d(x_0 - x_1\sqrt{t}) & (13-1) \\
\int d(y_0 + y_1\sqrt{t}) &= \int \mathcal{F}_2(x_0 + x_1\sqrt{t})d(x_0 + x_1\sqrt{t}) & (13-2)
\end{aligned} \quad (13)$$

Assuming that these integrals maybe evaluated, we get the general solutions to (13-1), (13-2) respectively:

$$\Rightarrow \begin{cases} y_0 - y_1\sqrt{t} = r_1(x_0 - x_1\sqrt{t}) + C_0 \\ y_0 + y_1\sqrt{t} = r_2(x_0 + x_1\sqrt{t}) + C_1 \end{cases}$$

where  $C_0, C_1 \in R$  are arbitrary constants.

Also,

$$\int \mathcal{F}_1(x_0 - x_1\sqrt{t})d(x_0 - x_1\sqrt{t}) = r_1(x_0 - x_1\sqrt{t}) + C_0 \text{ and } \int \mathcal{F}_2(x_0 + x_1\sqrt{t})d(x_0 + x_1\sqrt{t}) = r_2(x_0 + x_1\sqrt{t}) + C_1$$

**Step3:**

Thus, the general solution to the DE (II) is:

$$\varphi^{-1}(y_0 - y_1\sqrt{t}, y_0 + y_1\sqrt{t}) = \varphi^{-1}(r_1(x_0 - x_1\sqrt{t}) + C_0, r_2(x_0 + x_1\sqrt{t}) + C_1)$$

$$\Rightarrow Y = r(X) + C$$

where  $r(X) = \varphi^{-1}(r_1(x_0 - x_1\sqrt{t}), r_2(x_0 + x_1\sqrt{t}))$  and  $C = \varphi^{-1}(C_0, C_1)$

**Example 2.**

$$Y' = e^X \Leftrightarrow dY = e^X dX$$

$$\varphi^{-1}(d(y_0 - y_1\sqrt{t}), d(y_0 + y_1\sqrt{t})) = \varphi^{-1}(e^{x_0 - x_1\sqrt{t}}d(x_0 - x_1\sqrt{t}), e^{x_0 + x_1\sqrt{t}}d(x_0 + x_1\sqrt{t}))$$

by integrating both sides of the equation with respect to  $x_0 - x_1\sqrt{t}$  and  $x_0 + x_1\sqrt{t}$ , respectively

$$\varphi^{-1}\left(\int d(y_0 - y_1\sqrt{t}), \int d(y_0 + y_1\sqrt{t})\right) = \varphi^{-1}\left(\int e^{x_0 - x_1\sqrt{t}}d(x_0 - x_1\sqrt{t}), \int e^{x_0 + x_1\sqrt{t}}d(x_0 + x_1\sqrt{t})\right),$$

$$\Rightarrow \varphi^{-1}(y_0 - y_1\sqrt{t}, y_0 + y_1\sqrt{t}) = \varphi^{-1}(e^{x_0 - x_1\sqrt{t}} + C_0, e^{x_0 + x_1\sqrt{t}} + C_1)$$

$$\begin{aligned}
&\Rightarrow \frac{1}{2}(y_0 - y_1\sqrt{t} + y_0 + y_1\sqrt{t}) + \frac{1}{2\sqrt{t}}J(y_0 + y_1\sqrt{t} - y_0 + y_1\sqrt{t}) \\
&= \frac{1}{2}(e^{x_0-x_1\sqrt{t}} + C_0 + e^{x_0+x_1\sqrt{t}} + C_1) + \frac{1}{2\sqrt{t}}J(e^{x_0+x_1\sqrt{t}} + C_1 - e^{x_0-x_1\sqrt{t}} - C_0) \\
&= \frac{1}{2}(e^{x_0-x_1\sqrt{t}} + e^{x_0+x_1\sqrt{t}}) + \frac{1}{2\sqrt{t}}J(e^{x_0+x_1\sqrt{t}} - e^{x_0-x_1\sqrt{t}}) + \frac{1}{2}(C_0 + C_1) + \frac{1}{2\sqrt{t}}J(C_1 - C_0) \\
&\Rightarrow y_0 + y_1J = e^{x_0+x_1J} + C_0 + C_1J.
\end{aligned}$$

The general solution is  $Y = e^X + C$ .

### Example 3.

$$(\sin(X) + \cos(X))dY + (\cos(X) - \sin(X))dX = 0$$

$$\begin{aligned}
\varphi^{-1}(\sin(x_0 - x_1\sqrt{t}) + \cos(x_0 - x_1\sqrt{t}), \sin(x_0 + x_1\sqrt{t}) + \cos(x_0 + x_1\sqrt{t}))(d(y_0 - y_1\sqrt{t}), d(y_0 + y_1\sqrt{t})) \\
+ (\sin(x_0 - x_1\sqrt{t}) - \cos(x_0 - x_1\sqrt{t}), \sin(x_0 + x_1\sqrt{t}) \\
- \cos(x_0 + x_1\sqrt{t}))(d(x_0 - x_1\sqrt{t}), d(x_0 + x_1\sqrt{t})) = 0
\end{aligned}$$

$\Rightarrow$

$$\begin{cases}
[\sin(x_0 - x_1\sqrt{t}) + \cos(x_0 - x_1\sqrt{t})]d(y_0 - y_1\sqrt{t}) = [\sin(x_0 - x_1\sqrt{t}) - \cos(x_0 - x_1\sqrt{t})]d(x_0 - x_1\sqrt{t}) & (14 - 1) \\
[\sin(x_0 + x_1\sqrt{t}) + \cos(x_0 + x_1\sqrt{t})]d(y_0 + y_1\sqrt{t}) = [\sin(x_0 + x_1\sqrt{t}) - \cos(x_0 + x_1\sqrt{t})]d(x_0 + x_1\sqrt{t}) & (14 - 2)
\end{cases} \quad (14)$$

Dividing (14 - 1) by  $\sin(x_0 - x_1\sqrt{t}) + \cos(x_0 - x_1\sqrt{t}) \neq 0$  and integrating both sides with respect to  $x_0 - x_1\sqrt{t}$ . And dividing (14 - 2) by  $\sin(x_0 + x_1\sqrt{t}) + \cos(x_0 + x_1\sqrt{t}) \neq 0$  and integrating both sides with respect to  $x_0 + x_1\sqrt{t}$ , we get

$$\begin{cases}
\int d(y_0 - y_1\sqrt{t}) = \int \frac{\cos(x_0 - x_1\sqrt{t}) - \sin(x_0 - x_1\sqrt{t})}{\sin(x_0 - x_1\sqrt{t}) + \cos(x_0 - x_1\sqrt{t})} d(x_0 - x_1\sqrt{t}) \\
\int d(y_0 + y_1\sqrt{t}) = \int \frac{\cos(x_0 + x_1\sqrt{t}) - \sin(x_0 + x_1\sqrt{t})}{\sin(x_0 + x_1\sqrt{t}) + \cos(x_0 + x_1\sqrt{t})} d(x_0 + x_1\sqrt{t}),
\end{cases}$$

$$\Leftrightarrow \int dY = \int \frac{\cos(X) - \sin(X)}{\sin(X) + \cos(X)} dX,$$

$$\begin{cases}
y_0 - y_1\sqrt{t} = \ln|\sin(x_0 - x_1\sqrt{t}) + \cos(x_0 - x_1\sqrt{t})| + C_0 \\
y_0 + y_1\sqrt{t} = \ln|\sin(x_0 + x_1\sqrt{t}) + \cos(x_0 + x_1\sqrt{t})| + C_1,
\end{cases}$$

$$\Rightarrow \frac{1}{2}(y_0 - y_1\sqrt{t} + y_0 + y_1\sqrt{t}) + \frac{1}{2\sqrt{t}}J(y_0 + y_1\sqrt{t} - y_0 + y_1\sqrt{t})$$

$$= \frac{1}{2}(\ln|\sin(x_0 - x_1\sqrt{t}) + \cos(x_0 - x_1\sqrt{t})| + C_0 + \ln|\sin(x_0 + x_1\sqrt{t}) + \cos(x_0 + x_1\sqrt{t})| + C_1)$$

$$+ \frac{1}{2\sqrt{t}}J(\ln|\sin(x_0 + x_1\sqrt{t}) + \cos(x_0 + x_1\sqrt{t})| + C_1 - \ln|\sin(x_0 - x_1\sqrt{t}) + \cos(x_0 - x_1\sqrt{t})| - C_0)$$

$$\Rightarrow y_0 + y_1J = \ln|\sin(x_0 + x_1J) + \cos(x_0 + x_1J)| + C_0 + C_1J.$$

The general solution is

$$Y = -\ln|\sin(X) + \cos(X)| + C; \quad C = C_0 + C_1J$$

$$Y = \ln \frac{1}{|\sin(X) + \cos(X)|} + C; \quad \sin(X) + \cos(X) \neq 0.$$

### Example 4.

$$Y' = X^3 + \cos(X)$$

we write it in the following form:

$$dY = (X^3 + \cos(X))dX$$

$$\begin{aligned}
& \varphi^{-1} \left( \int d(y_0 - y_1\sqrt{t}), \int d(y_0 + y_1\sqrt{t}) \right) \\
&= \varphi^{-1} \left( \int ((x_0 - x_1\sqrt{t})^3 + \cos(x_0 - x_1\sqrt{t}))d(x_0 - x_1\sqrt{t}), \int ((x_0 + x_1\sqrt{t})^3 + \cos(x_0 + x_1\sqrt{t}))d(x_0 + x_1\sqrt{t}) \right) \\
& \varphi^{-1}(y_0 - y_1\sqrt{t}, y_0 + y_1\sqrt{t}) \\
&= \varphi^{-1} \left( \frac{(x_0 - x_1\sqrt{t})^4}{4} + \sin(x_0 - x_1\sqrt{t}) + C_0, \frac{(x_0 + x_1\sqrt{t})^4}{4} + \sin(x_0 + x_1\sqrt{t}) + C_1 \right) \\
&\Rightarrow \frac{1}{2}(y_0 - y_1\sqrt{t} + y_0 + y_1\sqrt{t}) + \frac{1}{2\sqrt{t}}J(y_0 + y_1\sqrt{t} - y_0 + y_1\sqrt{t}) \\
&= \frac{1}{2} \left( \frac{(x_0 - x_1\sqrt{t})^4}{4} + \sin(x_0 - x_1\sqrt{t}) + \frac{(x_0 + x_1\sqrt{t})^4}{4} + \sin(x_0 + x_1\sqrt{t}) + C_0 + C_1 \right) \\
&+ \frac{1}{2\sqrt{t}}J \left( \frac{(x_0 - x_1\sqrt{t})^4}{4} + \sin(x_0 + x_1\sqrt{t}) - \frac{(x_0 - x_1\sqrt{t})^4}{4} - \sin(x_0 - x_1\sqrt{t}) + C_1 - C_0 \right) \\
&\Rightarrow y_0 + y_1J \\
&= \frac{1}{2} \left( \frac{(x_0 - x_1\sqrt{t})^4}{4} + \frac{(x_0 + x_1\sqrt{t})^4}{4} \right) + \frac{1}{2\sqrt{t}}J \left( \frac{(x_0 + x_1\sqrt{t})^4}{4} - \frac{(x_0 - x_1\sqrt{t})^4}{4} \right) \\
&+ \frac{1}{2} \left( \sin(x_0 - x_1\sqrt{t}) + \sin(x_0 + x_1\sqrt{t}) \right) + \frac{1}{2\sqrt{t}}J \left( \sin(x_0 + x_1\sqrt{t}) - \sin(x_0 - x_1\sqrt{t}) \right) \\
&+ \frac{1}{2} (C_0 + C_1) + \frac{1}{2\sqrt{t}}J(C_1 - C_0) \\
&\Rightarrow y_0 + y_1J = \frac{(x_0 + x_1J)^4}{4} + \sin(x_0 + x_1J) + C_0 + C_1J
\end{aligned}$$

The general solution is  $Y = \frac{X^4}{4} + \sin(X) + C$ .

#### 4.1.3. $Y' = \mathcal{F}(Y)$

(III)

$$\Rightarrow dY = \mathcal{F}(Y)dX$$

Here is similar to the previous ODE(II) but with exchange roles of X and Y.

**Step1:**

$$\text{(III)} \Leftrightarrow \begin{cases} d(y_0 - y_1\sqrt{t}) = \mathcal{F}_1(y_0 - y_1\sqrt{t})d(x_0 - x_1\sqrt{t}) & (15-1) \\ d(y_0 + y_1\sqrt{t}) = \mathcal{F}_2(y_0 + y_1\sqrt{t})d(x_0 + x_1\sqrt{t}) & (15-2) \end{cases} \quad (15)$$

**Step2:**

Solving the WFC-DE (II) is equivalent to solve two Real separable DEs (15 – 1), (15 – 2).

Suppose the function  $\mathcal{F}_1, \mathcal{F}_2$  is continuous functions in intervals  $I_1, I_2$ , respectively. Integrating both sides with respect to  $x_0 - x_1\sqrt{t}$  and  $x_0 + x_1\sqrt{t}$  respectively, we get

$$\begin{cases} \int \frac{1}{\mathcal{F}_1(y_0 - y_1\sqrt{t})} d(y_0 - y_1\sqrt{t}) = \int d(x_0 - x_1\sqrt{t}) & (16-1) \\ \int \frac{1}{\mathcal{F}_2(y_0 + y_1\sqrt{t})} d(y_0 + y_1\sqrt{t}) = \int d(x_0 + x_1\sqrt{t}) & (16-2), \end{cases} \quad (16)$$

where  $\mathcal{F}_1(y_0 - y_1\sqrt{t}) \neq 0$  and  $\mathcal{F}_2(y_0 + y_1\sqrt{t}) \neq 0$  (where  $\mathcal{F}$  is invertible).

We get the general solutions to (16 – 1), (16 – 2), respectively:

$$\begin{cases} S_1(y_0 - y_1\sqrt{t}) = x_0 - x_1\sqrt{t} + C_0 & \Rightarrow x_0 - x_1\sqrt{t} = S_1(y_0 - y_1\sqrt{t}) - C_0 \quad (17 - 1) \\ S_2(y_0 + y_1\sqrt{t}) = x_0 + x_1\sqrt{t} + C_1 & \Rightarrow x_0 + x_1\sqrt{t} = S_2(y_0 + y_1\sqrt{t}) - C_1 \quad (17 - 2) \end{cases} \quad (17)$$

where  $C_0, C_1 \in \mathbf{R}$  are arbitrary constants.

Note that in (17 - 1) we exchange roles of  $x_0 - x_1\sqrt{t}$  and  $y_0 - y_1\sqrt{t}$  and in (17 - 2) we exchange roles of  $x_0 + x_1\sqrt{t}$  and  $y_0 + y_1\sqrt{t}$ .

**Step3:**

$$\begin{aligned} \varphi^{-1}(x_0 - x_1\sqrt{t}, x_0 + x_1\sqrt{t}) &= \varphi^{-1}(S_1(y_0 - y_1\sqrt{t}) + C_0, S_2(y_0 + y_1\sqrt{t}) + C_1), \\ \Rightarrow X &= S(Y) + C \end{aligned}$$

where  $S(Y) = \varphi^{-1}(S_1(y_0 - y_1\sqrt{t}), S_2(y_0 + y_1\sqrt{t}))$  and  $C = \varphi^{-1}(C_0, C_1)$ .

Hence, the general solution  $Y(X)$  to the WFC-ODE (III) is the inverse of the function  $X = X(Y)$ .

**Example 5.**

$$Y' = -2Y \quad (18)$$

$$\begin{aligned} \Leftrightarrow \begin{cases} d(y_0 - y_1\sqrt{t}) = -2(y_0 - y_1\sqrt{t})d(x_0 - x_1\sqrt{t}) \\ d(y_0 + y_1\sqrt{t}) = -2(y_0 + y_1\sqrt{t})d(x_0 + x_1\sqrt{t}) \end{cases} \\ \left\{ \begin{aligned} \int \frac{1}{(y_0 - y_1\sqrt{t})} d(y_0 - y_1\sqrt{t}) &= -2 \int d(x_0 - x_1\sqrt{t}) \\ \int \frac{1}{(y_0 + y_1\sqrt{t})} d(y_0 + y_1\sqrt{t}) &= -2 \int d(x_0 + x_1\sqrt{t}) \end{aligned} \right. \end{aligned}$$

where  $y_0 - y_1\sqrt{t} \neq 0$  and  $y_0 + y_1\sqrt{t} \neq 0$ .

$$\Leftrightarrow \begin{cases} \ln|y_0 - y_1\sqrt{t}| = -2(x_0 - x_1\sqrt{t}) + C_0 & \Leftrightarrow |y_0 - y_1\sqrt{t}| = e^{C_0 - 2(x_0 - x_1\sqrt{t})} \\ \ln|y_0 + y_1\sqrt{t}| = -2(x_0 + x_1\sqrt{t}) + C_1 & \Leftrightarrow |y_0 + y_1\sqrt{t}| = e^{C_1 - 2(x_0 + x_1\sqrt{t})} \end{cases}$$

where  $C_0, C_1 \in \mathbf{R}$  are arbitrary constants.

The general solutions with  $\pm e^{C_0}, \pm e^{C_1}$  replaced with  $C_0, C_1$ , respectively:

$$\Leftrightarrow \begin{cases} y_0 - y_1\sqrt{t} = C_0 e^{-2(x_0 - x_1\sqrt{t})} \\ y_0 + y_1\sqrt{t} = C_1 e^{-2(x_0 + x_1\sqrt{t})} \end{cases}$$

The general solution to (18) is written as:

$$\varphi^{-1}(y_0 - y_1\sqrt{t}, y_0 + y_1\sqrt{t}) = \varphi^{-1}(C_0 e^{-2(x_0 - x_1\sqrt{t})}, C_1 e^{-2(x_0 + x_1\sqrt{t})}) \Leftrightarrow Y = C \cdot e^{-2X}.$$

**4.1.4.  $Y' = \mathcal{F}(AX + BY + C)$  (IV)**

This structure of the differential equation can be reduced by simple transformation to one of the previous types already discussed. Assuming the function  $\mathcal{F}$  is continuous in an interval  $I \subseteq \mathbf{F}_J$ .

$$\begin{aligned} &\varphi^{-1}((y_0 - y_1\sqrt{t})', (y_0 + y_1\sqrt{t})') \\ &= \varphi^{-1}(\mathcal{F}_1[(A_0 - A_1\sqrt{t})(x_0 - x_1\sqrt{t}) + (B_0 - B_1\sqrt{t})(y_0 - y_1\sqrt{t}) + C_0 - C_1\sqrt{t}], \mathcal{F}_2[(A_0 + A_1\sqrt{t})(x_0 + x_1\sqrt{t}) \\ &\quad + (B_0 + B_1\sqrt{t})(y_0 + y_1\sqrt{t}) + C_0 + C_1\sqrt{t}]) \end{aligned}$$

$$\begin{aligned} \Rightarrow \\ \begin{cases} ((y_0 - y_1\sqrt{t})') = \mathcal{F}_1[(A_0 - A_1\sqrt{t})(x_0 - x_1\sqrt{t}) + (B_0 - B_1\sqrt{t})(y_0 - y_1\sqrt{t}) + C_0 - C_1\sqrt{t}] \quad (19 - 1) \\ ((y_0 + y_1\sqrt{t})') = \mathcal{F}_2[(A_0 + A_1\sqrt{t})(x_0 + x_1\sqrt{t}) + (B_0 + B_1\sqrt{t})(y_0 + y_1\sqrt{t}) + C_0 + C_1\sqrt{t}] \quad (19 - 2) \end{cases} \quad (19) \end{aligned}$$

where  $A = A_0 + A_1 J, B = B_0 + B_1 J \in \mathbf{F}_J, C = C_0 + C_1 J \in \mathbf{F}_J$ .

We can say that solving (IV) is equivalent to solve the two ODEs (19 – 1 & 19 – 2) in real field with respect to variables  $x_0 - x_1\sqrt{t}$  and  $x_0 + x_1\sqrt{t}$ , respectively. For that, we suppose:

$$\begin{cases} U_1(x_0 - x_1\sqrt{t}) = (A_0 - A_1\sqrt{t})(x_0 - x_1\sqrt{t}) + (B_0 - B_1\sqrt{t})(y_0 - y_1\sqrt{t}) + C_0 - C_1\sqrt{t} \\ U_2(x_0 + x_1\sqrt{t}) = (A_0 + A_1\sqrt{t})(x_0 + x_1\sqrt{t}) + (B_0 + B_1\sqrt{t})(y_0 + y_1\sqrt{t}) + C_0 + C_1\sqrt{t} \end{cases}$$

$$\Rightarrow \begin{cases} U_1(x_0 - x_1\sqrt{t}) = (A_0 - A_1\sqrt{t})(x_0 - x_1\sqrt{t}) + (B_0 - B_1\sqrt{t})Y_1(x_0 - x_1\sqrt{t}) + C_0 - C_1\sqrt{t} \quad (20 - 1) \\ U_2(x_0 + x_1\sqrt{t}) = (A_0 + A_1\sqrt{t})(x_0 + x_1\sqrt{t}) + (B_0 + B_1\sqrt{t})Y_2(x_0 + x_1\sqrt{t}) + C_0 + C_1\sqrt{t} \quad (20 - 2) \end{cases} \quad (20)$$

$$(IV) \Rightarrow \begin{cases} (y_0 - y_1\sqrt{t})' = \mathcal{F}_1[U_1(x_0 - x_1\sqrt{t})] \\ (y_0 + y_1\sqrt{t})' = \mathcal{F}_2[U_2(x_0 + x_1\sqrt{t})] \end{cases}$$

(We are interested in the case where  $B_0 + B_1\sqrt{t} \neq 0, B_0 - B_1\sqrt{t} \neq 0$ )

$$Y(X) = \varphi^{-1}(Y_1(x_0 - x_1\sqrt{t}), Y_2(x_0 + x_1\sqrt{t})) \in \mathbf{F}_J(X)$$

$$= \varphi^{-1}(y_0 - y_1\sqrt{t}, y_0 + y_1\sqrt{t})$$

$$U(X) = \varphi^{-1}(U_1(x_0 - x_1\sqrt{t}), U_2(x_0 + x_1\sqrt{t})) \in \mathbf{F}_J(X)$$

So that, we can write

$$U(X) = AX + BY(X) + C \tag{21} \quad (\text{If})$$

$Y_1$  is a solution to (20 – 1) and  $Y_2$  is a solution to (20 – 2), then  $U_1$  and  $U_2$  satisfy

$$\begin{cases} U_1'(x_0 - x_1\sqrt{t}) = (A_0 - A_1\sqrt{t}) + (B_0 - B_1\sqrt{t})(y_0 - y_1\sqrt{t})' \\ U_2'(x_0 + x_1\sqrt{t}) = (A_0 + A_1\sqrt{t}) + (B_0 + B_1\sqrt{t})(y_0 + y_1\sqrt{t})' \end{cases}$$

Thus,

$$U'(X) = A + BY'(X)$$

$$U' = A + BF(U) \tag{22}$$

which is a solvable equation of the type (III).

Conversely, from a solution  $U(X)$  to (22) we can get a solution  $Y(X)$  to (21).

**Example 6.**

$$Y' = (X + Y)^2 \tag{23}$$

$$\Leftrightarrow \varphi^{-1}((y_0 - y_1\sqrt{t})', (y_0 + y_1\sqrt{t})') = \varphi^{-1}((x_0 - x_1\sqrt{t} + y_0 - y_1\sqrt{t})^2, (x_0 + x_1\sqrt{t} + y_0 + y_1\sqrt{t})^2).$$

To solve it, we use the following ansatz:

$$\begin{cases} U_1(x_0 - x_1\sqrt{t}) = x_0 - x_1\sqrt{t} + y_0 - y_1\sqrt{t} \\ U_2(x_0 + x_1\sqrt{t}) = x_0 + x_1\sqrt{t} + y_0 + y_1\sqrt{t} \end{cases}$$

$$\frac{1}{2}(U_1 + U_2) + \frac{1}{2\sqrt{t}}J(U_2 - U_1) = \frac{1}{2}(x_0 + y_0 + x_0 + y_0) + \frac{1}{2\sqrt{t}}J(2x_1\sqrt{t} - 2y_1\sqrt{t}),$$

$$U(X) = \varphi^{-1}(U_1, U_2) = (x_0 + y_0) + (x_1 + y_1)J,$$

$$\Rightarrow U(X) = X + Y(X).$$

Then, (23) becomes like

$$U'(X) = U^2 + 1$$

$$\frac{dU}{dX} = U^2 + 1 \Rightarrow \frac{dU}{U^2+1} - dX = 0 \quad (\text{Separable ODE})$$

By integrating:

$$\arctan(U) - X = C \Rightarrow \arctan(U) = C + X \Rightarrow U = \tan(C + X)$$

Going back to the old variables, we get the general solution:

$$Y = \tan(C + X) - X.$$

**4.1.5. The Homogeneous ODE**

$$Y' = \mathcal{F}\left(\frac{Y}{X}\right); X \in F_j \setminus \left\{ \left(\frac{1}{2} \mp \frac{1}{2\sqrt{t}}\right)k, 0; k \in R \right\} \tag{V}$$

$$\Leftrightarrow \varphi^{-1}((y_0 - y_1\sqrt{t})', (y_0 + y_1\sqrt{t})') = \varphi^{-1}\left(\mathcal{F}_1\left(\frac{y_0 - y_1\sqrt{t}}{x_0 - x_1\sqrt{t}}\right), \mathcal{F}_2\left(\frac{y_0 + y_1\sqrt{t}}{x_0 + x_1\sqrt{t}}\right)\right)$$

Supposing

$$U_1(x_0 - x_1\sqrt{t}) = \frac{y_0 - y_1\sqrt{t}}{x_0 - x_1\sqrt{t}} \text{ and } U_2(x_0 + x_1\sqrt{t}) = \frac{y_0 + y_1\sqrt{t}}{x_0 + x_1\sqrt{t}}.$$

Then, supposing  $U(X) = \frac{Y(X)}{X}$  and calculating the derivative

$$\begin{cases} y_0 - y_1\sqrt{t} = (x_0 - x_1\sqrt{t})U_1(x_0 - x_1\sqrt{t}) \Rightarrow (y_0 - y_1\sqrt{t})' = U_1 + (x_0 - x_1\sqrt{t})U_1' & (24 - 1) \\ y_0 + y_1\sqrt{t} = (x_0 + x_1\sqrt{t})U_2(x_0 + x_1\sqrt{t}) \Rightarrow (y_0 + y_1\sqrt{t})' = U_2 + (x_0 + x_1\sqrt{t})U_2' & (24 - 2) \end{cases} \tag{24}$$

But we have  $\begin{cases} (y_0 - y_1\sqrt{t})' = \mathcal{F}_1(U_1) \\ (y_0 + y_1\sqrt{t})' = \mathcal{F}_2(U_2) \end{cases}$

It means  $Y(X) = XU(X) \Rightarrow Y'(X) = U(X) + XU'(X) = \mathcal{F}(U)$ .

Then,

$$\begin{cases} U_1'(x_0 - x_1\sqrt{t}) = \frac{\mathcal{F}_1(U_1) - U_1(x_0 - x_1\sqrt{t})}{x_0 - x_1\sqrt{t}} \\ U_2'(x_0 + x_1\sqrt{t}) = \frac{\mathcal{F}_2(U_2) - U_2(x_0 + x_1\sqrt{t})}{x_0 + x_1\sqrt{t}} \end{cases}$$

Hence, we get ODEs for the new function with separated variables,

$$U'(X) = \frac{\mathcal{F}(U) - U(X)}{X}$$

Every solution  $U_1(X)$ ,  $U_2(X)$  to last ODEs leads to a solution  $Y_1(x_0 - x_1\sqrt{t}) = (x_0 - x_1\sqrt{t})U_1(x_0 - x_1\sqrt{t})$ ,  $Y_2(x_0 + x_1\sqrt{t}) = (x_0 + x_1\sqrt{t})U_2(x_0 + x_1\sqrt{t})$  of the homogeneous ODE (24 - 1) and (24 - 2), respectively, so the general solution is  $Y = \varphi^{-1}(Y_1, Y_2)$ .

**Example 7.**

**Let the ODE:**

$$Y' = \frac{Y}{X} - \frac{X^2}{Y^2}; X \in F_j \setminus \left\{ \left(\frac{1}{2} \mp \frac{1}{2\sqrt{t}}\right)k, 0; k \in R \right\} \tag{25}$$

$$\Rightarrow \varphi^{-1}((y_0 - y_1\sqrt{t})', (y_0 + y_1\sqrt{t})') = \varphi^{-1}\left(\frac{y_0 - y_1\sqrt{t}}{x_0 - x_1\sqrt{t}} - \left(\frac{x_0 - x_1\sqrt{t}}{y_0 - y_1\sqrt{t}}\right)^2, \frac{y_0 + y_1\sqrt{t}}{x_0 + x_1\sqrt{t}} - \left(\frac{x_0 + x_1\sqrt{t}}{y_0 + y_1\sqrt{t}}\right)^2\right)$$

$$\Leftrightarrow \begin{cases} (y_0 - y_1\sqrt{t})' = U_1 - \frac{1}{U_1^2} = \mathcal{F}_1(U_1) \\ (y_0 + y_1\sqrt{t})' = U_2 - \frac{1}{U_2^2} = \mathcal{F}_2(U_2) \end{cases}$$

Where

$$U_1(x_0 - x_1\sqrt{t}) = \frac{y_0 - y_1\sqrt{t}}{x_0 - x_1\sqrt{t}} \text{ and } U_2(x_0 + x_1\sqrt{t}) = \frac{y_0 + y_1\sqrt{t}}{x_0 + x_1\sqrt{t}} \text{ (} U(X) = \frac{Y(X)}{X} \text{),}$$

$$Y(X) = XU(X), \tag{26}$$

$$\Rightarrow Y'(X) = U(X) + XU'(X) \Leftrightarrow \begin{cases} (y_0 - y_1\sqrt{t})' = U_1 + (x_0 - x_1\sqrt{t})U_1' \\ (y_0 + y_1\sqrt{t})' = U_2 + (x_0 + x_1\sqrt{t})U_2' \end{cases}$$

Then, the ODE (25) transforms into separable ODEs:

$$\begin{cases} U_1'(x_0 - x_1\sqrt{t}) = \frac{\mathcal{F}_1(U_1) - U_1}{x_0 - x_1\sqrt{t}} = -\frac{1}{(x_0 - x_1\sqrt{t})U_1^2} \\ U_2'(x_0 + x_1\sqrt{t}) = \frac{\mathcal{F}_2(U_2) - U_2}{x_0 + x_1\sqrt{t}} = -\frac{1}{(x_0 + x_1\sqrt{t})U_2^2} \end{cases} \quad \left( U' = -\frac{1}{XU^2} \right)$$

$$\Rightarrow \begin{cases} U_1^2 dU_1 = -\frac{d(x_0 - x_1\sqrt{t})}{x_0 - x_1\sqrt{t}} \\ U_2^2 dU_2 = -\frac{d(x_0 + x_1\sqrt{t})}{x_0 + x_1\sqrt{t}} \end{cases} \quad \left( U^2 dU = -\frac{dX}{X} \right)$$

By integrating:

$$\begin{cases} \int U_1^2 dU_1 + \int \frac{d(x_0 - x_1\sqrt{t})}{x_0 - x_1\sqrt{t}} = 0 \\ \int U_2^2 dU_2 + \int \frac{d(x_0 + x_1\sqrt{t})}{x_0 + x_1\sqrt{t}} = 0 \end{cases} \quad \left( \int U^2 dU + \int \frac{dX}{X} = 0 \right)$$

$$\Rightarrow \begin{cases} \frac{U_1^3}{3} + \log(x_0 - x_1\sqrt{t}) = C_0 \\ \frac{U_2^3}{3} + \log(x_0 + x_1\sqrt{t}) = C_1 \end{cases} \quad \left( \frac{U^3}{3} + \log(X) = C \right)$$

$$\Rightarrow \begin{cases} U_1^3 = -3\log(x_0 - x_1\sqrt{t}) + 3C_0 \\ U_2^3 = -3\log(x_0 + x_1\sqrt{t}) + 3C_1 \end{cases}$$

$$\xrightarrow{\text{from(26)}} \begin{cases} Y_1^3 = ((x_0 - x_1\sqrt{t})U_1)^3 = (x_0 - x_1\sqrt{t})^3 [-3\log(x_0 - x_1\sqrt{t}) + 3C_0] \\ Y_2^3 = ((x_0 + x_1\sqrt{t})U_2)^3 = (x_0 + x_1\sqrt{t})^3 [-3\log(x_0 + x_1\sqrt{t}) + 3C_1] \end{cases}$$

$$\Rightarrow \begin{cases} Y_1 = \sqrt[3]{(x_0 - x_1\sqrt{t})^3 (-3\log(x_0 - x_1\sqrt{t}) + 3C_0)} \\ Y_2 = \sqrt[3]{(x_0 + x_1\sqrt{t})^3 (-3\log(x_0 + x_1\sqrt{t}) + 3C_1)} \end{cases}$$

The general solution is  $Y = \varphi^{-1}(Y_1, Y_2) = \sqrt[3]{X^3 (-3\log(X) + 3C)}$ .

#### 4.2. Exact type

**Definition 19.** Let  $X \in F_j$  and  $Y(X) \in F_j(X)$ , the following DE:

$$M(X, Y) + N(X, Y)Y' = 0$$

Or

$$M(X, Y)dX + N(X, Y)dY = 0 \quad (27)$$

is called an *Exact* (or *total*) *WFC-DE* when there exists a function  $U(X, Y)$  of which it is the *Exact differential*, so that  $\frac{\partial U}{\partial X} = M(X, Y)$  and  $\frac{\partial U}{\partial Y} = N(X, Y)$ , i.e.  $dU = M(X, Y)dX + N(X, Y)dY = \frac{\partial U}{\partial X}dX + \frac{\partial U}{\partial Y}dY$ , where  $U(X, Y) = C$  is the general solution.

**Remark 6.** The previous definition is equivalent to say that (27) is an exact WFC-DE when its consequent equations by using  $\varphi$  for (28 - 1), (28 - 2) are exact DEs:

$$(27) \Leftrightarrow \begin{cases} M_1(x_0 - x_1\sqrt{t}, y_0 - y_1\sqrt{t})d(x_0 - x_1\sqrt{t}) + N_1(x_0 - x_1\sqrt{t}, y_0 - y_1\sqrt{t})d(y_0 - y_1\sqrt{t}) = 0 & (28 - 1) \\ M_2(x_0 + x_1\sqrt{t}, y_0 + y_1\sqrt{t})d(x_0 + x_1\sqrt{t}) + N_2(x_0 + x_1\sqrt{t}, y_0 + y_1\sqrt{t})d(y_0 + y_1\sqrt{t}) = 0 & (28 - 2) \end{cases}$$

where  $M = \varphi^{-1}(M_1, M_2), N = \varphi^{-1}(N_1, N_2)$  and  $U = \varphi^{-1}(U_1, U_2)$ . Since,

$$\begin{aligned} dU_1 &= M_1(x_0 - x_1\sqrt{t}, y_0 - y_1\sqrt{t})d(x_0 - x_1\sqrt{t}) + N_1(x_0 - x_1\sqrt{t}, y_0 - y_1\sqrt{t})d(y_0 - y_1\sqrt{t}) \\ &= \frac{\partial U_1}{\partial(x_0 - x_1\sqrt{t})}d(x_0 - x_1\sqrt{t}) + \frac{\partial U_1}{\partial(y_0 - y_1\sqrt{t})}d(y_0 - y_1\sqrt{t}) \\ dU_2 &= M_2(x_0 + x_1\sqrt{t}, y_0 + y_1\sqrt{t})d(x_0 + x_1\sqrt{t}) + N_2(x_0 + x_1\sqrt{t}, y_0 + y_1\sqrt{t})d(y_0 + y_1\sqrt{t}) \\ &= \frac{\partial U_2}{\partial(x_0 + x_1\sqrt{t})}d(x_0 + x_1\sqrt{t}) + \frac{\partial U_2}{\partial(y_0 + y_1\sqrt{t})}d(y_0 + y_1\sqrt{t}) \end{aligned}$$

Theorem. If  $M(X, Y)$  and  $N(X, Y)$  are continuously differentiable WFC functions in a connected domain, then (27) is an exact differential WFC if and only if  $\frac{\partial N}{\partial X} = \frac{\partial M}{\partial Y}$ .

**Proof.**

We know that WFC functions  $M(X, Y)$  and  $N(X, Y)$  are continuous if and only if  $M_1, M_2, N_1, N_2$  are continuous real functions. Also, (28-1) and (28-2) are exact DEs if and only if

$$\frac{\partial N_1}{\partial(x_0 - x_1\sqrt{t})} = \frac{\partial M_1}{\partial(y_0 - y_1\sqrt{t})}, \frac{\partial N_2}{\partial(x_0 + x_1\sqrt{t})} = \frac{\partial M_2}{\partial(y_0 + y_1\sqrt{t})}, \text{ respectively.}$$

So that,

$$\begin{aligned} \frac{\partial N}{\partial X} &= \varphi^{-1}\left(\frac{\partial N_1}{\partial(x_0 - x_1\sqrt{t})}, \frac{\partial N_2}{\partial(x_0 + x_1\sqrt{t})}\right) \\ &= \frac{1}{2}\left(\frac{\partial N_1}{\partial(x_0 - x_1\sqrt{t})} + \frac{\partial N_2}{\partial(x_0 + x_1\sqrt{t})}\right) + \frac{1}{2\sqrt{t}}J\left(\frac{\partial N_2}{\partial(x_0 + x_1\sqrt{t})} - \frac{\partial N_1}{\partial(x_0 - x_1\sqrt{t})}\right) \\ &= \frac{1}{2}\left(\frac{\partial M_1}{\partial(x_0 - x_1\sqrt{t})} + \frac{\partial M_2}{\partial(x_0 + x_1\sqrt{t})}\right) + \frac{1}{2\sqrt{t}}J\left(\frac{\partial M_2}{\partial(x_0 + x_1\sqrt{t})} - \frac{\partial M_1}{\partial(x_0 - x_1\sqrt{t})}\right) \\ &= \varphi^{-1}\left(\frac{\partial M_1}{\partial(x_0 - x_1\sqrt{t})}, \frac{\partial M_2}{\partial(x_0 + x_1\sqrt{t})}\right) = \frac{\partial M}{\partial Y}. \end{aligned}$$

**Example 8.**

$$\frac{Y}{\sqrt{X}}dX + 2\sqrt{X}dY = 0$$

It is an exact WFC-DE since  $\frac{\partial N}{\partial X} = \frac{\partial M}{\partial Y}$ , because of  $\frac{\partial N_1}{\partial(x_0 - x_1\sqrt{t})} = \frac{\partial M_1}{\partial(y_0 - y_1\sqrt{t})}, \frac{\partial N_2}{\partial(x_0 + x_1\sqrt{t})} = \frac{\partial M_2}{\partial(y_0 + y_1\sqrt{t})}$ ,

where

$$\begin{aligned} M = \frac{Y}{\sqrt{X}} \Rightarrow \frac{\partial M}{\partial Y} = \frac{1}{\sqrt{X}} \Rightarrow \varphi^{-1}\left(\frac{\partial M_1}{\partial(x_0 - x_1\sqrt{t})}, \frac{\partial M_2}{\partial(x_0 + x_1\sqrt{t})}\right) &= \left(\frac{1}{\sqrt{x_0 - x_1\sqrt{t}}}, \frac{1}{\sqrt{x_0 + x_1\sqrt{t}}}\right), \\ N = 2\sqrt{X} \Rightarrow \frac{\partial N}{\partial X} = \frac{1}{\sqrt{X}} \Rightarrow \varphi^{-1}\left(\frac{\partial N_1}{\partial(x_0 - x_1\sqrt{t})}, \frac{\partial N_2}{\partial(x_0 + x_1\sqrt{t})}\right) &= \left(\frac{1}{\sqrt{x_0 - x_1\sqrt{t}}}, \frac{1}{\sqrt{x_0 + x_1\sqrt{t}}}\right). \end{aligned}$$

To find the general solution, suppose

$$dU_1 = M_1(x_0 - x_1\sqrt{t}, y_0 - y_1\sqrt{t})d(x_0 - x_1\sqrt{t}) + N_1(x_0 - x_1\sqrt{t}, y_0 - y_1\sqrt{t})d(y_0 - y_1\sqrt{t})$$

$$= \frac{y_0 - y_1\sqrt{t}}{\sqrt{x_0 - x_1\sqrt{t}}} d(x_0 - x_1\sqrt{t}) + 2\sqrt{x_0 - x_1\sqrt{t}} d(y_0 - y_1\sqrt{t})$$

$$dU_2 = M_2(x_0 + x_1\sqrt{t}, y_0 + y_1\sqrt{t}) d(x_0 + x_1\sqrt{t}) + N_2(x_0 + x_1\sqrt{t}, y_0 + y_1\sqrt{t}) d(y_0 + y_1\sqrt{t})$$

$$= \frac{y_0 + y_1\sqrt{t}}{\sqrt{x_0 + x_1\sqrt{t}}} d(x_0 + x_1\sqrt{t}) + 2\sqrt{x_0 + x_1\sqrt{t}} d(y_0 + y_1\sqrt{t})$$

$$dU = M(X, Y) dX + N(X, Y) dY = \frac{Y}{\sqrt{X}} dX + 2\sqrt{X} dY = \frac{\partial U}{\partial X} dX + \frac{\partial U}{\partial Y} dY$$

where  $M = \varphi^{-1}(M_1, M_2)$ ,  $N = \varphi^{-1}(N_1, N_2)$  and  $U = \varphi^{-1}(U_1, U_2)$

$$\Rightarrow U(X, Y) = \int \frac{Y}{\sqrt{X}} dX + \int 2\sqrt{X} dY = 2Y\sqrt{X} ; X > 0.$$

### 4.3. Linear type

Definition 20. Let  $Y$  is the WFC dependent variable, the following DE is called the standard form of a linear first-order WFC-DE in the dependent WFC variable  $Y$ :

$$Y' + P(X)Y = Q(X) \quad (29)$$

assuming that the two given functions  $P(X)$  and  $Q(X)$  are both continuous on the interval  $I = \varphi^{-1}(I_1, I_2)$ .

$$(29) \Leftrightarrow \begin{cases} (y_0 - y_1\sqrt{t})' + (y_0 - y_1\sqrt{t})P_1(x_0 - x_1\sqrt{t}) = Q_1(x_0 - x_1\sqrt{t}) & (30-1) \\ (y_0 + y_1\sqrt{t})' + (y_0 + y_1\sqrt{t})P_2(x_0 + x_1\sqrt{t}) = Q_2(x_0 + x_1\sqrt{t}) & (30-2) \end{cases} \quad (30)$$

where  $Q = \varphi^{-1}(Q_1, Q_2)$ ,  $P = \varphi^{-1}(P_1, P_2)$ ,  $Q_1, P_1$  are continuous functions on  $I_1$  and  $Q_2, P_2$  are continuous functions on  $I_2$ .

Remark 7. When  $Q(X) = 0$ , the linear equation (29) is said to be homogeneous; otherwise, it is nonhomogeneous, i.e.,  $Q_1(x_0 - x_1\sqrt{t}) = 0$  and  $Q_2(x_0 + x_1\sqrt{t}) = 0$ . Such that, we have two homogeneous DEs:

$$Y' + P(X)Y = 0 \Leftrightarrow \begin{cases} (y_0 - y_1\sqrt{t})' + (y_0 - y_1\sqrt{t})P_1(x_0 - x_1\sqrt{t}) = 0 & (31-1) \\ (y_0 + y_1\sqrt{t})' + (y_0 + y_1\sqrt{t})P_2(x_0 + x_1\sqrt{t}) = 0 & (31-2) \end{cases} \quad (31)$$

Definition 21. Finding  $Y$  the solution to the WFC linear ODE (29) is equivalently to find the solutions  $y_0 - y_1\sqrt{t}$ ,  $y_0 + y_1\sqrt{t}$  to the linear ODEs (30-1), (30-2), respectively, where

$$Y = \varphi^{-1}(y_0 - y_1\sqrt{t}, y_0 + y_1\sqrt{t}).$$

### Find the General Solution to the Linear First-order WFC-ODE:

$$\text{FIRST STEP: Solve the homogeneous equation } Y' + P(X)Y = 0 \quad (32)$$

To do that we have to solve the homogeneous equations (31-1) and (31-2) which are equivalent to the following separable DEs:

$$\begin{cases} \frac{d(y_0 - y_1\sqrt{t})}{y_0 - y_1\sqrt{t}} + P_1 d(x_0 - x_1\sqrt{t}) = 0 & (33-1) \\ \frac{d(y_0 + y_1\sqrt{t})}{y_0 + y_1\sqrt{t}} + P_2 d(x_0 + x_1\sqrt{t}) = 0 & (33-2) \end{cases} \quad (33)$$

by integrating and solving (33-1) and (33-2) for  $y_0 - y_1\sqrt{t}$  and  $y_0 + y_1\sqrt{t}$ , respectively:

$$\begin{cases} Y_{h1} = C_1 e^{-\int P_1(x_0 - x_1\sqrt{t}) d(x_0 - x_1\sqrt{t})} \\ Y_{h2} = C_2 e^{-\int P_2(x_0 + x_1\sqrt{t}) d(x_0 + x_1\sqrt{t})} \end{cases} \\ \Rightarrow Y_h = \varphi^{-1}(Y_{h1}, Y_{h2}) = \varphi^{-1}(C_1, C_2) \varphi^{-1}(e^{-\int P_1 d(x_0 - x_1\sqrt{t})}, e^{-\int P_2 d(x_0 + x_1\sqrt{t})})$$

After calculating:

$$C = \frac{1}{2}[C_1 + C_2] + \frac{1}{2\sqrt{t}}J[C_2 - C_1]$$

$$e^{-\int P(X)d(X)} = \frac{1}{2}[e^{-\int P_1 d(x_0-x_1\sqrt{t})} + e^{-\int P_2 d(x_0+x_1\sqrt{t})}] + \frac{1}{2\sqrt{t}}J[e^{-\int P_2 d(x_0+x_1\sqrt{t})} - e^{-\int P_1 d(x_0-x_1\sqrt{t})}]$$

We find that the solution to homogeneous WFC-DE (32) is as follows:

$$Y_h = C e^{-\int P(X)dX}$$

**SECOND STEP:** Find the particular solutions to the nonhomogeneous equations (30 – 1) and (30 – 2), respectively are:

$$\begin{cases} Y_{r1} = e^{-\int P_1(x_0-x_1\sqrt{t})d(x_0-x_1\sqrt{t})} \int Q_1(x_0 - x_1\sqrt{t}) e^{\int P_1(x_0-x_1\sqrt{t})d(x_0-x_1\sqrt{t})} d(x_0 - x_1\sqrt{t}) \\ Y_{r2} = e^{-\int P_2(x_0+x_1\sqrt{t})d(x_0+x_1\sqrt{t})} \int Q_2(x_0 + x_1\sqrt{t}) e^{\int P_2(x_0+x_1\sqrt{t})d(x_0+x_1\sqrt{t})} d(x_0 + x_1\sqrt{t}) \end{cases}$$

$$\Rightarrow Y_r = \varphi^{-1}(Y_{r1}, Y_{r2}) = e^{-\int P(X)dX} \int Q(X) e^{\int P(X)dX} dX$$

**THIRD STEP:** We know that the general solutions to each (30 – 1) and (30 – 2) is the sum of two solutions:

$$\begin{cases} y_0 - y_1\sqrt{t} = Y_{h1} + Y_{r1} \\ y_0 + y_1\sqrt{t} = Y_{h2} + Y_{r2} \end{cases}$$

Then, the general solution to (29) can be written as the sum of two solutions:

$$Y = Y_h + Y_r$$

$$\text{where } = \varphi^{-1}(y_0 - y_1\sqrt{t}, y_0 + y_1\sqrt{t}),$$

$Y_h = \varphi^{-1}(Y_{h1}, Y_{h2})$  is the solution of the associated homogeneous equation (32) and

$Y_r = \varphi^{-1}(Y_{r1}, Y_{r2})$  is a particular solution to the nonhomogeneous (29).

### Example 9

Let the WFC-DE is  $XY' - 2Y - X^3e^X = 0$

We can write it as follows:

$$Y' - \frac{2}{X}Y = X^2e^X.$$

It is a Linear 1<sup>st</sup> order WFC-DE, we can solve it using the previous steps, where

$$P(X) = -\frac{2}{X} = \varphi^{-1}\left(-\frac{2}{x_0 - x_1\sqrt{t}}, -\frac{2}{x_0 + x_1\sqrt{t}}\right)$$

$$Q(X) = X^2e^X = \varphi^{-1}\left((x_0 - x_1\sqrt{t})^2 e^{x_0-x_1\sqrt{t}}, (x_0 + x_1\sqrt{t})^2 e^{x_0+x_1\sqrt{t}}\right)$$

The solution to homogeneous associated DE is  $Y_h(X) = \varphi^{-1}(Y_{h1}, Y_{h2}) = CX^2$

$$\text{where } \begin{cases} Y_{h1} = C_1 e^{\int \frac{2}{x_0-x_1\sqrt{t}} d(x_0-x_1\sqrt{t})} = C_1 (x_0 - x_1\sqrt{t})^2 \\ Y_{h2} = C_2 e^{\int \frac{2}{x_0+x_1\sqrt{t}} d(x_0+x_1\sqrt{t})} = C_2 (x_0 + x_1\sqrt{t})^2 \end{cases}; C = \varphi^{-1}(C_1, C_2)$$

The particular solution to nonhomogeneous associated DE is

$$Y_r(X) = e^{\int X^2 dX} \int X^2 e^X e^{-\int X^2 dX} dX = X^2 \int e^X dX = X^2 e^X$$

where,

$$\begin{cases} Y_{r1} = e^{\int \frac{2}{x_0-x_1\sqrt{t}} d(x_0-x_1\sqrt{t})} \int (x_0 - x_1\sqrt{t})^2 e^{x_0-x_1\sqrt{t}} e^{-\int \frac{2}{x_0-x_1\sqrt{t}} d(x_0-x_1\sqrt{t})} d(x_0 - x_1\sqrt{t}) = (x_0 - x_1\sqrt{t})^2 e^{x_0-x_1\sqrt{t}} \\ Y_{r2} = e^{\int \frac{2}{x_0+x_1\sqrt{t}} d(x_0+x_1\sqrt{t})} \int (x_0 + x_1\sqrt{t})^2 e^{x_0+x_1\sqrt{t}} e^{-\int \frac{2}{x_0+x_1\sqrt{t}} d(x_0+x_1\sqrt{t})} d(x_0 + x_1\sqrt{t}) = (x_0 + x_1\sqrt{t})^2 e^{x_0+x_1\sqrt{t}} \end{cases}$$

The general solution to the given DE is

$$\begin{aligned} Y &= Y_h + Y_r \\ &= CX^2 + X^2 e^X. \end{aligned}$$

## 5. Conclusion

In this paper, we have defined the Weak Fuzzy Complex Differential Equation depending on a special isomorphic transformation function which helps us to deal easily with two ODEs in  $R$  then come back to  $F_j$  to get the general solution to WFC-ODEs. Therefore, we introduce some types of WFC-ODEs of first-order first-degree (Separable - Exact - Linear) with their general solutions and some explicit examples. In the future, we aim to solve WFC Initial Value Problems.

## References

- [1] M. Abualhomos, W.M.M. Salameh, M. Bataineh, M.O. Al-Qadri, A. Alahmade, A. Al-Husban, An Effective Algorithm for Solving Weak Fuzzy Complex Diophantine Equations in Two Variables, (2024). <https://doi.org/10.54216/IJNS.230431>.
- [2] A.M.A. Alfahal, M. Abobala, Y.A. Alhasan, R.A. Abdulfatah, Generating Weak Fuzzy Complex and Anti Weak Fuzzy Complex Integer Solutions for Pythagoras Diophantine Equation  $X^2 + Y^2 = Z^2$ , International Journal of Neutrosophic Science 22 (2023) 8–14. <https://doi.org/10.54216/IJNS.220201>.
- [3] Y.A. Alhasan, A.M.A. Alfahal, R.A. Abdulfatah, G. Nordo, M.M.A. Zahra, On Some Novel Results About Weak Fuzzy Complex Matrices, International Journal of Neutrosophic Science 21 (2023) 134–140. <https://doi.org/10.54216/IJNS.210112>.
- [4] Y.A. Alhasan, L. Xu, R.A. Abdulfatah, A.M.A. Alfahal, The Geometrical Characterization for The Solutions of a Vectorial Equation By Using Weak Fuzzy Complex Numbers and Other Generalizations Of Real Numbers, International Journal of Neutrosophic Science 21 (2023) 155–159. <https://doi.org/10.54216/IJNS.210415>.
- [5] R. Ali, On The Weak Fuzzy Complex Inner Products On Weak Fuzzy Complex Vector Spaces, Neoma Journal of Mathematics and Computer Science 1 (2023). <https://doi.org/10.5281/zenodo.7953682>.
- [6] G. Birkhoff, G.C. Rota, ORDINARY DIFFERENTIAL EQUATIONS, Wiley, 1991.
- [7] F.C. Galarza, M.L. Flores, D.P. Rivero, M. Abobala, On Weak Fuzzy Complex Pythagoras Quadruples, International Journal of Neutrosophic Science 22 (2023) 108–113. <https://doi.org/10.54216/IJNS.220209>.
- [8] W. Gander, M.J. Gander, F. Kwok, Scientific computing-An Introduction using Maple and MATLAB, Springer, 2014.
- [9] A. Hatip, an Introduction to Weak Fuzzy Complex Numbers, Galoitica: Journal of Mathematical Structures and Applications 3 (2023) 08–13. <https://doi.org/10.54216/gjmsa.030101>.
- [10] L. Razouk, S. Mahmoud, M. Ali, On the Foundations of Weak Fuzzy Complex-Real Functions, 5 (2024) 116–140, <https://doi.org/10.22105/jfea.2024.435955.1369>.
- [11] L. Razouk, S. Mahmoud, M. Ali, A Computer Program For The System Of Weak Fuzzy Complex Numbers And Their Arithmetic Operations Using Python, Galoitica: Journal of Mathematical Structures and Applications 8 (2023) 45–5 <https://doi.org/10.54216/gjmsa.080104>.
- [12] Abdallah Shihadeh, Khaled Ahmad Mohammad Matarneh, Raed Hatamleh, Randa Bashir Yousef Hijazeen, Mowafaq Omar Al-Qadri, Abdallah Al-Husban.(2024). An Example of Two-Fold Fuzzy Algebras Based on Neutrosophic Real Numbers. Neutrosophic Sets and Systems, vol.67, pp. 169-178. <https://doi.org/10.5281/zenodo.11151930>.
- [13] Abdallah Shihadeh, Khaled Ahmad Mohammad Matarneh, Raed Hatamleh, Mowafaq Omar Al-Qadri, Abdallah Al-Husban.(2024), On The Two-Fold Fuzzy n-Refined Neutrosophic Rings For  $2 \leq n \leq 3$ , Neutrosophic Sets and Systems, Vol. 68, pp.8-25. <https://doi.org/10.5281/zenodo.11406449>.
- [14] W. Walter, Ordinary Differential Equations, Springer New York, 1998. <https://doi.org/10.1007/978-1-4612-0601-9>.
- [15] D.G. Zill, a First Course in Differential Equations with Modeling Application, Brooks/Cole, Cengage Learning., 2008.
- [16] Heilat, A. S. (2023). Cubic Trigonometric B-spline Method for Solving a Linear System of Second Order Boundary Value Problems. European Journal of Pure and Applied Mathematics, 16(4), 2384-2396. <https://doi.org/10.29020/nybg.ejpm.v16i4.4947>
- [17] Heilat, A. S., & Hailat, R. S. (2020). Extended cubic B-spline method for solving a system of non-linear second-order boundary value problems. J. Math. Comput. Sci, 21, 231-242. <http://dx.doi.org/10.22436/jmcs.021.03.06>.
- [18] Heilat, A.S., Qazza, A., Hatamleh, R., Saadeh, R., Alomari, M.W. (2023). An application of Hayashi's inequality in numerical integration. Open Mathematics, 21(1), 20230162. <https://doi.org/10.1515/math-2023-0162>.
- [19] Heilat, A.S., Batiha, B., Qawasmeh, Hatamleh R. (2023). Hybrid Cubic B-spline Method for Solving A Class of Singular Boundary Value Problems. European Journal of Pure and Applied Mathematics, 16(2), 751-762. <https://doi.org/10.29020/nybg.ejpm.v16i2.4725>.

- [20] Ayman Hazaymeh, Ahmad Qazza, Raed Hatamleh, Mohammad W Alomari, Rania Saadeh.(2023). On Further Refinements of Numerical Radius Inequalities, *Axiom-MDPI*, 12(9), 807.
- [21] T.Qawasmeh, A. Qazza, R. Hatamleh, M.W. Alomari, R. Saadeh. (2023). Further accurate numerical radius inequalities, *Axiom-MDPI*, 12 (8), 801.
- [22] S. Hussain, S. Khan, and M. Younis, "Numerical simulation of sine-Gordon equation using cubic B-spline collocation method," *Applied Numerical Mathematics*, vol. 158, pp. 240–252, 2021, doi: 10.1016/j.apnum.2020.09.012.
- [23] M. A. Al-Sharif, H. M. Al-Masri, and F. A. Al-Hamadi, "Numerical Methods for Solving Fuzzy Differential Equations with Applications," *American Journal of Business and Operations Research*, vol. 8, no. 1, pp. 45-56, 2021. doi:10.1234/ajbor.2021.045.
- [24] Alrwashdeh, Dima. , Alkhouli, Talat. , Soiman, Ahmed. , Allouf, Ali. , Edduweh, Hussein. , Al-Husban, Abdallah.(2025). On Two Novel Generalized Versions of Diffie-Hellman Key Exchange Algorithm Based on Neutrosophic and Split-Complex Integers and their Complexity Analysis. *International Journal of Neutrosophic Science*, 25(2), pp. 01-10.
- [25] Abubaker, A. A., Abualhomos, M., Matarneh, K., & Al-Husban, A. (2025). A Numerical Approach For The Algebra of Two-Fold. *Neutrosophic Sets and Systems*, 75, 181-195.
- [26] A., Ahmad. , Hatamleh, Raed. , Matarneh, Khaled. , Al-Husban, Abdallah. On the Irreversible k-Threshold Conversion Number for Some Graph Products and Neutrosophic Graphs. (2025). *International Journal of Neutrosophic Science*, 25(2), pp. 183-196.
- [27] Al-Husban, A., Djenina, N., Saadeh, R., Ouannas, A., & Grassi, G. (2023). A new incommensurate fractional-order COVID 19: modelling and dynamical analysis. *Mathematics*, 11(3), 555.
- [28] M. Alkahtani, M. Alsaedi, and F. A. Rihan, "A fractional-order model of COVID-19 with quarantine, isolation, and environmental viral load," *Chaos, Solitons & Fractals*, vol. 140, p. 110220, 2020, doi: 10.1016/j.chaos.2020.110220.
- [29] Hamadneh, T., Hioual, A., Alsayyed, O., Al-Khassawneh, Y. A., Al-Husban, A., & Ouannas, A. (2023). The FitzHugh–Nagumo Model Described by Fractional Difference Equations: Stability and Numerical Simulation. *Axioms*, 12(9), 806.
- [30] Heilat, A. S., Karoun, R. C., Al-Husban, A., Abbas, A., Al Horani, M., Grassi, G., & Ouannas, A. (2023). The new fractional discrete neural network model under electromagnetic radiation: Chaos, control and synchronization. *Alexandria Engineering Journal*, 76, 391-409.
- [31] Al-Husban, A., Karoun, R. C., Heilat, A. S., Al Horani, M., Khennaoui, A. A., Grassi, G., & Ouannas, A. (2023). Chaos in a two dimensional fractional discrete Hopfield neural network and its control. *Alexandria Engineering Journal*, 75, 627-638.
- [32] Abu Falahah, I., Hioual, A., Al-Qadri, M. O., Al-Khassawneh, Y. A., Al-Husban, A., Hamadneh, T., & Ouannas, A. (2023). Synchronization of Fractional Partial Difference Equations via Linear Methods. *Axioms*, 12(8), 728.
- [33] Abu Falahah, I., Hioual, A., Al-Qadri, M. O., Al-Khassawneh, Y. A., Al-Husban, A., Hamadneh, T., & Ouannas, A. (2023). Synchronization of Fractional Partial Difference Equations via Linear Methods. *Axioms*, 12(8), 728.
- [34] Hamadneh, T., Abbas, A., Falahah, I. A., Al-Khassawneh, Y. A., Heilat, A. S., Al-Husban, A., & Ouannas, A. (2023). Complexity and chaos analysis for two-dimensional discrete-time predator–prey Leslie–Gower model with fractional orders. *Axioms*, 12(6), 561.
- [35] Hamadneh, T., Hioual, A., Alsayyed, O., Al-Khassawneh, Y. A., Al-Husban, A., & Ouannas, A. (2023). Finite Time Stability Results for Neural Networks Described by Variable-Order Fractional Difference Equations. *Fractal and Fractional*, 7(8), 616.
- [36] Abu Falahah, I., Hioual, A., Al-Qadri, M. O., Al-Khassawneh, Y. A., Al-Husban, A., Hamadneh, T., & Ouannas, A. (2023). Synchronization of Fractional Partial Difference Equations via Linear Methods. *Axioms*, 12(8), 728.