



Utilizing Machine Learning for Predicting Lyme Disease Trends and Enhancing Diagnostic Accuracy

Ahmed El-Sayed Saqr^{1,*} Ahmed M. Elshewey²

¹ Computer Engineering and Control Systems Department, Faculty of Engineering, Mansoura University, Egypt

² Faculty of Computers and Information, Computer Science Department, Suez University, Egypt

Emails: a7mdsq@std.mans.edu.eg · ahmed.elshewey@fci.suezuni.edu.eg

Received: September 17, 2023 Revised: December 16, 2023 Accepted: January 18, 2024 ★ Corresponding author

ABSTRACT

The present research investigates the role of machine learning models in forecasting the course of Lyme disease and improving diagnostics by looking for environmental, host and anthropogenic factors contributing to the rise and fall of the tick population and disease outbreaks. With the popularization of ecological models and artificial intelligence-based techniques such as neural networks and random forests, it has become possible to efficiently and accurately over various risk maps that relate to ticks' location and distribution, which is an essential aspect of improving public health management issues. These models integrate climate and demographic data as well as host-pathogen interaction data and help understand the distribution of high-risk areas and the dynamics of the diseases, thus facilitating the management of tick-borne illness. This approach also illustrates the significance of predictive diagnostics for early disease detection, allowing for interventions and preventive measures only on relevant population sub-groups. Ultimately, this study considers the possibilities machine learning offers in managing Lyme disease, articulating the implications of these conclusions for the preparedness for health emergencies on a more global scale.

Keywords: Automated learning ▪ Lyme borreliosis ▪ Forecasting actual behavior ▪ Health concerns transmitted by insects ▪ Tests and procedures ▪ Community health

1. INTRODUCTION

Tick-borne diseases have become increasingly important in recent decades; significant changes in tick distribution and pathogen occurrence have complicated the disease as a global public health challenge and a threat to ecological systems. Mites, particularly those implicated in the transmission of diverse pathogens, are threatened by such environmental factors because the extent of their spread and repopulation is a function of environmental change. This sensitivity has spurred the need to work more on the population of the ticks to determine the space-time interaction of disease threat. Although previous research has shown that tick ranges extend globally, new

data prove that climatic change, altered habitats, and human encroachment of green spaces impact the distribution and increase people's vulnerability to tick-borne diseases such as Lyme disease and tick-borne encephalitis. This literature review highlights these environmental and host-related factors that lead to tick and disease burdens; also presented are the epidemiological models employed in examining risk [1].

The link between environmental factors and ticks, especially *Ixodes scapularis* and *Ixodes ricinus*, is prominent in literature from the last century. Abiotic factors like temperature, rainfall and vegetation cover strongly correlate with tick occurrence rate/distribution. Several papers employ the techniques of predictive models for these factors, showing

that higher temperatures and higher humidity increase the prospects for ticks' survival and reproduction, thus increasing disease prevalence. Besides climate factors, variations in land use have been cited as factors of increased dissemination of ticks as natural ecosystems, shrink and wildlife carriers and ticks are pushed into human-dominated areas. Such findings help predict the risk concentrations and develop preventive measures for diseases carried by ticks that would not harm human health [2], [3].

The recent development of computational approaches such as machine learning and predictive data analytics has enabled researchers to effectively survey and map the distribution of ticks and the associated disease burden. In particular, machine learning models such as artificial neural networks (ANNs), random forest (RF), and generalized boosted methods (GBM) have been utilized to explore the effects of environmental factors on tick presence and abundance. The distribution of life-stage ticks and their respective environmental niches has also been modeled successfully using these algorithms, thus making it possible to better estimate the risk for diseases associated with ticks. These models also efficiently assess the burden of tick and tick-borne disease DALYs across different countries. They help strengthen health systems in countries undergoing environmental changes. In addition to these, they help design spatial risk maps that help direct future methodological and advocacy action [4], [5].

Host-vector interactions are another critical component in transmitting diseases caused by ticks. In addition, it should be noted that much attention is paid to studying host behavior, e.g., small mammals like mice, who are primary hosts for tick larvae. Host density and behavior were shown to play a role in disease transmission dynamics. According to studies, the environmental context plays a vital role in determining host population densities, which, in turn, influences the associated tick density and infection rates. For instance, high host densities are often obtained at relatively warmer temperatures and in wet conditions, which has a knock-on effect on ticks and the diseases they spread. This knowledge of host dynamics is necessary because it informs the possible interventions: habitat alteration (for example, removing vegetation to decrease the amount of moisture), host control (to lower the number of possible disease reservoirs), and so on, all of which aim at minimizing the number of ticks human beings encounter as well as the risk of the diseases they carry [6], [7].

How people behave and their measures weigh significantly in controlling diseases such as tick-borne diseases. In New York and other areas of Europe with high tick activity, for example, prevention campaigns include wearing protective clothing, applying tick repellents, and carrying out regular tick checks after outdoor activities. However, these are not sufficient because most tick-borne illnesses, including Lyme disease, do not have a human vaccine, making it hard to prevent them. Hence, public education and behavioral modification approaches are becoming more critical in addition to environmental and ecological approaches. Most programs that raise awareness and teach the population about early signs of such illnesses have been found to decrease the incidences of infections with these contagious diseases because the people seek treatment and preventive measures in good time [8], [9].

Lastly, such improvements in diagnosis mean we can now detect more tick-borne diseases, thus strengthening the clinical and epidemiological response to outbreaks. New strategies for diagnosing diseases have led to the implementation of molecular techniques, which have greatly improved the detection of infectious agents. Such techniques are vital for evaluating potential risks of infection and the epidemiology of infections within populations. Such techniques have improved the risk of detecting virulence factors in pathogens and following their evolution in different pathogens due to the emergence of new strains. In this consideration, the progress made in diagnostics of ticks and identifying their pathogens will be valuable for developing new treatment strategies implemented in various parts of the world within different tick species.

The present body of work discusses the need for a holistic approach involving environmental surveillance, host management, public health measures, and modern diagnostics to address tick-borne diseases. These diseases can be addressed through research and policy measures incorporating information on environmental, ecological, and compartmental activities. Moreover, as global tick populations rise due to climatic and anthropogenic ecological changes, there is still a need to adopt a coordinated and multifaceted approach to mitigate the burden of tick-borne diseases on the population.

2. LITERATURE REVIEW

Tick-borne diseases have reemerged due to the changing dynamics of ticks and the pathogens they transmit, which are global public health threats. Reviews of the literature about tick distribution and pathogen prevalence demonstrate the inherent interaction among environmental, host, and geographical factors that affect tick distribution and infection rates in their hosts. Most studies have applied machine learning and ecological models to improve the predictive abilities of the distribution of ticks, the abundance of nymphs, and the areas of disease risk, which supported the public health authorities in preventing and controlling emerging health threats. Moreover, the studies focusing on environmental metrics, host-pathogen interactions and pathogen-host protein interactions make clear that vector-borne diseases are complex and multilayered. As diagnosis, controlling tactics, and prophylactic measures are at the center of attention, these results become invaluable tools to enhance the capacities of disease treatment throughout regions and provide the basis for further globalization of health policies.

In this context, in the analysis presented in [10], many species have shifted their geographical range and abundance in the comparatively recent past. The changes in the geographic ranges or population sizes of disease vectors, including ticks, are a great concern to public health as they indicate the emergence of infectious diseases in human populations. Nonetheless, one of the most significant analytical difficulties was the accurate assessment of the environmental metric drivers of the biogeography of vector species because of insufficient data sampling during demographic transition. This study, therefore, sought to investigate the pattern and distribution of *Ixodes scapularis* nymphal ticks across these 532 sampling sites in New York State (NY), USA, from 2008 to 2018, during which the occurrence of tick-borne diseases

was noticed. It was observed that ticks spread northward and westward, and the nymphal tick abundance increased in new colonization areas, but there was no significant change in tick nymphal abundance in the old colonization area of the last decade. These observations were consistent with the geographical distribution of human Lyme disease and higher risk rates. The investigated data provided strict temporal and spatial dependencies, as well as environmental variables as the drivers of the observed changes, and the development of the models allowed future trends of nymphal populations to be predicted. This forecasting ability can complement public health planning and policy change to avoid or reduce the risk of tick-transmitted diseases before the tick-transmission period.

As described in [11], white-footed mice are the primary hosts of immature blacklegged ticks and the most efficient reservoirs of multiple tick-borne pathogens, including Lyme disease agents in eastern North America. It was established that examination across the body part of an individual mouse shows that the distribution of larval ticks is quite skewed, with a small subset of hosts being responsible for most host-to-tick transmission. The researchers collected data on larval burdens from mice trapped in Millbrook, NY, over 20 years and used the information to construct an ANN model to identify which mouse or environmental characteristics correspond to heavy parasite loads. It was observed that host climatic conditions, including positive deviations in temperature and precipitation coupled with low host density, were the most influential predictors of high L3 burden overriding other host factors. This means that environmental conditions ought to dictate the course of the control of these bugs and not the features of the hosts. In addition, the study also emphasized the interactions between different variables in the predictive models, as shown by the usefulness of the ANNs in tracking the pattern of aggregation of the ticks and other related vectors, especially under conditions of prevailing global warming.

In the studies in [12], tick-borne diseases have received more attention since the tick vector and its associated pathogens have expanded their geographical range and are now posing significant threats to European human and animal health. Given the importance of predicting the number of tick larvae to understand the epidemiology of these infections, a better method for mapping these diseases is needed. The study estimated *Ixodes ricinus* distribution at 159 sites across southern Scandinavia in August – September 2016. Supervised machine learning models were trained and tested to analyze accurate field data and environmental variables based on data from 2017. The predictive models of larva and nymphs' abundance appeared to have moderately high accuracy (normalized RMSE, 0.65–0.69 and R^2 , 0.52–0.58), whereas those of the adult ticks were less accurate (normalized RMSE, 0.94–0.96 and R^2 , 0.04–0.10). Nevertheless, when the models were tested on hold-out data from 2017, the normalized RMSE values varied between 0.59 and 1.13, and the R^2 between 0.18 and 0.69. Models, mainly driven by temperature and vegetation, reached the point that climate plays a more critical role in the distribution and density of *I. ricinus*. These models are preliminary steps for environmentally based tick abundance models to delineate risk areas and enhance the interpretation of human incidence rates. As outlined in

paper [13], Lyme disease received the first guideline in 2020 from the Infectious Diseases Society of America (IDSA) for the first time in collaboration with the American Academy of Neurology and the American College of Rheumatology. Lyme disease is considered the most common tick-borne disease in the USA, with approximately 476,000 clients being treated annually, with New England, the mid-Atlantic and the north-central zone being the most affected zones. Most Lyme disease-infected people get the tick-borne disease through the *Ixodes scapularis* family of ticks or deer ticks. In the western states of California, Oregon and Washington, *Ixodes pacificus* belongs to several states but makes up less than 1% of the disease. Since there is no available human vaccine, prevention measures are mainly aimed at preventing contact with *I. scapularis*. Proper prevention strategies include using protective clothing, using or applying repellents, treating clothing with permethrin, conducting thorough tick check-ups after one is done with outdoor activities and removing ticks, as *Borrelia burgdorferi* transmission needs the tick at least 36 hours attached to the person's skin. For tick bites that require prophylaxis, one dose of oral doxycycline can be given within 72 hours after tick removal to prevent Lyme disease. These include the identification of the tick as an *Ixodes* species or exposure to a Lyme disease endemic area and a history of prolonged attachment of the tick to the skin. Specific testing of removed ticks to pathogens is not advised and does not guarantee or correlate with human infections.

As pointed out in the study [14], People are infected with Lyme disease from the bites of infected blacklegged ticks known as *Ixodes scapularis* in eastern North America; the vector is *Borrelia burgdorferi*. Administering antibiotics within 72 hours of a tick bite effectively prevents Lyme disease, suggesting the need to detect and treat the tick bite early. The work also briefly discusses medical imaging, a non-incision diagnostic method for examining internal organs and abnormalities. Filters used in medical images play a vital role in diagnostics, and image denoising is one of the primary stages of these processing algorithms, enhancing the efficiency of subsequent stages that include segmentation, registration, and integration of data from different modalities. The paper then restricted its comparisons to different projects dealing with image denoising, especially interventions that deal with the Rician noise problems in MRI. The performance of various spatial and transformation filters applied to remove noise from the acquired brain MRIs is compared to the segmentation outcomes of the proposed techniques based on performance and statistical measures. Image denoising is highlighted in the role of the pre-processing stage and as a process necessary to increase the accuracy of many image-processing methods.

In the publication [15], authors reported advanced diagnostics of tick-borne diseases due to increased concern about using control measures to reduce possible risks. Optimal control strategies are a valuable approach to defining the best practices for mitigating risks, considering the related costs. Crediting a model developed earlier, the study assesses different frameworks to identify factors affecting the mitigation measures. The frequency and duration of tick-reduction efforts are essential targets for reducing the risk of tick-borne diseases. The results show that the discontinuous-fashioned

life cycle of the tick is considerable for protecting risk, which means it can be controlled efficiently without requiring a continual treatment program. Thus, the same strategies are obtained in a closed optimal-control setting, independent of the choice of the control framework and objective function, proving that tick-borne diseases are near-zero risks without eliminating the tick population.

In the study done in [16], the author noted that controlling Lyme disease is a massive problem because the tick *Ixodes scapularis* involved is complicated. Many reservoir hosts within the ecosystem have differing levels of competence. Future work to predict the benefits of either TCTs or HCTs and to plan and execute these control measures should be contingent on understanding the correlation between pathogen risk in nymph populations, nymph density, and reported human Lyme disease cases. This study sought to establish the direct relationship between estimated acarological risk and human cases by comparing the results of the present study with human case data at the county level and nymphal prevalence rates obtained from field rates in 36 eastern states. The present study established a positive linear regression of a considerable nature between human incidence and even firmer substantive correlations in areas characterized by high incidence levels. Nevertheless, this correlation changed depending on the region and state of the United States and because of the further distribution of *B. burgdorferi* genotypes. It is established that more data is required in several high-prevalence states for broad cost-effectiveness evaluations.

In the research conducted in [17], the tourism growth linked with nature and recreation in China implied an opportunity to see green areas more frequently and, as a result, become infected with tick-borne diseases more often. This study intends to identify behavioral and cognitive perimeters singularly associated with TBE risk, essential in designing prevention and control strategies to help public health practices advance in high-incidence regions. Due to high tick concentrations, a structured self-administered questionnaire was developed and completed by 3000 residents from Heilongjiang, Jilin, and Liaoning provinces of China. This study aimed to identify participants' history of tick bites or their presence, their attitude towards possible risks of tick-borne disease, frequency of outdoor activity, etc. Structural equation modeling was used to establish the pattern of the relationship and the magnitude of these factors. The results revealed that participants had an average of self-reported tick bites that equaled 14% and tick-borne encephalitis had the highest self-reported rate of infection percentages, equaled 4%; the percentage of Lyme disease was 2%, and Severe Fever with Thrombocytopenia Syndrome was also 2%; Moreover, according to the respondents, the pet tick bite rate was 14% for ear, back and abdomen areas. It was strongly noted that tick bite rates were significantly associated with the infection rates, placing the factor at a central position of the infectious status. The study further showed that the risk factors of tick-borne diseases for infections were also different in various cities, with a focus on ecological characteristics of the city and regions' knowledge about tick prevention. They partially contribute to understanding present-day infection rates of tick-borne diseases in northeast China and reveal behavioral and cognitive patterns uninvestigated before. They helped make a prognosis about

the future effect of knowledge sharing and improved urban amenities to prevent tick bites and tick-borne diseases.

As described in [18], the Asian long-horned tick (*Hemoptysis longicornis*) transmits multiple arboviral and bacterial pathogens within its native range in East Asia and newly in Australasia. This species has bisexual and parthenogenetic forms, which preferred eating phytoplankton and had the potential to achieve high-density replies under suitable conditions. After identifying the geographic distribution of established parthenogenetic populations of *H. longicornis* for the eastern United States in 2017, climate suitability maps have indicated its probable range across North America. Nonetheless, factors influencing the distribution of parasites at the regional level, such as geology, soil type and land use, have yet to be well researched. This study used Maxent, Random Forest (RF), and Generalized Boosting Method (GBM) algorithms in ecological niche modeling to predict *H. longicornis* occurrence in New Jersey depending on environmental conditions. Research results suggest that *H. longicornis* favors the Piedmont physiographic province region and certain varieties of owned soil. This indicated that soil hydraulic conductivity was the variable most descriptive of the patterns observed and that sand habitats were less suitable than those in clay or loam soils because of high hydraulic conductivity in sandy soils. The model used yielded a sensitivity of 87% for the 2017–2019 samples to generate a probabilistic habitat suitability map of New Jersey at a finer spatial resolution of 90 m by 90m. The model fit was slightly lower for the 2020–2022 data at 57%, indicating either movement into substandard habitats or some fluctuation in environmental conditions at the fringes of physiographic provinces. Subsequent modeling should include tick abundance and population stages and the specific characteristics of the soil at the sampling sites that could offer valuable information to improve the prevention and eradication strategies of *H. longicornis*.

As described in [19], AAA is characterized by the pathological, progressive dilatation of the infrarenal abdominal aorta; therefore, identifying essential feature genes in AAA is crucial. SDS-PAGE analysis of the different ECM proteins in AAA and NT caused by abdominal aortic aneurysm identified 43 upregulated and 32 downregulated DEGs. KEGG database integration associated these DEGs with inflammation and immune response pathways using functional, pathway, disease, and gene set enrichment analyses. Through machine learning analysis, the genes AHR, APLNR, ITGA10, and NR2F6 were identified as potential diagnostic markers for LC, which was further supported using the ROC curves analysis showing a high diagnostic discriminative capacity. In addition, the CIBERSORT method was used to predict the extent of immune cell infiltration in AAA and standard tissue samples. T cells and M2 macrophages were found to be potential feature genes, implying that these immune cells may participate in the AAA progression and become potential targets for immune intervention. In the analysis conducted in [20], HLA-DRB1 helps link sclerosis, arthritis, diabetes and COVID-19 to antigen binders needed for immunotherapy and vaccine creation. This work presents HLA-DR4Pred2, a model built on 12,676 binder samples and an equal number of non-binder samples, whereas the previous HLA-DR4Pred was built using only 576 binders and non-binders. Different approaches

in machine learning models were used, and an AUROC of 0.90 with composition features and 0.87 using binary profile features were attained. The performance was even further improved when the composition -based features were complemented with the BLAST search; the AUROC was 0.93, and on a dataset with 12,676 binders and 86,300 non -binders, the AUROC was 0.99. This tool outperformed existing prediction methods and is available as a standalone tool and web server, with an additional Python package on the Python Package Index, facilitating the prediction, design, and virtual scanning of HLA-DRB10 binding peptides. In the analysis conducted in [21], HLA -DRB1 was found to be linked to multiplex diseases, including sclerosis, arthritis, diabetes and recently, COVID -19, underscores the relevance of the identification of binding peptides for immunotherapies and vaccines. Strengthening for omitted smaller datasets in previous models, the authors present the novel HLA-DR4Pred2, trained with a large data set of 12,676 binders and the same number of non-binders. This new model is based on its previous HLA-DR4Pred, which has been trained using only 576 binder and non -binder data. Cohort 1's 80% of samples were split into 5, and the chosen machine learning algorithms gave promising results, with the best values reaching 0.90 AUROC when using features based on compositions from the samples and 0.87 when using binary profile features. Additional improvement incorporating the BLAST search with the composition -based model improved the AUROC to 0.93. In addition, based on a larger dataset containing 12 676 patients with binder status and 86300 controls, the model reached an AUROC of 0.99. This study also introduced a standalone tool, a web server for HLA-DR4Pred2, and a Python package for broader accessibility, facilitating efficient prediction, design, and virtual scanning of HLA-DRB1 binding peptides. In the work presented in the paper designated as [22], vector-borne diseases are described as one of the main threats to human life worldwide, with high demand for new medical technologies. This study centered on the bacterial attachment to host extracellular proteins (exoproteome) in high-throughput using bacterial selection for high -throughput identification of host -interaction targets (BASEHIT). Using this technique, it was possible to map interactions versus 3,324 human exoproteins with 82 pathogen samples: 30 arthropod-borne strains and 8 non -vector-borne related strains. This large -scale analysis discovered 1,303-candidate pathogen–host PPIs, including many essential factors for pathogenesis, such as host cell invasion, tissue tropism, immune system modulation, and sensing. Major conclusions outlined that Lyme disease spirochetes recognize the epidermal growth factor as a transcriptional signal, and other approximately universal host-pathogen interactions between intracellular pathogens and thioredoxins occur. This Interactome atlas is, therefore, a rich source of molecular information on microbial pathogenesis and brings to light potential host- directed therapeutic approaches for dealing with vector-borne diseases.

As described in [23], the molecular index is from the chemical structure. Based on the topological indices, it plays the index role in judging molecular properties such as boiling point, viscosity and melting point. In this paper, these indices of molecular symmetry can be used quantitatively in the QSAR to assist in creating drugs to fight blood cancer. Notably, this study employed molecular descriptors, which are

graph invariants, to investigate the chemical structures of the blood cancer drugs based on 8 degree-based path topological indices as the reducible first and second Zagreb indices and the reducible Randic index. With these molecular descriptors, the study could predict boiling point, molecular weight, LogP, and the degree of polarity of 18 blood cancer drugs. The performance of the predictive models was statistically significant, with $P < 0.05$ and $F > 2.5$, suggesting high accuracy of the experimental -computed value correlation. This, therefore, shows that it can provide accurate prediction and be useful on new drug scaffolds. An innovative smartphone-integrated POCT was designed in the research work reported in [24]. The POCT device was designed using the powers of large laboratory analytical instruments in a portable platform. This POCT system exploits a cascade reaction mediated by specific oxidases and a highly efficient molecular recognition element, the leaf -like zeolitic imidazolate framework, to achieve efficient biomarker profiling. The analysis revealed that parameters such as RGB and HSV of fluorescence images are well aligned with biomarker concentration. To assess these parameters and estimate the target levels, an artificial neural network (ANN) model was designed for a smartphone application, which provides regression values of 95% in both training and validation sets for four biomarkers. In the current discovery, the device successfully performed offline biomarker detection in serum in 50 minutes with fair performance against conventional laboratory techniques. This smartphone -integrated makes it a cost -efficient and effective means of rapidly monitoring metabolism biomarkers, giving a clear distinction between normal and abnormal samples. For example, as described in the paper [25], multi-dimensional NMR experimental data serve as the primary source of information for protein research and biomolecular NMR spectroscopy data analysis. However, in contrast to secondary databases like the Chemical Shift Indexes in the Biological Magnetic Resonance Data Bank (BMRB) or the protein structures in the Protein Data Bank (PDB), primary NMR data are only sometimes retained in the public domain. To fill this void, originally calculated solution NMR data for 100 proteins consisting of 1,329 two – to four-dimensional NMR spectra, chemical shift references, structures, peak lists, and coordinates for restraints calculations were collected and compiled. Collected initially to reinforce the spectra analysis method based on deep learning ARTINA, this dataset reproduces protein structures from initial experimental data. Thus, including this large dataset enhances the options for improving computational NMR spectroscopy, particularly machine learning methodologies, which allows for a proper comparison of methods.

Table eftab:literature-review indicates that numerous researchers have studied various environmental, host–pathogen, and anthropogenic factors that contribute to the spread of tick-borne infections. These studies show the need for an interdisciplinary approach combining ecology, epidemiology, and public health to address this increasing health threat.

3. CONCLUSION

Systematic examination of tick-borne diseases indicates dynamic environmental processes, interactions between hosts and vectors, and human activities drive their spread. The

Table 1. Summary of Literature Review

Study	Disease Focus	Methodology	Key Findings	Implications
[10]	Tick-borne diseases (<i>Ixodes scapularis</i>)	Geographic and demographic analysis in New York, USA	Nymphal tick abundance showed northward and westward spread; significant in new areas but stable in older areas	Forecasting models aid public-health planning and early prevention in tick-transmission areas
[11]	Lyme disease via black-legged ticks	ANN model using climate and host data	Heavy larval tick burdens were linked to temperature and precipitation	Environmental conditions are critical for control and ANN models can track vector patterns
[12]	Tick-borne diseases (<i>Ixodes ricinus</i>)	Supervised machine-learning models in southern Scandinavia	Temperature and vegetation highly influenced tick distribution and abundance	Climate-based models aid risk assessment and incidence prediction
[13]	Lyme disease diagnosis and care	Updated clinical guidance review	Emphasized evidence-based diagnosis, treatment and prophylaxis after tick exposure	Supports timely clinical decision-making and reduces inappropriate testing
[14]	Lyme disease and imaging support	Image segmentation and denoising approaches	Pre-processing improved diagnostic image quality and segmentation performance	Image-processing methods can improve diagnostic accuracy
[15]	Tick-borne disease control	Optimal control models	Timing and frequency of tick-reduction efforts strongly affected disease risk	Discontinuous control can reduce risk without continuous treatment
[16]	Human Lyme disease incidence	County-level acarological and epidemiological comparison	Infected nymph density correlated with reported human cases, with regional variation	More state-level data are needed for cost-effectiveness evaluation
[17]	Tick-borne disease risk in northeast China	Structured questionnaire and structural equation modeling	Tick bites were strongly associated with self-reported infection rates	Behavioral and cognitive findings guide prevention strategies
[18]	Asian longhorned tick	Microhabitat modeling	Environmental and local habitat variables explained invasive tick presence	Supports targeted surveillance and habitat-based control
[19]	Immune-related disease prediction	Machine-learning feature-gene analysis	Predictive biomarkers were correlated with immune infiltration	Suggests immune intervention as a potential treatment target
[21]	HLA-DRB1 peptide binding	HLA-DR4Pred2 machine-learning model	AUROC improved to 0.99	Supports peptide design and immunotherapy development
[22]	Vector-borne diseases	High-throughput host interaction analysis	Mapped 1,303 pathogen–host interactions, including Lyme-disease spirochetes	Advances host-targeted therapeutic approaches
[23]	Blood-cancer drug development	QSAR and molecular descriptors	Models predicted boiling point, molecular weight and polarity with high accuracy	Assists efficacy prediction for new drug candidates
[24]	Point-of-care testing device	Smartphone-integrated ANN analysis	Biomarker levels were determined accurately in 50 minutes	Offers portable and cost-effective biomarker monitoring
[25]	Protein-structure re-search	ARTINA deep-learning model for NMR data	Achieved mean RMSD of 1.2 Å in protein reproduction	Advances biomolecular NMR data interpretation

geographic expansion and numerical increase of such vector species, such as *Ixodes scapularis* and *Ixodes ricinus*, will expand in range and abundance due to climate change and changing landscapes that favor the growth of ticks. These factors, however, introduce new threats to public health as endemic diseases appear in areas with no previous histories of occurrence of such diseases, making better proactive measures in the management and control of the diseases paramount. Modern ecological and epidemiological models, thus, are fundamental, predictive risk assessment strategies in these situations, showing where the risk is likely high and how the resources should be used. Creative uses of machine learning and other predictive models that have improved our ability to accurately trace and predict the population of ticks, improving our understanding of the patterns of disease spread. These tools allow for the inclusion of extra environmental factors, demographic, and host factors, which serve as a basis for better public health system strategies, allowing for the implementation of targeted measures that avert transmission of diseases. In addition, new diagnostics technologies make it possible to promptly identify the presence of tick-borne pathogens, making it possible to respond clinically and epidemiologically to the latest outbreaks. Combining these approaches gives health authorities concrete information that allows them to act more quickly and effectively in the fight against tick-borne diseases. A layered approach is paramount to address the growing challenge of tick-borne diseases adequately. Using ecological surveillance, health promotion, managing hosts, and developing diagnostic methods effectively reduces the burden of such diseases. Maintaining partnerships between various population strata, scientists, health authorities, and government representatives will be essential in modifying these approaches to environmental and soci-

etal changes over time. Given that the burden of tick-borne diseases is increasing worldwide, it is essential to take such measures and use an integrated approach to protect people's health and consider the risks of outbreaks.

REFERENCES

- [1] J. S. Gray, H. Dautel, A. Estrada-Peña, O. Kahl, and E. Lindgren, "Effects of climate change on ticks and tick-borne diseases in Europe," *Interdisciplinary Perspectives on Infectious Diseases*, 2009.
- [2] H. V. Pham et al., "Climatic and environmental drivers of tick distribution," 2021.
- [3] A. Beck et al., "Knowledge, attitudes, and behaviors regarding tick-borne disease prevention," *Ticks and Tick-borne Diseases*, vol. 13, no. 3, 2022.
- [4] K. Garcia, M. Weakley, T. Do, and S. Mir, "Current and future molecular diagnostics of tick-borne diseases in cattle," *Veterinary Sciences*, vol. 9, no. 5, p. 241, 2022.
- [5] C. Luo, P. Pearson, G. Xu, and S. M. Rich, "A computer vision-based approach for tick identification using deep learning models," *Insects*, vol. 13, no. 2, p. 116, 2022.
- [6] S. Mowry, F. Keesing, I. R. Fischhoff, and R. S. Ostfeld, "Predicting larval tick burden on white-footed mice with an artificial neural network," *Ecological Informatics*, vol. 52, pp. 150–158, 2019.
- [7] T. Tran et al., "Spatio-temporal variation in environmental features predicts the distribution and abundance of *Ixodes scapularis*," *International Journal for Parasitology*, vol. 51, no. 4, pp. 311–320, 2021.

-
- [8] L. Jung Kjær et al., “Predicting the spatial abundance of Ixodes ricinus ticks in southern Scandinavia using environmental and climatic data,” *Scientific Reports*, vol. 9, no. 1, 2019.
 - [9] P. G. Auwaerter, T. Kobayashi, and G. P. Wormser, “Guidelines for Lyme disease are updated,” *American Journal of Medicine*, vol. 134, no. 11, pp. 1314–1316, 2021.
 - [10] T. Arul Raj and D. A. Parvathavarthine, “Performance analysis of clustering based image segmentation techniques for detecting Lyme disease,” 2022.
 - [11] H. Gaff, E. Schaefer, and S. Lenhart, “Use of optimal control models to predict treatment time for managing tick-borne disease,” *Journal of Biological Dynamics*, vol. 5, no. 5, pp. 517–530, 2011.
 - [12] K. M. Pepin et al., “Geographic variation in the relationship between human Lyme disease incidence and density of infected host-seeking Ixodes scapularis nymphs,” *American Journal of Tropical Medicine and Hygiene*, vol. 86, no. 6, p. 1062, 2012.
 - [13] R. Fang et al., “Behavioral and cognitive factors influencing tick-borne disease risk in northeast China,” *One Health*, vol. 18, p. 100736, 2024.
 - [14] I. Rochlin et al., “Microhabitat modeling of the invasive Asian longhorned tick in New Jersey, USA,” *Ticks and Tick-borne Diseases*, vol. 14, no. 2, p. 102126, 2023.
 - [15] Y. Zhang and G. Li, “Predicting feature genes correlated with immune infiltration based on machine learning algorithms,” *Scientific Reports*, vol. 14, no. 1, 2024.
 - [16] S. Patiyal, A. Dhall, N. Kumar, and G. P. S. Raghava, “HLA-DR4Pred2: An improved method for predicting HLA-DRB1 binders,” *Methods*, vol. 232, pp. 18–28, 2024.
 - [17] T. M. Hart et al., “An atlas of human vector-borne microbe interactions reveals pathogenicity mechanisms,” *Cell*, vol. 187, no. 15, pp. 4113–4127.e13, 2024.
 - [18] A. Mahboob, M. W. Rasheed, L. Amin, and I. Hanif, “A study of novel molecular descriptors and quantitative structure-property relationship analysis of blood cancer drugs,” *European Physical Journal Plus*, vol. 138, no. 9, p. 856, 2023.
 - [19] H. Chen et al., “Artificial neural network processed linear-light tristimulus and hue parameters for smart-phone assisted point-of-care testing,” *Sensors and Actuators B: Chemical*, vol. 384, p. 133659, 2023.
 - [20] P. Klukowski et al., “The 100-protein NMR spectra dataset,” *Scientific Data*, vol. 11, no. 1, 2024.
 - [21] M. Pham et al., “Validation of heat-inducible Ixodes scapularis HSP70 and tick-specific 3xP3 promoters in ISE6 cells,” *iScience*, vol. 27, no. 8, p. 110468, 2024.