



## On Imperfect Duplets In Some Refined Neutrosophic Rings

Katy D. Ahmad, Islamic University Of Gaza, Palestine

Mikail Bal, Gaziantep University, Turkey

Arwa A. Hajjari, Cairo University, Egypt.

Rozina Ali, Cairo University, Egypt

Co-: [Katyon765@gmail.com](mailto:Katyon765@gmail.com)

[Rozyyy123n@gmail.com](mailto:Rozyyy123n@gmail.com)

**Abstract:** This paper solves the imperfect duplets problem in refined neutrosophic rings, where it presents the necessary and sufficient conditions for a pair  $(x, y)$  to be an imperfect duplet in any refined neutrosophic ring. Also, this work introduces a full description of the structure of imperfect duplets in numerical refined neutrosophic rings such as refined neutrosophic ring of integers  $Z(I_1, I_2)$ , refined neutrosophic ring of rationales  $Q(I_1, I_2)$ , and refined neutrosophic ring of real numbers  $R(I_1, I_2)$ .

**Keywords:** Refined Neutrosophic Ring, imperfect Duplet, Imperfect triplet

### 1. Introduction

Neutrosophy is a generalization of intuitionistic fuzzy logic founded by F.Smarandache to deal with indeterminacy in science and real life problems.

Neutrosophic algebra began with the efforts of Kandasamy and Smarandache where the concept of neutrosophic ring was presented in [7] as a generalization of classical rings. These rings were handled by many authors such as [1,2,9,11-19].

Recently, there is an increasing interest in the generalizations of neutrosophic rings, where refined neutrosophic rings were defined by Agboola et.al [5,6].

If  $(R, +, \cdot)$  is a ring, then the corresponding refined neutrosophic ring  $R(I_1, I_2)$  is defined as follows:

$R(I_1, I_2) = \{(a, bI_1, cI_2); a, b, c \in R\}$ . The operations on  $R(I_1, I_2)$  are defined as follows:

Addition:  $(a, bI_1, cI_2) + (x, yI_1, zI_2) = (a + x, (b + y)I_1, (c + z)I_2)$ .

Multiplication:  $(a, bI_1, cI_2) \cdot (x, yI_1, zI_2) = (ax, (ay + bx + by + bz + cy)I_1, (az + cz + cx)I_2)$

The notion of neutrosophic duplets and neutrosophic triplets was defined and handled by Smarandache et.al in [3,8,10], where they opened an interesting research direction about finding these elements in rings.

Through this paper, we extend the previous efforts to solve the problem of duplets into the case of refined neutrosophic rings, where we present the condition of imperfect duplets in any refined neutrosophic ring even when it is not commutative. In particular, we determine all possible imperfect duplets in the refined neutrosophic ring of integers  $Z(I_1, I_2)$ , refined neutrosophic ring of rationales  $Q(I_1, I_2)$ , and refined neutrosophic ring of real numbers  $R(I_1, I_2)$ .

## 2. Preliminaries

### Definition 2.1: [5]

(a) If  $X$  is a set then  $X(I_1, I_2) = \{(a, bI_1, cI_2): a, b, c \in X\}$  is called the refined neutrosophic set generated by  $X, I_1, I_2$ .

(b) Let  $(R, +, \times)$  be a ring,  $(R(I_1, I_2), +, \times)$  is called the refined neutrosophic ring generated by  $R, I_1, I_2$ .

### Definition 2.2: [12]

Let  $R$  be any ring,  $x, y$  are two arbitrary elements in  $R$ . We call them a duplet with  $y$  acts as an identity if and only if

$$xy = yx = x.$$

### Definition 2.3: [10]

Let  $R$  be any ring,  $x, y, z$  three arbitrary elements in  $R$ . We call them a triplet with  $y$  acts as an identity if and only if

$$xy = yx = x, zy = yz = z, xz = zx = y.$$

## 3. Main discussion :

### Definition 3.1:

Let  $x, y, z$  be three elements in a refined neutrosophic ring, we have:

$x, y$  are called an imperfect refined neutrosophic duplet with  $y$  acts as an identity if and only if  $xy = yx = x$ .

**Example 3.2:**

In  $Z_{10}(I_1, I_2)$  the refined neutrosophic ring of integers modulo 10, we have:

$x = (2,0,0), y = (6,0,0)$  is an imperfect refined neutrosophic duplet with  $y$  acts as an identity, that is because  $xy = yx = x$ .

**Theorem 3.3:**

Let  $x = (x_0, x_1I_1, x_2I_2), y = (y_0, y_1I_1, y_2I_2)$  be any two elements in  $R(I_1, I_2)$ , then  $(x, y)$  is an imperfect refined neutrosophic duplet with  $y$  acts as an identity if and only if

$(x_0, y_0), (x_0 + x_2, y_0 + y_2), (x_0 + x_1 + x_2, y_0 + y_1 + y_2)$  are three imperfect duplets in the classical ring  $R$  with  $y_0, y_0 + y_2, y_0 + y_1 + y_2$  acts as identities.

Proof:

We assume that  $(x, y)$  is an imperfect refined neutrosophic duplet with  $y$  acts as an identity, then  $xy = yx = x$ . From this condition we get:

$$\begin{aligned} xy &= (x_0y_0, I_1[x_0y_1 + x_1y_0 + x_1y_1 + x_1y_2 + x_2y_1], I_2[x_0y_2 + x_2y_0 + x_2y_2]) = (x_0y_0, I_1[(x_0 + x_1 + x_2)(y_0 + y_1 + y_2) - (x_0 + x_2)(y_0 + y_2)], I_2[(x_0 + x_2)(y_0 + y_2) - x_0y_0]) = x, \text{ and } yx = \\ &= (y_0x_0, I_1[y_1x_0 + y_0x_1 + y_1x_1 + y_1x_2 + y_2x_1], I_2[y_2x_0 + y_0x_2 + y_2x_2]) = (y_0x_0, I_1[(y_0 + y_1 + y_2)(x_0 + x_1 + x_2) - (y_0 + y_2)(x_0 + x_2)], I_2[(y_0 + y_2)(x_0 + x_2) - y_0x_0]) = x. \end{aligned}$$

This implies that:  $x_0y_0 = x_0 = y_0x_0, (x_0 + x_2)(y_0 + y_2) - x_0y_0 = (y_0 + y_2)(x_0 + x_2) - y_0x_0 = x_2, (x_0 + x_1 + x_2)(y_0 + y_1 + y_2) - (x_0 + x_2)(y_0 + y_2) = (y_0 + y_1 + y_2)(x_0 + x_1 + x_2) - (y_0 + y_2)(x_0 + x_2) = x_1$

Thus  $(x_0 + x_2)(y_0 + y_2) = (y_0 + y_2)(x_0 + x_2) = x_0 + x_2, \text{ and } (x_0 + x_1 + x_2)(y_0 + y_1 + y_2) = (y_0 + y_1 + y_2)(x_0 + x_1 + x_2) = x_0 + x_1 + x_2$ . Hence  $(x_0, y_0), (x_0 + x_2, y_0 + y_2), (x_0 + x_1 + x_2, y_0 + y_1 + y_2)$  are three imperfect duplets in the classical ring  $R$  with  $y_0, y_0 + y_2, y_0 + y_1 + y_2$  acts as identities.

The converse can be proved by the same.

Now, we find the set of all imperfect duplets in some refined neutrosophic rings.

**Theorem 3.4:**

Let  $Q(I_1, I_2)$  be the refined neutrosophic ring of rationales, the all non-trivial imperfect refined neutrosophic duplets have exactly the following 6 forms

$$1-)\{(0, x_1I_1, 0), (y_0, y_1I_1, y_2I_2); y_0 + y_1 + y_2 = 1, x_1 \neq 0\},$$

$$2-)\{(0, x_1I_1, -x_1I_1), (y_0, y_1I_1, y_2I_2); y_0 + y_2 = 1 \text{ and } x_1 \neq 0\},$$

$$3-)\{(0, x_1I_1, x_2I_1), (y_0, 0, y_2I_2); y_0 + y_2 = 1\},$$

$$4-)\{(x_0, 0, -x_0I_2), (1, y_1I_1, y_2I_2)\},$$

$$5-)\{(x_0, x_1I_1, -x_0I_2), (1, y_1I_1, -y_1I_2); x_0 \neq 0\},$$

$$6-)\{(x_0, x_1I_1, x_2I_2), (1, y_1I_1, 0); x_0 + x_1 + x_2 = 0\}.$$

Proof:

According to Theorem 3.3, we have

$$x_0y_0 = x_0 \text{ (equation I)}, (x_0 + x_2)(y_0 + y_2) = x_0 + x_2 \text{ (equation II)}, (x_0 + x_1 + x_2)(y_0 + y_1 + y_2) = x_0 + x_1 + x_2 \text{ (equation III)}.$$

From equation I, we get  $x_0 = 0$  or  $y_0 = 1$ . Firstly, we assume that  $x_0 = 0$ , from (II) we get

$x_2 = 0$  or  $y_0 + y_2 = 1$ , from equation (III), we get  $x_1 + x_2 = 0$  or  $y_0 + y_1 + y_2 = 1$ . Now we discuss the following possible cases:

If ( $x_2 = 0$ , with  $x_1 + x_2 = 0$ ) we get a trivial duplet since  $x = (0,0,0)$ .

If ( $x_2 = 0$ , with  $y_0 + y_1 + y_2 = 1$ ) we get the following duplet  $x = (0, x_1I_1, 0)$ ,  $y = (y_0, y_1I_1, y_2I_2)$  with  $y_0 + y_1 + y_2 = 1$  and  $x_1 \neq 0$ .

If ( $x_2 \neq 0$  and  $y_0 + y_2 = 1$ , with  $x_1 + x_2 = 0$ ) we get the following duplet

$$x = (0, x_1I_1, -x_1I_2), y = (y_0, y_1I_1, y_2I_2) \text{ with } y_0 + y_2 = 1.$$

If ( $x_2 \neq 0$  and  $x_1 + x_2 \neq 0$  with  $y_0 + y_2 = 1$ , with  $y_0 + y_1 + y_2 = 1$ ), then  $y_1=0$ , we get the following duplet

$$x = (0, x_1I_1, x_2I_2), y = (y_0, 0, y_2I_2); y_0 + y_2 = 1 .$$

Now, we discuss the second case of equation (I), we suppose that  $y_0 = 1$  and  $x_0 \neq 0$ .

From equation (II) we get  $x_0 + x_2 = 0$  or  $y_2 = 0$ . From equation (III) we get  $x_0 + x_1 + x_2 = 0$  or  $y_1 + y_2 = 0$ . We discuss the possible cases

If ( $x_0 + x_2 = 0$  and  $y_2 \neq 0$  with  $x_0 + x_1 + x_2 = 0$ ) we get  $x_1 = 0$  and  $x_2 = -x_0$ , thus the corresponding duplet is

$$x = (x_0, 0, -x_0I_2), y = (1, y_1I_1, y_2I_2).$$

If ( $x_0 + x_2 = 0$  and  $y_2 \neq 0$  with  $y_1 + y_2 = 0$ ) we get  $y_1 = -y_2, x_2 = -x_0$ , thus the corresponding duplet is  $x = (x_0, x_1I_1, -x_0I_2), y = (1, y_1I_1 - y_1I_2)$ .

If ( $x_0 + x_2 \neq 0$  and  $y_2 = 0$  with  $y_1 + y_2 = 0$ ) we get a trivial duplet since  $y = (1, 0, 0)$ .

If ( $x_0 + x_2 \neq 0$  and  $y_2 = 0$  with  $x_0 + x_1 + x_2 = 0$ ) we get the following duplet

$$x = (x_0, x_1I_1, x_2I_2), y = (1, y_1I_1, 0). \text{ Thus the proof is complete.}$$

### Example 3.5:

In the first form of duplets we put  $x_1 = 2, y_0 = y_1 = y_2 = \frac{1}{3}$ . It is clear that  $xy = yx = x$ .

### Theorem 3.6:

Let  $R(I_1, I_2)$  be the refined neutrosophic ring of real numbers, the all non-trivial imperfect refined neutrosophic duplets have exactly the following 6 forms

$$1-)\{(0, x_1I_1, 0), (y_0, y_1I_1, y_2I_2); y_0 + y_1 + y_2 = 1, x_1 \neq 0\},$$

$$2-)\{(0, x_1I_1, -x_1I_1), (y_0, y_1I_1, y_2I_2); y_0 + y_2 = 1 \text{ and } x_1 \neq 0\},$$

$$3-)\{(0, x_1I_1, x_2I_1), (y_0, 0, y_2I_2); y_0 + y_2 = 1\},$$

$$4-)\{(x_0, 0, -x_0I_2), (1, y_1I_1, y_2I_2)\},$$

$$5-)\{(x_0, x_1I_1, -x_0I_2), (1, y_1I_1, -y_1I_2); x_0 \neq 0\},$$

$$6-)\{(x_0, x_1I_1, x_2I_2), (1, y_1I_1, 0); x_0 + x_1 + x_2 = 0\}.$$

Proof:

The proof is exactly similar to Theorem 3.4 .

**Theorem 3.7:**

Let  $Z(I_1, I_2)$  be the refined neutrosophic ring of integers, the all non-trivial imperfect refined neutrosophic duplets have exactly the following 6 forms

$$1-) \{(0, x_1 I_1, 0), (y_0, y_1 I_1, y_2 I_2); y_0 + y_1 + y_2 = 1, x_1 \neq 0\},$$

$$2-) \{(0, x_1 I_1, -x_1 I_1), (y_0, y_1 I_1, y_2 I_2); y_0 + y_2 = 1 \text{ and } x_1 \neq 0\},$$

$$3-) \{(0, x_1 I_1, x_2 I_1), (y_0, 0, y_2 I_2); y_0 + y_2 = 1\},$$

$$4-) \{(x_0, 0, -x_0 I_2), (1, y_1 I_1, y_2 I_2); x_0 \neq 0\},$$

$$5-) \{(x_0, x_1 I_1, -x_0 I_2), (1, y_1 I_1, -y_1 I_2); x_0 \neq 0\},$$

$$6-) \{(x_0, x_1 I_1, x_2 I_2), (1, y_1 I_1, 0); x_0 + x_1 + x_2 = 0\}.$$

Proof:

The proof is exactly similar to Theorem 3.4.

**Example 3.8:**

We construct an example about an imperfect duplet in  $Q(I_1, I_2)$  .

Consider the corresponding imperfect duplet  $x = (0, \frac{1}{3} I_1, -\frac{1}{3} I_2), y = (0, -I_1, I_2)$ .

**Example 3.9:**

We construct an example about an imperfect duplet in  $Z(I_1, I_2)$ .

$$x = (1, I_1, -2I_2), y = (1, I_1, 0).$$

**Conclusion**

In this paper, we have studied the problem of finding imperfect duplets in a refined neutrosophic ring.

Where we have determined a general condition for imperfect duplets in such rings. In particular, we have

presented the structure of all duplets in the refined neutrosophic rings of integers, reals, and rationales respectively.

### **References**

- [1] T.Chalapathi and L. Madhavi,. "Neutrosophic Boolean Rings", Neutrosophic Sets and Systems, Vol. 33, pp. 57-66, 2020.
- [2] S. A. Edalatpanah., " Systems of Neutrosophic Linear Equations", Neutrosophic Sets and Systems, Vol. 33, pp. 92-104, 2020.
- [3] Kandasamy, V., Kandasamy, I., and Smarandache, F., " Algebraic Structure of Neutrosophic Duplets In Neutrosophic Rings  $\langle Z \cup I \rangle$ ,  $\langle Q \cup I \rangle$ , and  $\langle R \cup I \rangle$  ", Neutrosophic Sets and Systems, Vol.23, pp.85-95, 2018.
- [4] Kandasamy, V. W. B., Ilanthenral, K., and Smarandache, F., " Semi-Idempotents in Neutrosophic Rings", Mathematics Journal (MDPI), Vol. 7, 2019.
- [5] Adeleke, E.O., Agboola, A.A.A.,and Smarandache, F., "Refined Neutrosophic Rings I", International Journal of Neutrosophic Science, Vol. 2(2), pp. 77-81, 2020.
- [6] Adeleke, E.O., Agboola, A.A.A., and Smarandache, F., "Refined Neutrosophic Rings II", International Journal of Neutrosophic Science, Vol. 2(2), pp. 89-94, 2020.
- [7] Kandasamy, V.W.B., and Smarandache, F., "Some Neutrosophic Algebraic Structures and Neutrosophic N-Algebraic Structures", Hexis, Phonex, Arizona 2006.
- [8] Kandasamy, V., Kandasamy, I., and Smarandache, F., "Neutrosophic Duplets of  $\{Z_p^n, \times\}$  and  $\{Z_{pq}, \times\}$  and Their Properties", Symmetry, 2018.
- [9] Yingcang, Ma., Xiaohong Zhang ., Smarandache, F., and Juanjuan, Z., The Structure of Idempotents in Neutrosophic Rings and Neutrosophic Quadruple Rings, Symmetry, Vol. 11, 2019.
- [10] Kandasamy, V., Kandasamy, I., and Smarandache, F., "Neutrosophic Triplets In Neutrosophic Rings", Mathematics, 2019.

[11] M. Ali, F. Smarandache, M. Shabir and L. Vladareanu, "Generalization of Neutrosophic Rings and Neutrosophic Fields," *Neutrosophic Sets and Systems*, vol. 5, pp. 9-14, 2014.

[12] Florentin Smarandache, *Neutrosophic Duplet Structures*, Joint Fall 2017 Meeting of the Texas Section of the APS, Texas Section of the AAPT, and Zone 13 of the Society of Physics Students, The University of Texas at Dallas, Richardson, TX, USA, October 20-21, 2017, <http://meetings.aps.org/Meeting/TSF17/Session/F1.32>

[13] Abobala, M., "On The Characterization of Maximal and Minimal Ideals In Several Neutrosophic Rings", *Neutrosophic Sets and Systems*, Vol. 45, 2021.

[14] Abobala, M., "Neutrosophic Real Inner Product Spaces", *Neutrosophic Sets and Systems*, Vol. 43, 2021.

[15] Abobala, M., On Refined Neutrosophic Matrices and Their Applications In Refined Neutrosophic Algebraic Equations, *Journal Of Mathematics*, Hindawi, 2021

[16] Abobala, M., On The Representation of Neutrosophic Matrices by Neutrosophic Linear Transformations, *Journal of Mathematics*, Hindawi, 2021.

[17] Abobala, M., "On Some Algebraic Properties of n-Refined Neutrosophic Elements and n-Refined Neutrosophic Linear Equations", *Mathematical Problems in Engineering*, Hindawi, 2021

[18] Abobala, M., Partial Foundation of Neutrosophic Number Theory, *Neutrosophic Sets and Systems*, Vol. 39 , 2021.

[19] Abobala, M., "A Study Of Nil Ideals and Kothe's Conjecture In Neutrosophic Rings", *International Journal of Mathematics and Mathematical Sciences*, hindawi, 2021

[20] Abobala, M., and Hatip, A., "An Algebraic Approach To Neutrosophic Euclidean Geometry", *Neutrosophic Sets and Systems*, Vol. 43, 2021.

[21] Abobala, M., Bal, M., and Hatip, A., " A Review On Recent Advantages In Algebraic Theory Of Neutrosophic Matrices", *International Journal of Neutrosophic Science*, Vol. 17, 2021.

[22] Hajjari, A., and Ali, R., " A Contribution To Kothe's Conjecture In Refined Neutrosophic Rings", *International Journal of Neutrosophic Science*", Vol. 16, 2021.

- [23] Olgun, N., Hatip, A., Bal, M., and Abobala, M., " A Novel Approach To Necessary and Sufficient Conditions For The Diagonalization of Refined Neutrosophic Matrices", *International Journal of Neutrosophic Science*, Vol. 16, pp. 72-79, 2021.
- [24] Smith, J. L., and Gallian, J. A., "Factoring Finite Factor Rings", *Mathematics Magazine*, vol. 58, 1985.
- [25] J. A. Gallian, *Contemporary Abstract Algebra*, 7th ed., Cengage Learning, 2012.
- [26] Ibrahim, M., and Abobala, M., "An Introduction To Refined Neutrosophic Number Theory", *Neutrosophic Sets and Systems*, Vol. 45, 2021.
- [27] Chalapathi, T., and Kumar, K., "Neutrosophic Units of Neutrosophic Rings and Fields", *Neutrosophic Sets and Systems*, Vol. 21, 2018.
- [28] Abobala, M., "On Some Special Elements In Neutrosophic Rings and Refined Neutrosophic Rings", *Journal of New Theory*, vol. 33, 2020.
- [29] Singh, P.K., " Data With Turiyam Set for Fourth Dimension Quantum Information Processing", *Journal of Neutrosophic and Fuzzy Systems*, vol.1, 2022.
- [30] Singh, P.K., " Anti-geometry and NeutroGeometry Characterization of Non-Euclidean Data", *Journal of Neutrosophic and Fuzzy Systems*, vol. 1, 2022.
- [31] Smarandache, F., and Broumi, M., "Neutro-Intelligent Set is a particular case of the refined neutrosophic set", *Journal of Neutrosophic and Fuzzy Systems*, Vol. 1, 2022.
- [32] Sankari, H, and Abobala, M, " A Contribution to m-Power Closed Groups", *UMM-Alqura University Journal for Applied Sciences*, KSA, 2020.
- [33] Ahmad, K., Bal, M., and Aswad, M., " The kernel of Fuzzy and Anti Fuzzy Groups", *Journal of Neutrosophic and Fuzzy Systems*, Vol.1, 2022.
- [34] Ahmad, K., Bal, M., and Aswad, M., " A Short Note on The Solution Of Fermat's Diophantine Equation In Some Neutrosophic Rings", *Journal of Neutrosophic and Fuzzy Systems*, Vol. 1, 2022.
- [35] Abobala, M., "On Some Neutrosophic Algebraic Equations", *Journal of New Theory*, Vol. 33, 2020.
- [36] Abobala, M., A Study of Maximal and Minimal Ideals of n-Refined Neutrosophic Rings, *Journal of Fuzzy Extension and Applications*, Vol. 2, pp. 16-22, 2021.