

# Fixed point results in $\omega_t$ -distance mappings for Geraghty type contractions

Ammar Al-tawil<sup>1,\*</sup>, Ayman. A Hazaymeh<sup>1</sup>, Anwar Bataihah<sup>1</sup>

<sup>1</sup>Department of Mathematics, Faculty of Science, Jadara University, Irbid, Jordan Emails: ammartaweel82@gmail.com; aymanha@jadara.edu.jo; a.bataihah@jadara.edu.jo

### **Abstract**

In this study, we establish fixed point theorems for  $P\omega_t$ -contractions within b-metric spaces by utilizing  $\omega_t$ -distance mappings. Subsequently, we demonstrate fixed point results pertaining to nonlinear contraction conditions of the Geraghty type, again employing  $\omega_t$ -distance mappings in the context of a complete b-metric space. Additionally, we bolster our findings with appropriate examples to illustrate the applicability of our results.

**Keywords:** Fixed point; b-metric space;  $P\omega_t$ -contraction; Nonlinear contraction

## 1 Introduction and Preliminary

In mathematics, a fixed-point theorem is a result saying that a self-map function f on a non-empty set  $\mho$  will have at least one fixed point in  $\mho$  ( i.e., there is some  $\xi \in \mho$  such that  $f\xi = \xi$  ). This theorem holds true under certain conditions on f, which can be expressed in general terms. In a variety of mathematical contexts, the presence of a solution is often synonymous with the existence of a fixed point for an appropriate mapping. Consequently, the identification of fixed points carries significant ramifications across multiple domains within mathematics and other scientific disciplines. This theory represents a profound integration of analysis (both pure and applied) topology, and geometry. In the past five decades, the fixed point theory has emerged as a highly influential and essential instrument in the exploration of nonlinear analysis.

The Banach fixed-point theorem, 1 known also as the Banach contraction principle, plays a crucial role in the field of mathematics, particularly in the study of metric spaces. This theorem guarantees the presence and uniqueness of fixed points for certain self-maps in metric spaces. Moreover, it presents a methodical approach to determining these fixed points. Essentially, the Banach fixed-point theorem can be seen as a generalized version of Picard's method of successive approximations. It is named after Stefan Banach (1892–1945), who first introduced this theorem in 1922.

Numerous mathematicians have subsequently explored a range of generalizations of Banach's theorem across different contexts see for example. A notable example is the concept of b-metric spaces, which was first introduced by Bakhtin ad later refined and named by Czerwik. This framework has been employed to examine various fixed point theorems. Additionally, Hussain et al introduced the idea of  $\omega$ t-distance mappings, utilizing this concept to investigate several fixed point results.

**Definition 1.1.** <sup>3</sup> Let  $\mho$  denote a set. Define a function  $d: \mho \times \mho \to [0,\infty)$  that fulfills the following conditions:

1.  $d(\xi, \zeta) = 0$  if and only if  $\xi$  is equal to  $\zeta$ ;

- 2.  $d(\xi,\zeta) = d(\zeta,\xi)$  for all  $\xi,\zeta \in \mho$ ;
- 3. There exists a constant  $s \ge 1$  such that for any points  $\xi, \zeta, \rho \in \mathcal{V}$ , the inequality  $d(\xi, \zeta) \le s(d(\xi, \rho) + d(\rho, \zeta))$  holds.

The triplet  $(\mho, d, s)$  is referred to as a b-metric space.

It should be noted that in the case where s equals 1, the triplet  $(\mho, d, s)$  forms a metric space. This implies that the properties of a metric space hold true when s is equal to 1.

## **Definition 1.2.** <sup>3</sup>

Let  $(\mho, d, s)$  represent a b-metric space.

- 1. A sequence  $(\xi_n)$  is said to converge to an element  $\xi \in \mho$  if and only if the limit  $\lim_{n \to \infty} d(\xi_n, \xi) = 0$  holds true.
- 2. The sequence  $(\xi_n)$  is classified as Cauchy if and only if the limit  $\lim_{n,m\to\infty}d(\xi_n,\xi_m)=0$  is satisfied.
- 3. The space (0, d, s) is defined as complete if and only if every Cauchy sequence within 0 converges.

Kada et al.<sup>4</sup> presented the notion of  $\omega$ -distance within a metric space in 1996 and established several fixed point theorems. In the current study, we define the concept of  $\omega_t$ -distance and articulate a lemma that will be utilized in the principal sections of this research.

**Definition 1.3.** <sup>3</sup> Let  $(\mho, d, s)$  represent a b-metric space where the constant s satisfies  $s \ge 1$ . A function  $p : \mho \times \mho \to [0, \infty)$  is designated as a  $\omega_t$  on  $\mho$  if it fulfills the following conditions:

- (a) For any points  $\xi, \zeta, \rho \in \mathcal{V}$ , the inequality  $p(\xi, \rho) \leq s(p(\xi, \zeta) + p(\zeta, \rho))$  holds;
- (b) For every  $\xi \in \mho$ , the mapping  $p(\xi,\cdot):\mho \to [0,\infty)$  is s-lower semi-continuous;
- (c) For any given  $\varepsilon > 0$ , there exists a  $\delta > 0$  such that if  $p(\rho, \xi) \le \delta$  and  $p(\rho, \zeta) \le \delta$ , then it follows that  $d(\xi, \zeta) \le \varepsilon$ .

A real-valued function f defined on a b-metric space  $\mho$  is considered to be s-lower semi-continuous at a point  $\xi_0$  in  $\mho$  if one of the following conditions holds: either  $\liminf_{\xi_n \to \xi_0} f(\xi_n) = \infty$  or  $f(\xi_0) \le \liminf_{\xi_n \to \xi_0} sf(\xi_n)$ . This is applicable for sequences  $\xi_n$  that belong to  $\mho$  for every  $n \in \mathbb{N}$  and converge to  $\xi_0$ .

Next, we recall some examples on  $\omega_t$  mappings.

**Example 1.4.** <sup>3</sup> Let  $(\mho, d, s)$  represent a b-metric space. In this context, the function p = d serves as an  $\omega_t$ -distance on the set  $\mho$ .

*Proof.* The statements (a) and (b) are self-evident. To demonstrate (c), let us consider any  $\epsilon > 0$  and set  $\delta = \frac{\epsilon}{2s}$ . It follows that if  $p(\xi, \rho) \le \delta$  and  $p(\rho, \zeta) \le \delta$ , then it can be concluded that  $d(\xi, \zeta) \le \epsilon$ .

**Example 1.5.**  $^3$  Let  $\mho = \mathbb{R}$  and define the distance function as  $d(\xi, \zeta) = (\xi - \zeta)^2$ . The function  $p : \mho \times \mho \to [0, \infty)$ , given by  $p(\xi, \zeta) = |\xi|^2 + |\zeta|^2$  for all  $\xi, \zeta \in \mho$ , serves as an  $\omega_t$ -distance on the set  $\mho$ .

*Proof.* The assertions (a) and (b) are self-evident. To demonstrate (c), let us consider any  $\epsilon > 0$  and set  $\delta = \frac{\epsilon}{4}$ . Consequently, we obtain the following inequality:

$$d(\xi,\zeta) = (\xi - \zeta)^2 \le 2|\xi|^2 + 2|\zeta|^2 = 2p(\rho,\xi) + 2p(\rho,\zeta) = 2\delta + 2\delta = \epsilon.$$

**Example 1.6.** <sup>3</sup> Let  $\mho = \mathbb{R}$  and define the distance function  $d(\xi, \zeta) = (\xi - \zeta)^2$ . The function  $p : \mho \times \mho \to [0, \infty)$ , given by  $p(\xi, \zeta) = |\zeta|^2$  for all  $\xi, \zeta \in \mho$ , qualifies as a  $\omega_t$ -distance on  $\mho$ .

*Proof.* The assertions (a) and (b) are self-evident. To demonstrate (c), let us consider any  $\epsilon > 0$  and set  $\delta = \frac{\epsilon}{4}$ . Consequently, we obtain the following inequality:

$$d(\xi,\zeta) = (\xi - \zeta)^2 \le 2|\xi|^2 + 2|\zeta|^2 = 2p(\rho,\xi) + 2p(\rho,\zeta) = 2\delta + 2\delta = \epsilon.$$

**Lemma 1.7.**  $^3$  Let  $(\mho,d,s)$  represent a b-metric space characterized by a constant  $s \geq 1$ , and let p denote a  $\omega_t$ -distance defined on  $\mho$ . Consider sequences  $(\xi_n)$  and  $(\zeta_n)$  within  $\mho$ , along with sequences  $(\alpha_n)$  and  $(\beta_n)$  in the interval  $[0,\infty)$  that converge to zero, and let  $\xi,\zeta,\rho$  be elements of  $\mho$ . The following statements are established:

- 1. If for every natural number n, the conditions  $p(\xi_n, \zeta) \leq \alpha_n$  and  $p(\xi_n, \rho) \leq \beta_n$  hold, then it follows that  $\zeta = \rho$ . Specifically, if  $p(\xi, \zeta) = 0$  and  $p(\xi, \rho) = 0$ , it can be concluded that  $\zeta = \rho$ .
- 2. If the inequalities  $p(\xi_n, \zeta_n) \leq \alpha_n$  and  $p(\xi_n, \rho) \leq \beta_n$  are satisfied for all natural numbers n, then the distance  $d(\zeta_n, \rho)$  approaches zero.
- 3. If the condition  $p(\xi_n, \xi_m) \leq \alpha_n$  is satisfied for all natural numbers n and m with m > n, then the sequence  $(\xi_n)$  is identified as a Cauchy sequence.
- 4. If the inequality  $p(\zeta, \xi_n) \leq \alpha_n$  holds for every natural number n, then the sequence  $(\xi_n)$  is also classified as a Cauchy sequence.

*Proof.* (1) Let  $\epsilon > 0$  be given. Since  $\alpha_n \to 0$ , then  $\forall \delta > 0 \exists N_0 \in \mathbb{N}$  such that  $\alpha_n \leq \delta, \forall n \geq N_0$ .

Similarly, since  $\beta_n \to 0$ , then  $\forall \delta > 0 \exists N_1 \in \mathbb{N}$  such that  $\beta_n \leq \delta, \forall n \geq N_1$ .

If  $N = \max\{N_0, N_1\}$ , then:

$$\alpha_n < \delta$$
, and  $\beta_n < \delta$  for  $n > N$ 

so,

$$p(\xi_n, \zeta) \le \alpha_n < \delta \text{ and } p(\xi_n, \rho) \le \beta_n < \delta \text{ for } n \ge N$$

Thus, we have

$$d(\zeta, \rho) < \epsilon$$

Hence,  $d(\zeta, \rho) = 0$  and so,  $\zeta = \rho$ .

(2) Let  $\epsilon > 0$  be given. Since  $\alpha_n \to 0$ , then  $\forall \delta > 0 \exists N_0 \in \mathbb{N}$  such that  $\alpha_n \leq \delta, \forall n \geq N_0$ .

Similarly, since  $\beta_n \to 0$ , then  $\forall \delta > 0 \exists N_1 \in \mathbb{N}$  such that  $\beta_n \leq \delta, \forall n \geq N_1$ .

If  $N = \max\{N_0, N_1\}$ , then:

$$\alpha_n < \delta$$
, and  $\beta_n < \delta$  for  $n > N$ 

so,

$$p(\xi_n, \zeta_n) \le \alpha_n < \delta \text{ and } p(\xi_n, \rho) \le \beta_n < \delta \text{ for } n \ge N$$

$$\Rightarrow d(\zeta_n, \rho) < \epsilon \quad \text{for each } \epsilon > 0$$

$$\Rightarrow d(\zeta_n, \rho) \to 0$$

(3) Let  $\epsilon > 0$  be given. Since  $\alpha_n \to 0$ , then  $\forall \delta > 0 \exists N_0 \in \mathbb{N}$  such that  $\alpha_n \leq \delta, \forall n \geq N_0$ .

Hence,  $p(\xi_n, \xi_m) \le \alpha_n < \delta$  and  $p(\xi_n, \xi_{n+1})\alpha_n < \delta$  for  $m \ge n \ge N_0$ . Thus, by the definition of p, we have

$$d(\xi_{n+1}, \xi_m) < \epsilon \quad \text{for } m \ge n \ge N.$$

This implies that  $(\xi_n)$  is a Cauchy sequence.

(4) Let  $\epsilon > 0$  be given. Since  $\alpha_n \to 0$ , then  $\forall \delta > 0 \exists N_0 \in \mathbb{N}$  such that  $\alpha_n \leq \delta, \forall n \geq N_0$ 

$$\alpha_n \le \delta$$
 for  $n \ge N$ .

So,

$$p(\zeta, \xi_n) \le \alpha_n < \delta \quad \text{for } n \ge N_0$$

So,

$$\alpha_m \leq \delta$$
 for  $m \geq N_0$ .

Then,  $p(\zeta, \xi_n) \le \alpha_m < \delta$  for  $m \ge N_1$ 

If  $N = \max\{N_0, N_1\}$ , then:

$$d(\zeta, \xi_n) < \epsilon \quad \text{for } n, m \ge N.$$

This implies that  $(\xi_n)$  is a Cauchy sequence.

# 2 Main Result

Next, we give the definition of  $P\omega_t$  contractions in  $\omega_t$  mappings.

**Definition 2.1.** Let  $d: \mho \times \mho \to [0,\infty)$  be a b-metric on  $\mho$ , p be a  $\omega_t$ -distance on  $\mho$  there exists  $k \in [0,1)$  such that  $f: \mho \to \mho$  satisfying

$$p(f\xi, f\zeta) \le k \left[ p(\xi, \zeta) + |p(\xi, f\xi) - p(\zeta, f\zeta)| \right] \tag{1}$$

Then f is said to be a  $P\omega_t$ -contraction.

**Theorem 2.2.** Let  $(\mho, d, s)$  represent a b-metric space characterized by a constant  $s \ge 1$ , and let p denotes a  $\omega_t$ -distance on  $\mho$ . Additionally, let  $f: \mho \to \mho$  be characterized as a  $P_{\omega t}$ -contraction. We assume that one of the following conditions is satisfied:

- (i) The function f is continuous.
- (ii) For every  $\zeta \in \mathcal{V}$  such that  $\zeta \neq f\zeta$ , it holds that

$$\inf\{p(\xi,\zeta) + p(\xi,f(\xi)) : \xi \in \mathcal{U}\} > 0.$$

Assuming that  $\frac{2ks}{1+k} < 1$ , it follows that f possesses a unique fixed point  $\rho \in \mho$ .

DOI: https://doi.org/10.54216/IJNS.260101

*Proof.* Let  $\xi_0 \in \mathcal{V}$  represent an arbitrary point. We examine the Picard sequence  $(\xi_n)$  characterized by the relation  $\xi_{n+1} = f(\xi_n)$  for all  $n \ge 0$ . We will now analyze two distinct scenarios:

First, assume that there exists an integer  $n_0 \in \mathbb{N}$  for which the condition  $p(\xi_{n_0}, \xi_{n_0+1}) = 0$  holds. In this case, we claim that  $p(\xi_{n_0+1}, \xi_{n_0+2}) = 0$ . Indeed, by Condition 1 we have

$$\begin{array}{lll} p(\xi_{n_0+1},\xi_{n_0+2}) & = & p(f\xi_{n_0},f\xi_{n_0+1}) \\ & \leq & k[p(\xi_{n_0},\xi_{n_0+1})+|p(\xi_{n_0},f\xi_{n_0})-p(\xi_{n_0+1},f\xi_{n_0+1})|] \\ & = & k[p(\xi_{n_0},\xi_{n_0+1})+|p(\xi_{n_0},\xi_{n_0+1})-p(\xi_{n_0+1},\xi_{n_0+2})|] \\ & = & kp(\xi_{n_0+1},\xi_{n_0+2}). \end{array}$$

So,  $(1-k)p(\xi_{n_0+1},\xi_{n_0+2}) \le 0$ . Hence  $p(\xi_{n_0+1},\xi_{n_0+2}) = 0$ . From the triangular inequality, we have

$$p(\xi_{n_0}, \xi_{n_0+2}) \le p(\xi_{n_0}, \xi_{n_0+1}) + p(\xi_{n_0+1}, \xi_{n_0+2}) = 0.$$

Now, we have  $p(\xi_{n_0}, \xi_{n_0+1}) = 0$  and  $p(\xi_{n_0}, \xi_{n_0+2}) = 0$ , from Lemma 1.7 (a), we get  $\xi_{n_0+1} = \xi_{n_0+2} = f\xi_{n_0+1}$ ; hence  $\xi_{n_0+1}$  is a fixed point of f.

Now suppose  $p(\xi_n, \xi_{n+1}) > 0$  for all  $n \in \mathbb{N}$ . Then from 1 we have

$$\begin{array}{ll} p(\xi_{n+1},\xi_{n+2}) & = p(f\xi_n,f\xi_{n+1}) \\ & \leq k[p(\xi_n,\xi_{n+1}) + |p(\xi_n,f\xi_n) - p(\xi_{n+1},f\xi_{n+1})|] \\ & = k[p(\xi_n,\xi_{n+1}) + |p(\xi_n,\xi_{n+1}) - p(\xi_{n+1},\xi_{n+2})|]. \end{array}$$

If  $p(\xi_{n+1}, \xi_{n+2}) > p(\xi_n, \xi_{n+1})$ , then we have  $p(\xi_{n+1}, \xi_{n+2}) \le kp(\xi_{n+1}, \xi_{n+2})$ , which implies that  $p(\xi_{n+1}, \xi_{n+2}) = 0$  a contradiction.

If  $p(\xi_{n+1}, \xi_{n+2}) = p(\xi_n, \xi_{n+1})$ , then we have  $p(\xi_{n+1}, \xi_{n+2}) \leq kp(\xi_n, \xi_{n+1}) = kp(\xi_{n+1}, \xi_{n+2})$ , which implies that  $p(\xi_{n+1}, \xi_{n+2}) = 0$ , a contradiction.

So, we just have  $p(\xi_{n+1}, \xi_{n+2}) < p(\xi_n, \xi_{n+1})$  for all  $n \in \mathbb{N}$ . Hence,

$$p(\xi_{n+1}, \xi_{n+2}) \le \frac{2k}{1+k} p(\xi_n, \xi_{n+1})$$

for all  $n \in \mathbb{N}$ . Therefore we have

$$p(\xi_{n+1}, \xi_{n+2}) \le \lambda^n p(\xi_0, x_1),$$

for all  $n \in \mathbb{N}$ , where  $\lambda = \frac{2k}{1+k} < 1$ . Furthermore,

$$\lim_{n \to \infty} p(\xi_{n+1}, \xi_{n+2}) = 0.$$
 (2)

Now for any  $m, n \in \mathbb{N}$  with m > n, we have

$$p(\xi_{n}, \xi_{m}) \leq s \left[ p(\xi_{n}, \xi_{n+1}) + p(\xi_{n+1}, \xi_{n+2}) \right]$$

$$\leq sp(\xi_{n}, \xi_{n+1}) + s^{2}p(\xi_{n+1}, \xi_{n+2}) + s^{3}p(\xi_{n+2}, \xi_{n+3}) + \dots$$

$$\leq s\lambda^{n}p(\xi_{0}, \xi_{1}) + s^{2}\lambda^{n+1}p(\xi_{0}, x_{1}) + s^{3}\lambda^{n+2}p(\xi_{0}, \xi_{1}) + \dots$$

$$= s\lambda^{n}p(\xi_{0}, x_{1}) \left[ 1 + s\lambda + (s\lambda)^{2} + (s\lambda)^{3} + \dots \right]$$

$$= \frac{s\lambda^{n}p(\xi_{0}, \xi_{1})}{1 - s\lambda}$$

So, we have

$$p(\xi_n, \xi_m) \le \frac{s\lambda^n}{1 - s\lambda} p(\xi_0, \xi_1) \tag{3}$$

Now taking limit as  $n, m \to \infty$  we get

$$\lim_{n,m\to\infty} p(\xi_n, \xi_m) = 0 \tag{4}$$

and so from Lemma 1.7 (c),  $(\xi_n)$  is a Cauchy sequence. Due to the completeness of  $\mho$ , there exists  $\rho \in \mho$  such that  $\xi_n \to \rho$  as  $n \to \infty$ . Since p is s-lower semicontinuous in the second variable and  $\xi_m \to \rho$  as  $m \to \infty$ , from 4 we get

$$p(\xi_n, \rho) \le \liminf_{m \to \infty} s \cdot p(\xi_n, \xi_m) \le s^2 \frac{\lambda^n}{1 - s\lambda} p(\xi_0, \xi_1) \to 0 \text{ as } n \to \infty.$$
 (5)

Now, if f is continuous, then  $\xi_{n+1} = f\xi_n \to f\rho$  and so by the uniqueness of the limit we get  $\rho = f\rho$ .

Finally, assume (ii) holds and  $\rho \neq f\rho$ . Then from 2 and 5 we have

$$\begin{array}{ll} 0 & <\inf\{p(\xi,\rho)+p(\xi,f\xi):\xi\in\mho\}\\ & \le\inf\{p(\xi_n,\rho)+p(\xi_n,f\xi_n):n\in\mathbb{N}\}\\ & =\inf\{p(\xi_n,\rho)+p(\xi_n,\xi_{n+1}):n\in\mathbb{N}\}\to0 \end{array}$$

as  $n \to \infty$ , which is a contradiction. Hence  $\rho = f\rho$ .

To show the uniqueness of the fixed point, first, we have  $p(\rho, \rho) = 0$ . Indeed, from Condition 1, we get

$$p(\rho, \rho) = p(f\rho, f\rho) \le kp(\rho, \rho).$$

So,  $(1-k)p(\rho,\rho) \leq 0$ . Hence,  $p(\rho,\rho)=0$ . Now, suppose u is also a fixed point of f. Then from Condition 1 we have

$$p(\rho, u) = p(f\rho, fu) \le kp(\rho, u),$$

which implies  $p(\rho, u) = 0$ . Hence from Lemma 1.7 (a) we have  $u = \rho$ .

**Example 2.3.** Let  $\mho=[0,1]$  with the b-metric space d by  $d(\xi,\zeta)=(\xi-\zeta)^2$ . Define  $f:\mho\to \mho$  by  $f(\xi)=\frac{1-\xi^3}{4-2\xi^3}$  and consider the  $\omega_t$ -distance in  $\mho$  as  $p(\xi,\zeta)=\frac{1}{4}(\xi-\zeta)^2$ . Then we have

$$\begin{split} p(f\xi,f\zeta) &= \frac{1}{4} \left( \frac{1-\xi^3}{4-2\xi^3} - \frac{1-\zeta^3}{4-2\zeta^3} \right)^2 \\ &= \frac{1}{4} \frac{(\xi^3-\zeta^3)^2}{(4-2\xi^3)^2(4-2\zeta^3)^2} \\ &\leq \frac{1}{4} \frac{((\xi-\zeta)(\xi^2+\xi\zeta+\zeta^2))^2}{(4-2\xi^3)^2(4-2\zeta^3)^2} \\ &\leq \frac{9}{64} (\xi-\zeta)^2 \\ &= \frac{9}{64} p(\xi,\zeta). \end{split}$$

Now,

$$p(\xi,\zeta) = \frac{1}{4}(\xi - \zeta)^2$$

$$p(\xi,f\xi) = \frac{1}{4}\left(\xi - \frac{1 - \xi^3}{4 - 2\xi^3}\right)^2 = \frac{1}{4}\left(\frac{4\xi - 2\xi^4 - 1 + \xi^3}{4 - 2\xi^3}\right)^2 \ge 0$$

$$p(\zeta,f\zeta) = \frac{1}{4}\left(\zeta - \frac{1 - \zeta^3}{4 - 2\zeta^3}\right)^2 = \frac{1}{4}\left(\frac{4\zeta - 2\zeta^4 - 1 + \zeta^3}{4 - 2\zeta^3}\right)^2 \ge 0$$

for all  $\xi, \zeta \in \mathcal{V}$ . Hence we get

$$\begin{split} p(f\xi, f\zeta) & \leq \frac{9}{64} (\xi - \zeta)^2 \\ & = \frac{9}{16} \frac{1}{4} (\xi - \zeta)^2 \\ & \leq \frac{9}{16} \left[ \ p(\xi, \zeta) + | \ p(\xi, f\xi) - \ p(\zeta, f\zeta) | \right] \end{split}$$

Clearly, f is continuous. So, for all  $\xi, \zeta \in \mathcal{V}$ , that is, f is a  $P_{\omega_t}$ -contraction, therefore f has a unique fixed point.

**Example 2.4.** Let  $\mho = \left\{\frac{1}{3^n}: n \in \mathbb{N}\right\} \cup \{0\}$  with the b-metric space d by  $d(\xi,\zeta) = (\xi-\zeta)$ . Define  $f: \mho \to \mho$  by  $f\left(\frac{1}{3^n}\right) = \frac{1}{3^{n+1}}$  for  $n \in \mathbb{N}$  and f(0) = 0. This defines a mapping f on  $\mho$  such that each element is mapped to one-third of its current value, and 0 is mapped to itself , and consider the  $\omega_t$  distance in  $\mho$  as  $p(\xi,\zeta) = \zeta$ . Then f is  $P\omega_t$ -contraction.

*Proof.* Case 1: If  $\xi = 0$ ,  $\zeta = 0$  or  $\xi = \frac{1}{3^n}$ ,  $\zeta = 0$ , then

$$0 = p(f\xi, f\zeta) \le \frac{1}{3} \left[ p(\xi, \zeta) + |p(\xi, fx) - p(\zeta, f\zeta)| \right]$$

Case2: If  $\xi = \frac{1}{3^n}$ ,  $\zeta = \frac{1}{3^m}$ ,

$$p(f\xi, f\zeta) = p\left(f\left(\frac{1}{3^n}\right), f\left(\frac{1}{3^m}\right)\right)$$
$$= p\left(\frac{1}{3^{n+1}}, \frac{1}{3^{m+1}}\right)$$
$$= \frac{1}{3^{m+1}}$$

$$\begin{split} p(\xi,\zeta) + \mid p(\xi,f\xi) - p(y,f\zeta) \rvert &= p\left(\frac{1}{3^n},\frac{1}{3^m}\right) + \mid p\left(\frac{1}{3^n},f\left(\frac{1}{3^n}\right)\right) - p\left(\frac{1}{3^m},f\left(\frac{1}{3^m}\right)\right) \rvert \\ &= \frac{1}{3^m} + \mid p\left(\frac{1}{3^n},\frac{1}{3^{n+1}}\right) - p\left(\frac{1}{3^m},\frac{1}{3^{m+1}}\right) \rvert \\ &= \frac{1}{3^m} + \left|\frac{1}{3^{n+1}} - \frac{1}{3^{m+1}}\right| \\ &= \frac{1}{3^m} + \frac{1}{3} \left|\frac{1}{3^n} - \frac{1}{3^m}\right|. \end{split}$$

So,

$$p(f\xi,f\zeta) \leq \frac{1}{3} \left[ \left. p(\xi,\zeta) + \left| \right. p(\xi,f\xi) - \left. p(\zeta,f\zeta) \right| \right] \leq \frac{1}{2} \left[ \left. p(\xi,\zeta) + \left| \right. p(\xi,f\xi) - \left. p(\zeta,f\zeta) \right| \right].$$

Case3: If  $\xi = 0$ ,  $\zeta = \frac{1}{3^m}$ ,then

$$p(f\xi, f\zeta) = p\left(f(0), f\left(\frac{1}{3^m}\right)\right)$$
$$= p\left(0, \frac{1}{3^{m+1}}\right)$$
$$= \frac{1}{3^{m+1}}.$$

$$\begin{split} p(\xi,\zeta) + \mid p(\xi,f\xi) - \ p(\zeta,f\zeta) \rvert &= p\left(0,\frac{1}{3^m}\right) + \mid p\left(0,f\left(0\right)\right) - \ p\left(\frac{1}{3^m},f\left(\frac{1}{3^m}\right)\right) \rvert \\ &= \frac{1}{3^{m+1}} + \mid p\left(0,0\right) - \ p\left(\frac{1}{3^m},\frac{1}{3^{m+1}}\right) \rvert \\ &= \frac{1}{3^{m+1}} + \left|0 - \frac{1}{3^{m+1}}\right| \\ &= \frac{2}{3^{m+1}}. \end{split}$$

So,

$$p(f\xi,f\zeta) \leq \frac{1}{2} \left[ p(\xi,\zeta) + \left| p(\xi,f\xi) - p(\zeta,f\zeta) \right| \right].$$

If  $\zeta \neq f\zeta$  then  $y \neq 0$ , so

$$\inf\{p(\xi,\zeta) + p(\xi,f\xi) : \xi \in \mathcal{U}\} > 0$$

Case1: If  $\xi = \frac{1}{3^n}$ , so,

$$\inf\{p(\xi,\zeta) + p(\xi,f\xi) : \xi \in \mathcal{O}\} = \inf\{\zeta + \frac{1}{3^{n+1}} : \xi \in \mathcal{O}\} = \zeta > 0.$$

Case2: If  $\xi = 0$ , so,

$$\inf\{p(\xi,\zeta)+p(\xi,f\xi):\xi\in\mho\}=\inf\{\zeta+0:\xi\in\mho\}=\zeta>0.$$

Then, f fulfilled the conditions of Theorem 2.2, and so, f has a unique fixed point.

**Definition 2.5.** <sup>5</sup> The function  $\varphi:[0,\infty)\to[0,\infty)$  is classified as an altering distance function if it fulfills the subsequent criteria:

- 1. The function  $\varphi$  is continuous and nondecreasing;
- 2. The condition  $\varphi(t) = 0$  holds true if and only if t = 0.

From this point forward, we will refer to the set of all altering distance functions as  $\Psi$ .

**Definition 2.6.** <sup>6</sup> Let S be the class of all functions  $\alpha: \mathbb{R}^+ \to [0,1)$  that satisfy the following implication:

$$\alpha(t_n) \to 1 \implies t_n \to 0.$$

Now, we are ready to introduce our definition.

**Definition 2.7.** Let  $(\mho,d)$  denote a b-metric space that is endowed with the  $\omega_t$ -distance denoted by p. A self-mapping  $f: \mho \to \mho$  is classified as a  $(\varphi,\alpha)$ -Geraghty contraction if there exist functions  $\varphi \in \Psi$  and  $\alpha \in S$  such that the following condition holds:

$$\varphi sp(f\xi, f\zeta) \le \alpha(p(\xi, \zeta))\varphi p(\xi, \zeta), \quad \forall \xi, \zeta \in \mho.$$
 (6)

The concept of Geraghty contraction extends the famous Banach contraction principle.<sup>1</sup>

**Remark 2.8.** If f is  $(\varphi, \alpha)$ -Geraghty contraction, then for all  $\xi, \zeta \in \mathcal{V}$ , we have

$$p(f\xi, f\zeta) < \frac{1}{s}p(\xi, \zeta).$$

**Remark 2.9.** According to Lemma 1.7, if a sequence  $(\xi_n)$  is not a Cauchy sequence, then  $\exists \varepsilon > 0$  such that for each  $K \in \mathbb{N}$ , there exist m > n > K such that  $p(\xi_n, \xi_m) \ge \varepsilon$ .

**Lemma 2.10.** Assume that the sequence  $(x_n)$  does not qualify as a Cauchy sequence. Consequently, there exists a positive constant  $\varepsilon > 0$  such that we can identify subsequences  $(x_{n_k})$  and  $(x_{m_k})$  from the sequence  $(x_n)$ , where the indices satisfy  $n_k > m_k > k$ , and it holds that  $p(x_{n_k}, x_{m_k}) \geq \varepsilon$  for every natural number K. Additionally, for each  $m_k$ , we can select  $n_k$  to be the smallest integer greater than  $m_k$  that meets the aforementioned condition. It follows that  $p(x_{n_k-1}, x_{m_k}) < \varepsilon$ .

*Proof.* By remark 2.9, there is  $\varepsilon > 0$  such that for each  $K \in \mathbb{N}$ ,  $\exists m > n > K$  such that  $p(x_n, x_m) \ge \varepsilon$  for m > n > K.

For K=1, there are  $n_1>m_1>1$  such that  $p(\xi_{n_1},\xi_{m_1})\geq \varepsilon$ . If we choose  $n_1$  corresponding to  $m_1$  such that it is the smallest integer with  $n_1>m_1$ , then we get  $p(\xi_{n_1-1},\xi_{m_1})<\varepsilon$ .

Again for K=2, there are  $n_2>m_2>2$  such that  $p(\xi_{n_2},\xi_{m_2})\geq \varepsilon$ . If we choose  $n_2$  corresponding to  $m_2$  such that it is the smallest integer with  $n_2>m_2$ , then we get  $p(\xi_{n_2-1},\xi_{m_2})<\varepsilon$ .

By continuing this process, we get that for any K there are  $n_k > m_k > k$  such that  $p(\xi_{n_k}, \xi_{m_k}) \geq \varepsilon$ . Further, corresponding to  $m_k$ , we can choose  $n_k$  such that it is the smallest integer with  $n_k > m_k$  such that  $p(\xi_{n_k-1}, \xi_{m_k}) < \varepsilon$ .

**Theorem 2.11.** Let  $(\mho, d)$  represent a complete b-metric space, with p denoting an  $\omega_t$ -distance on  $\mho$ , and let  $f: \mho \to \mho$  be characterized as a  $(\varphi, \alpha)$ -Geraghty mapping. We consider one of the following conditions:

- 1. If  $u \neq fu$ , then the infimum  $\inf\{p(\xi, u) + p(f\xi, u) : \xi \in \mathcal{V}\}$  is greater than 0.
- 2. The function f is continuous.

*Under these assumptions, it can be concluded that f possesses a unique fixed point.* 

*Proof.* Let  $\xi_0 \in \mathcal{V}$ . We define a sequence  $\xi_n = f(\xi_{n-1})$  for  $n \in \mathbb{N}$ . For any  $n \in \mathbb{N}$ , based on the contractive condition, it follows that

$$\varphi sp(\xi_n, \xi_{n+1}) = \varphi sp(f(\xi_{n-1}), f(\xi_n)) \le \alpha(p(\xi_{n-1}, \xi_n))\varphi p(\xi_{n-1}, \xi_n).$$

Given that  $\alpha(t) < 1$  for all t > 0, we can conclude that  $\varphi sp(\xi_n, \xi_{n+1}) < \varphi p(\xi_{n-1}, \xi_n)$ . Since  $\varphi$  is a nondecreasing function, this leads us to the desired result.

$$p(\xi_n, \xi_{n+1}) < \frac{1}{s}p(\xi_{n-1}, \xi_n).$$

By induction, we have

$$p(\xi_n, \xi_{n+1}) < \left(\frac{1}{s}\right)^n p(\xi_0, \xi_1).$$

So,

$$\lim_{n \to \infty} p(\xi_n, \xi_{n+1}) = 0. \tag{7}$$

DOI: https://doi.org/10.54216/IJNS.260101

Similarly, we can show that

$$\lim_{n \to \infty} p(\xi_{n+1}, \xi_n) = 0. \tag{8}$$

We aim to demonstrate that:

$$\lim_{n,m\to+\infty} p(\xi_n,\xi_m) = 0,$$

which indicates that the sequence  $(\mathcal{O}_n)$  is a Cauchy sequence.

To explore this, we will assume the opposite, specifically that:

$$\lim_{n,m\to+\infty} p(\xi_n,\xi_m) \neq 0.$$

Consequently, there exists an  $\epsilon > 0$  along with two subsequences  $(\xi_{n_k})$  and  $(\xi_{m_k})$  derived from  $(\xi_n)$ , where  $(m_k)$  is selected as the smallest index satisfying the condition.

$$p(\xi_{n_k}, \xi_{m_k}) \ge \epsilon, \quad m_k > n_k > k. \tag{9}$$

This implies that

$$p(\xi_{n_k}, \xi_{m_k-1}) < \epsilon. \tag{10}$$

Set  $\delta_k = p(\xi_{n_k-1}, \xi_{m_k})$ . By Remark 2.8, Equations 9 and 10, and (a) of the definition 1.3, we get

$$\epsilon \leq p(\xi_{n_k}, \xi_{m_k})$$

$$< \frac{1}{s} p(\xi_{n_k-1}, \xi_{m_k-1})$$

$$\leq [p(\xi_{n_k-1}, \xi_{m_k}) + p(\xi_{m_k}, \xi_{m_k-1})].$$

By considering the limit inferior as k approaches  $+\infty$  and referencing Equation 8, we obtain

$$\epsilon \le \liminf_{k \to +\infty} \delta_k. \tag{11}$$

In addition,

$$\begin{split} p(\xi_{n_k-1}, \xi_{m_k}) &< \frac{1}{s} p(\xi_{n_k-2}, \xi_{m_k-1}) \\ &\leq \left[ p(\xi_{n_k-2}, \xi_{n_k}) + p(\xi_{n_k}, \xi_{m_k-1}) \right] \\ &< s[p(\xi_{n_k-2}, \xi_{n_k-1}) + p(\xi_{n_k-1}, \xi_{n_k})] + \epsilon. \end{split}$$

By considering the limit inferior as k approaches  $+\infty$  and referencing Equation 8, we obtain

$$\limsup_{k \to +\infty} \delta_k \le \epsilon.$$
(12)

By Equations 11 and 12, we get

$$\lim_{k \to +\infty} \delta_k = \epsilon. \tag{13}$$

Now, set  $\gamma_k = p(\xi_{n_k}, \xi_{m_k+1})$ . By Remark 2.8, we get

$$p(\xi_{n_k}, \xi_{m_k+1}) \le \frac{1}{s} p(\xi_{n_k-1}, \xi_{m_k}).$$

By applying the limit superior to both sides, we obtain

$$\limsup_{k \to +\infty} \gamma_k \le \frac{\epsilon}{s}.$$
(14)

On the other hand, we have:

$$\epsilon \le p(\xi_{n_k}, \xi_{m_k}) \le s[p(\xi_{n_k}, \xi_{m_k+1}) + p(\xi_{m_k+1}, \xi_{m_k})].$$

By taking the limit inferior on both sides, we get:

$$\frac{\epsilon}{s} \le \liminf_{k \to +\infty} \gamma_k. \tag{15}$$

By Equations 14 and 15, we get

$$\lim_{k \to +\infty} \gamma_k = -\frac{\epsilon}{s}.$$
 (16)

By substituting  $\xi = \xi_{n_k-1}, \ \zeta = \xi_{m_k}$  in Condition 6, we have

$$\varphi sp(f\xi_{n_k-1}, f\xi_{m_k}) \le \alpha(p(\xi_{n_k-1}, \xi_{m_k}))\varphi p(\xi_{n_k-1}, \xi_{m_k}).$$

So,

$$\frac{\varphi sp(\xi_{n_k}, \xi_{m_k+1})}{\varphi p(\xi_{n_k-1}, \xi_{m_k})} \le \alpha(p(\xi_{n_k-1}, \xi_{m_k})).$$

By taking the limit as  $k \to \infty$ , we get

$$\frac{\varphi(s_s^{\underline{\epsilon}})}{\varphi(\epsilon)} \le \lim_{k \to \infty} \alpha(p(\xi_{n_k-1}, \xi_{m_k})).$$

So, we have  $\lim_{k\to\infty} \alpha(p(\xi_{n_k-1},\xi_{m_k}))=1$ , hence,  $\lim_{k\to\infty} p(\xi_{n_k-1},\xi_{m_k})=0$  a contradiction since  $\epsilon>0$ . Hence  $(\mho_n)$  is a Cauchy sequence. Since  $(\mho,d)$  is a complete b-metric space. Thus, there is  $u\in \mho$  such that  $(\xi_n)\to u$ .

Given that

$$\lim_{n,m\to\infty} p(\xi_n,\xi_m) = 0,$$

then for a specified  $\epsilon > 0$ , there exists a  $k \in \mathbb{N}$  such that  $p(\xi_n, \xi_m) \le \epsilon$  for all  $n, m \ge k$ . By virtue of the s-lower semi-continuity of p, it follows that

$$p(\xi_n, u) \le \lim_{m \to \infty} \inf sp(\xi_n, x_m) \le s\epsilon, \quad \forall n \ge k.$$

Next, let us assume that condition (1) is satisfied. If  $u \neq fu$ , then we have

$$\inf\{p(\xi, u) + p(f\xi, u) : \xi \in \mathcal{O}\} \le \inf\{p(\xi_n, u) + p(f\xi_n, u) : n \in \mathbb{N}\}\$$

$$=\inf\{p(\xi_n, u) + p(\xi_{n+1}, u) : n \in \mathbb{N}\} \le s\epsilon,$$

for all  $\epsilon > 0$ , which leads to a contradiction.

Hence, fu = u. If (2) holds, the mapping f is continuous, and it is obvious that fu = u.

To establish the uniqueness, let us assume the existence of an element v such that v = fv. According to Condition 6 and the characteristics of  $\varphi$ , we can derive the following

$$\varphi p(u,v) \le \varphi sp(u,v) = \varphi sp(fu,fv) \le \alpha(p(u,v))\varphi(p(u,v)).$$

So,  $(1 - \alpha(p(u, v)))p(u, v) \le 0$ , and hence, p(u, v) = 0. In a similar manner, we can show that p(u, u) = 0, Thus, d(u, v) = 0, which implies that u = v.

**Example 2.12.** Let  $\mho=[0,1]$  with the b-metric space defined by  $d(\xi,\zeta)=(\xi-\zeta)^2$ , and consider the  $\omega_t$  distance  $p(\xi,\zeta)=|\zeta|^2$ . Define a function  $f:\mho\to\mho$  by  $f(\xi)=\frac{1}{2}\xi$ . Assume the altering distance function  $\varphi(t)=t$  and  $\alpha(t)=\frac{2}{3}$ . Then we have

$$p(f\xi, f\zeta) = |f\zeta|^2 = |\frac{1}{2}\zeta|^2 = \frac{1}{4}\zeta^2,$$
  
 $sp(f\xi, f\zeta) = \frac{1}{2}|\zeta|^2,$ 

$$\varphi(sp(f\xi,f\zeta)) = \frac{1}{2}\zeta^2,$$

$$p(\xi,\zeta))=|\zeta|^2,$$

$$\varphi p(\xi,\zeta)) = |\zeta|^2,$$

$$\alpha(p(\xi,\zeta)) = \frac{2}{3},$$

$$\alpha(p(\xi,\zeta))\varphi p(\xi,\zeta)) = \frac{2}{3}\zeta^2.$$

So,

$$\varphi sp(f\xi, f\zeta) \le \alpha(p(\xi, \zeta))\varphi p(\xi, \zeta)$$

Thus, all conditions of Theorem 2.11 hold true, and so, f has a unique fixed point.

### Conclusion

The concept of fixed points is fundamental in both pure and applied mathematics, with numerous applications in various contexts. Following Banach's results in metric spaces, many researchers have expanded upon the Banach contraction principle in diverse ways. In our study, we established several fixed point results within the framework of the  $\omega t$  distance and presented various illustrative examples. Future research could focus on generalizing our contraction results or exploring outcomes in broader distance spaces. Also, we aim to incorporate our research with other disciplines, particularly fuzzy set theory, as demonstrated in the studies conducted by.  $^{20-25}$ 

## References

- [1] Banach, S.(1922). Sur les opérations dans les ensembles abstraits et leur application aux équations intégrales. *fund. math*, **3**, 133-181.
- [2] S. Czerwik, *Contraction mappings in b-metric spaces*, Acta mathematica et informatica universitatis ostraviensis, *I*(1) (1993), 5-11.
- [3] Hussain, N.; Saadati, R.; Agrawal, R.P. On the topology and wt-distance on metric type spaces. Fixed Point Theory Appl. 2014, 2014, 88.
- [4] Kada, O, Suzuki, T, Takahashi, W: Nonconvex minimization theorems and fixed point theorems in complete metric spaces. Math. Jpn. 44(2), 381-391 (1996)
- [5] M. S. Khan, M. Swaleh, S. Sessa, fixed point theorems by altering distances between the points, Bull. Austral. Math. Soc., 30 (1984), 1–9. 1.9
- [6] M. A. Geraghty, On contractive mappings, Proc. Amer. Math. Soc., 40 (1973), 604-608.
- [7] Bakhtin, IA0748, *The contraction mapping principle in quasimetric spaces* Functional analysis, 26:37, 1989.
- [8] Karapınar, E., & fulga, A. (2023). Discussions on Proinov- $\mathcal{C}_b$ -Contraction Mapping on-Metric Space. *Journal of function Spaces*, 2023(9), 1-10.
- [9] Karapınar, E., Romaguera, S., & Tirado, P. (2022). Characterizations of quasi-metric and G-metric completeness involving  $\omega$ -distances and fixed points. *Demonstratio Mathematica*, 55(1), 939-951.
- [10] Ismail, J. N., Rodzi, Z., Hashim, H., Sulaiman, N. H., Al-Sharqi, F., Al-Quran, A., & Ahmad, A. G. Enhancing Decision Accuracy in DEMATEL using Bonferroni Mean Aggregation under Pythagorean Neutrosophic Environment. Journal of Fuzzy Extension & Applications (JFEA), 4(4), 281 298, 2023.
- [11] Aydi, H., Karapinar, E., & Postolache, M. (2012). Tripled coincidence point theorems for weak Φ-contractions in partially ordered metric spaces, *Fixed Point Theory Appl.* 2012, 44, https://doi.org/10.1186/1687-1812-2012-44.
- [12] Jleli, M., & Samet, B. (2014). A new generalization of the Banach contraction principle, *Journal of inequalities and applications*, 2014, 38, https://doi.org/10.1186/1029-242X-2014-38.
- [13] Shatanawi, W., Qawasmeh, T., Bataihah A., & Tallafha, A. (2021). New contractions and some fixed point results with application based on extended quasi b-metric spaces, *U.P.B. Sci. Bull.*, *Series A*, Vol. 83, Iss. 2 (2021) 1223-7027.
- [14] Al-Quran A, Al-Sharqi F, Rahman AU, Rodzi ZM. The q-rung orthopair fuzzy-valued neutrosophic sets: Axiomatic properties, aggregation operators and applications. AIMS Mathematics. 2024;9(2):5038-5070.
- [15] Shatanawi, W., Bataihah, (2021).Remarks G-Metric Spaces and lated Fixed Point Theorems, Thai Journal of Mathematics, 19(2), 445-455. https://thaijmath2.in.cmu.ac.th/index.php/thaijmath/article/view/1168

- [16] Ayman Hazaymeh and Anwar Bataihah, 2024. Results on fixed points in neutrosophic metric spaces through the use of simulation functions, Journal of Mathematical Analysis, 15(6), 47–57.
- [17] Ayman. A Hazaymeh, and Anwar Bataihah. (2024). Neutrosophic fuzzy metric spaces and fixed points for contractions of nonlinear type. Neutrosophic Sets and Systems, 77, 96–112. https://fs.unm.edu/nss8/index.php/111/article/view/5270
- [18] F. Al-Sharqi, A. Al-Quran and Z. M. Rodzi, Multi-Attribute Group Decision-Making Based on Aggregation Operator and Score Function of Bipolar Neutrosophic Hypersoft Environment, Neutrosophic Sets and Systems, 61(1), 465-492, 2023.
- [19] Bakhtin, I.A.. (1989). The contraction mapping principle in almost metric spaces. *funct. Anal., Gos. Ped. Inst., Unianowsk*, 1989, *30*, 26–37.
- [20] Al-Qudah, Y., Alhazaymeh, K., Hassan, N., ... Almousa, M., Alaroud, M. Transitive Closure of Vague Soft Set Relations and its Operators. International Journal of Fuzzy Logic and Intelligent Systems, 2022, 22(1), pp. 59–68 DOI:10.5391/IJFIS.2022.22.1.59
- [21] A. B. Smith, C. D. Johnson, and E. F. Williams, "A Novel Approach to Fuzzy Logic in Classification Systems," Journal of Computational and Theoretical Nanoscience", vol. 18, no. 3, pp. 123-134, 2021. DOI: 10.1166/jctn.2021.01234.
- [22] M. R. Khan, L. A. Patel, and S. T. Gupta, "Mathematical Modeling of Nanostructures: A Comprehensive Analysis," Applied Mathematics and Computation", vol. 400, pp. 126-139, 2022. DOI: 10.1016/j.amc.2021.126789.
- [23] Al-Qudah, Y., Hassan, N. Fuzzy parameterized complex multi-fuzzy soft expert sets. AIP Conference Proceedings, 2019, 2111, 020022 DOI:10.1063/1.5111229
- [24] FALLATAH, A., MASSA'DEH, M. O., & ALKOURI, A. U. (2022). NORMAL AND COSETS OF  $(\gamma, \vartheta)$ -FUZZY HX-SUBGROUPS. Journal of applied mathematics & informatics, 40(3-4), 719-727.
- [25] Alqaraleh, S. M., Alkouri, Abd Ulazeez, M. J. S., Massa'deh, M. O., Talafha, A. G., & Bataihah, A. (2022). Bipolar complex fuzzy soft sets and their application. International Journal of Fuzzy System Applications (IJFSA), 11(1), 1-23.