



## A Contribution to the Group of Units' Problem in Some 2-Cyclic Refined Neutrosophic Rings

Basheer Abd Al Rida Sadiq

Al-Imam Al-Kadhumi College For Islamic Science University, Iraq

E-mail: [basheer.abdrida@alkadhumi-col.edu.iq](mailto:basheer.abdrida@alkadhumi-col.edu.iq)

### Abstract:

This paper is dedicated to study the group of units problem in some 2-cyclic refined neutrosophic rings, where it presents a full classification of the group of units in 2-cyclic refined neutrosophic ring of integers  $Z_2(I)$ , 2-cyclic refined neutrosophic ring of integers modulo 2 and 3  $(Z_2)_2(I)$ ,  $(Z_3)_2(I)$  and 2-cyclic refined neutrosophic ring of real numbers  $R_2(I)$ , as direct products of the cyclic groups. On the other hand, this work provides a necessary and sufficient condition for the invertibility of any square 2-cyclic refined neutrosophic real matrix. Also, we will illustrate some examples to clarify the validity of our work.

**Keywords:** 2-cyclic refined rings; group of units; invertible; neutrosophic ring

### 1. Introduction

In the theory of rings, the problem of determining units (invertible elements) in a ring with unity is still open. The classification of the group of units is still unknown in general. After the arrival of neutrosophy in 1995, it was used to define some new kinds of algebraic structures, where we find concepts such as neutrosophic rings, refined neutrosophic modules and spaces, refined neutrosophic ring of matrices, n-refined neutrosophic rings [3-6,7-12], and n-cyclic refined neutrosophic rings [2]. The condition of invertibility in n-cyclic refined neutrosophic rings is still an open question in general for any positive integer n.

This open question has motivated us to study the case of 2-cyclic refined neutrosophic rings. We determine the units in 2-cyclic refined neutrosophic ring of integers  $Z_2(I)$ , 2-cyclic refined neutrosophic ring of integers modulo 2 and 3  $(Z_2)_2(I)$ ,  $(Z_3)_2(I)$  and 2-cyclic refined neutrosophic ring of real numbers  $R_2(I)$ , with the corresponding structure of the group of units by using a computational method depends only on the definition of multiplication operation. Also, we have determined a necessary and sufficient condition for the invertibility of any square 2-cyclic refined neutrosophic real matrix. All rings through this paper are considered with unity 1.

## 2. Preliminaries

### Definition 1: [2]

Let  $(R, +, \times)$  be a ring and  $I_k; 1 \leq k \leq n$  be  $n$  sub-indeterminacies. We define  $R_n(I) = \{a_0 + a_1I + \dots + a_nI_n; a_i \in R\}$  to be  $n$ -cyclic refined neutrosophic ring.

Operations on  $R_n(I)$  are defined as:

$$\sum_{i=0}^n x_i I_i + \sum_{i=0}^n y_i I_i = \sum_{i=0}^n (x_i + y_i) I_i, \sum_{i=0}^n x_i I_i \times \sum_{i=0}^n y_i I_i = \sum_{i,j=0}^n (x_i \times y_j) I_i I_j = \sum_{i,j=0}^n (x_i \times y_j) I_{(i+j \bmod n)}.$$

$\times$  is the multiplication on the ring  $R$ .

### Example 2:

(a) The 2-cyclic refined neutrosophic ring of integers is defined as follows:

$$Z_2(I) = \{t_0 + t_1 I_1 + t_2 I_2; t_i \in Z\}.$$

(b) Addition on  $Z_2(I)$  can be clarified as follows:

$$(a + bI_1 + cI_2) + (m + nI_1 + tI_2) = (a + m) + I_1(b + n) + I_2(c + t).$$

(c) Multiplication on  $Z_2(I)$  can be clarified as follows:

$$\begin{aligned} (a + bI_1 + cI_2)(m + nI_1 + tI_2) &= am + anI_1 + atI_2 + bmI_1 + bnI_2 + btI_1 + cmI_2 + cnI_1 + ctI_2 \\ &= am + I_1(an + bm + bt + cn) + I_2(at + bn + cm + ct). \end{aligned}$$

Where  $I_1 I_1 = I_{(1+1 \bmod 2)} = I_2, I_2 I_2 = I_{(2+2 \bmod 2)} = I_2, I_1 I_2 = I_{(1+2 \bmod 2)} = I_1$ .

The elements of  $Z_2$  are taken by the form  $\{1, 2\}$  instead of  $\{0, 1\}$  in the definition of indices  $I$  in sub-indeterminacies  $I_i$ . See [2].

### Definition 3:

Let  $R$  be any ring with unity. An arbitrary element  $x \in R$  is called a unit if and only if there exists  $y \in R$  such that  $xy = yx = 1$ . The element  $y$  is called the inverse of  $x$ .

### Example 4:

Let  $Z_{4_2}(I) = \{a + bI_1 + cI_2; a, b, c \in Z_4\}$  be the 2-cyclic refined neutrosophic ring of integers modulo 4. The element  $x = 1 + 2I_2$  is a unit, that is because there exists  $y = x = 1 + 2I_2$ , such that  $xy = yx = 1 + 4I_2 + 4I_2 = 1$ . It is clear that the inverse of  $x$  is  $x$  itself.

## 3. Main concepts and results

### Theorem

Let  $R$  be any ring with unity, let  $R_2(I)$  be its corresponding 2-cyclic refined neutrosophic ring.

Let  $A = a_0 + a_1I_1 + a_2I_2$  be a unit in  $R_2(I)$ , then there exists  $B = b_0 + b_1I_1 + b_2I_2 \in R_2(I)$  such that

- 1).  $a_0b_0 = 1$ .
- 2).  $(a_0 + a_1 + a_2)(b_0 + b_1 + b_2) = 1$ .

Proof.

Suppose that  $A$  is a unit, there exists  $B = b_0 + b_1I_1 + b_2I_2 \in R_2(I)$ , such that  $A.B = 1$ .

Thus,  $a_0b_0 + I_1(a_1b_2 + a_2b_1 + a_0b_1 + a_1b_0) + I_2(a_2b_0 + a_0b_2 + a_2b_2 + a_1b_1) = 1$ , then

$$\begin{cases} a_0b_0 = 1 \dots (1) \\ a_1b_2 + a_2b_1 + a_0b_1 + a_1b_0 = 0 \dots (2) \\ a_2b_0 + a_0b_2 + a_2b_2 + a_1b_1 = 0 \dots (3) \end{cases}$$

By Adding (1) and (2) and (3), we get.

$$(a_0 + a_1 + a_2)(b_0 + b_1 + b_2) = 1, \text{ hence the proof is complete.}$$

### Theorem:

Let  $R_2(I) = \{t_0 + t_1I_1 + t_2I_2; t_i \in Z\}$  be the 2-cyclic refined neutrosophic ring of integers, then it has exactly 8 units.

Proof.

Suppose that  $A = a_0 + a_1I_1 + a_2I_2 \in Z_2(I)$  is a unit, then  $B = A^{-1} = b_0 + b_1I_1 + b_2I_2$ , where  $a_0b_0 = 1 (I)$  and  $(a_0 + a_1 + a_2)(b_0 + b_1 + b_2) = 1(II)$ .

From equation (II), we get:

$$a_0 + a_1 + a_2 = b_0 + b_1 + b_2 = 1 \text{ or}$$

$$a_0 + a_1 + a_2 = b_0 + b_1 + b_2 = -1.$$

We discuss the possible cases.

Case 1: if  $a_0 = b_0 = 1$  and  $a_0 + a_1 + a_2 = b_0 + b_1 + b_2 = 1$ , hence  $a_1 + a_2 = b_1 + b_2 = 0$ , thus  $a_2 = -a_1, b_2 = -b_1$

The possible unit according to this case has the form  $A = 1 + a_1I_1 - a_1I_2$ ,  $A^{-1} = 1 + b_1I_1 - b_1I_2$ .

$$A.A^{-1} = 1 \Rightarrow 1 + b_1I_1 - b_1I_2 + a_1I_1 + a_1b_1I_2 - a_1b_1I_1 - a_1I_2 - a_1b_1I_1 + a_1b_1I_2 = 1$$

$$\Rightarrow I_1(b_1 + a_1 - 2a_1b_1) + I_2(-a_1 - b_1 + 2a_1b_1) = 0, \text{ so that } b_1 + a_1 - 2a_1b_1 = 0 \Rightarrow b_1 = a_1(2b_1 - 1).$$

This means that  $2b_1 - 1 / b_1$  which is possible if and only if  $b_1 = 0$  or  $b_1 = 1$ .

If  $b_1 = 0$  then  $0 = a_1(-1) \Rightarrow a_1 = 0$ .

If  $b_1 = 1$  then  $1 = a_1(1) \Rightarrow a_1 = 1$ .

Thus, we get the following two units  $A_1 = 1, A_2 = 1 + I_1 - I_2$ , where  $A_2^{-1} = A_2$

Case 2: if  $a_0 = b_0 = 1$  and  $a_0 + a_1 + a_2 = b_0 + b_1 + b_2 = -1$ , hence  $a_2 = -2 - a_1$  and  $b_2 = -2 - b_1$ .

The possible units are  $A = 1 + a_1I_1 + (-2 - a_1)I_2, A^{-1} = B = 1 + b_1I_1 + (-2 - b_1)I_2$ .

$$A.A^{-1} = A.B = 1 \Rightarrow 1 + b_1I_1 - 2I_2 - b_1I_2 + a_1I_1 + a_1b_1I_2 - 2a_1I_1 - a_1I_2 - a_1b_1I_1 - 2I_2 - a_1I_2 - 2b_1I_1 - a_1b_1I_1 + 4I_2 + a_1b_1I_2 + 2a_1I_2 + 2b_1I_2 = 1,$$

$$\Rightarrow I_1(-b_1 - a_1 - 2a_1b_1) + I_2(a_1 + b_1 + 2a_1b_1) = 0, \text{ so that } a_1 + b_1 + 2a_1b_1 = 0 \Rightarrow b_1 = a_1(-2b_1 - 1).$$

This is possible if and only if  $a_1 = b_1 = 0$  or  $b_1 = -1$ , which implies that  $-1 = a_1(1) = a_1$ .

The possible unit is  $A_3 = 1 - I_1 + I_2$  with  $A_3^{-1} = A_3$ , and  $A_4 = 1 - 2I_2$  with  $A_4^{-1} = A_4$ .

Case 3: if  $a_0 = b_0 = -1$  and  $a_0 + a_1 + a_2 = b_0 + b_1 + b_2 = -1$ , then  $a_2 = -a_1$  and  $b_2 = -b_1$ .

By a similar argument to the case 1, we get:

$$A_5 = -1 \text{ or } A_5 = -1 + I_1 - I_2, \text{ where } A_6^{-1} = A_6.$$

Case 4: if  $a_0 = b_0 = -1$  and  $a_0 + a_1 + a_2 = b_0 + b_1 + b_2 = 1$ , then  $a_2 = 2 - a_1$  and  $b_2 = 2 - b_1$ .

By a similar argument to the case 1, we get:

$$A_5 = -1 \text{ or } A_5 = -1 + I_1 - I_2, \text{ where } A_6^{-1} = A_6$$

The possible units have the form:

$$A = -1 + a_1I_1 + (2 - a_1)I_2, B = A^{-1} = -1 + b_1I_1 + (2 - b_1)I_2.$$

$$A.B = 1 \Rightarrow 1 - b_1I_1 + (2 - a_1)I_2 - a_1I_1 + a_1b_1I_2 + I_1(2a_1 - a_1b_1) + (-2 + a_1)I_2 - 2b_1I_1 - a_1b_1I_1 + (4 - 2b_1 + a_1b_1)I_2 = 1,$$

$$\Rightarrow I_1(b_1 + a_1 - 2a_1b_1) + I_2(-b_1 - a_1 + 2a_1b_1) = 0 \Rightarrow b_1 + a_1 - 2a_1b_1 = 0 \Rightarrow b_1 = a_1(2b_1 - 1),$$

$$\Rightarrow 2b_1 - 1 / b_1 \text{ so that } a_1 + b_1 + 2a_1b_1 = 0 \Rightarrow b_1 = a_1(-2b_1 - 1).$$

Which is possible if and only if  $a_1 = b_1 = 0$  or  $b_1 = 1 = a_1$

Hence  $A_7 = -1 + I_1 + I_2, A_7^{-1} = A_7$  and  $A_8 = 1 + 2I_2$  and  $A_8^{-1} = A_8$ .

**Theorem .**

The group of units of  $Z_2(I)$  is isomorphic to  $Z_2 \times Z_2 \times Z_2$ .

Proof. According to the previous theorem we get 8 units. Each one has order 2. This implies that the group of units is isomorphic to  $Z_2 \times Z_2 \times Z_2$ .

### Theorem.

The group of units of the finite 2-cyclic refined neutrosophic ring  $(Z_2)_2(I)$  is isomorphic to  $Z_2$ .

Proof.

Let  $A = a_0 + a_1I_1 + a_2I_2$  be a unit in  $(Z_2)_2(I)$ , then  $A^{-1} = b_0 + b_1I_1 + b_2I_2$  such that  $a_0b_0 = 1$  and

$$(a_0 + a_1 + a_2)(b_0 + b_1 + b_2) = 1.$$

This means that  $a_0 = b_0 = 1$  and  $a_0 + a_1I_1 + a_2I_2 = b_0 + b_1I_1 + b_2I_2 = 1$ , thus  $a_2 = -a_1$ ,  $b_2 = -b_1$ .

$$A = 1 + a_1I_1 - a_2I_2, A^{-1} = 1 + b_1I_1 - b_2I_2.$$

$$A \cdot A^{-1} = 1 \Rightarrow 1 + b_1I_1 + a_1I_1 + a_1b_1I_2 - a_1b_1I_1 - a_1I_2 - a_1b_1I_1 + a_1b_1I_2 = 1.$$

Hence.

$$\Rightarrow I_1(a_1 + b_1 - 2a_1b_1) + I_2(-a_1 - b_1 + 2a_1b_1) = 0, \text{ so that } a_1 + b_1 - 2a_1b_1 = 0 \Rightarrow a_1 + b_1 = 0 \Rightarrow$$

$$a_1 = -b_1.$$

This means that.  $A = 1 + a_1I_1 - a_1I_2, A^{-1} = 1 - a_1I_1 + a_1I_2$ .

If  $a_1 = 0$ , then  $A_1 = A_1^{-1} = 1$ .

If  $a_1 = 1$ , then  $A_2 = 1 + I_1 - I_2, A_2^{-1} = 1 - I_1 + I_2 = 1 + I_1 - I_2 = A_2$ .

$A_2 \cdot A_2^{-1} = 1 \Rightarrow 1 - I_1 + I_2 + I_1 - I_2 + I_1 - I_2 + I_1 - I_2 = 1 \Rightarrow 2I_1 - 2I_2 = 0$  it is true.

**Remark 2.1.**  $R_n(I) \cong \underbrace{R \times R \times \dots R}_{n\text{-times}}$  in general.

### Theorem

The group of units of  $(Z_3)(I)$  is isomorphic to  $Z_2 \times Z_2 \times Z_2$ .

Proof.

If  $A = a_0 + a_1I_1 + a_2I_2 \in Z_2(I)$  is a unit, then  $B = A^{-1} = b_0 + b_1I_1 + b_2I_2$ , such that  $a_0b_0 = 1$  (I) and

$$(a_0 + a_1 + a_2)(b_0 + b_1 + b_2) = 1(II).$$

From equation (II), we get:

$$a_0 = b_0 = 1 \text{ or } a_0 = b_0 = 2$$

From equation (II), we get:

$$a_0 + a_1 + a_2 = b_0 + b_1 + b_2 = 1 \text{ or } a_0 + a_1 + a_2 = b_0 + b_1 + b_2 = 2.$$

We discuss the possible cases.

Case 1: if  $a_0 = b_0 = 1$  and  $a_0 + a_1 + a_2 = b_0 + b_1 + b_2 = 1$ , we get  $a_1 + a_2 = b_1 + b_2 = 0$ , hence

$$a_2 = -a_1, b_2 = -b_1$$

The possible units have the form  $A = 1 + a_1I_1 - a_1I_2, A^{-1} = B = 1 + b_1I_1 - b_1I_2$ .

$$\Rightarrow AA^{-1} = 1 \Rightarrow I_1(b_1 + a_1 - 2a_1b_1) + I_2(-a_1 - b_1 + 2a_1b_1) = 0, \text{ so that } a_1 + b_1 - 2a_1b_1 = 0$$

$$\Rightarrow a_1 + b_1 = 2a_1b_1 (*).$$

Equation (\*) has the following solutions:

$$a_1 = b_1 = 0, a_1 = b_1 = 1, \text{ hence}$$

$$A_1 = 1, A_2 = 1 + I_1 - I_2, \text{ where } A_2^{-1} = A_2.$$

Case 2: if  $a_0 = b_0 = 1$  and  $a_0 + a_1 + a_2 = b_0 + b_1 + b_2 = 2 \Rightarrow a_1 + a_2 = b_1 + b_2 = 1$ , so that

$$A = 1 + a_1I_1 + (1 - a_1)I_2, A^{-1} = B = 1 + b_1I_1 + (1 - b_1)I_2.$$

$$A.A^{-1} = 1 \Rightarrow 1 + a_1I_1 + (1 - a_1)I_2 + a_1b_1I_2 + I_2(b_1 - b_1a_1) + (1 - b_1)I_2 + (a_1 - a_1b_1)I_1 + I_2(1 + a_1b_1 - a_1 - b_1) = 1,$$

$$\Rightarrow I_1(a_1 + b_1 + b_1 - b_1a_1 + a_1 - a_1b_1) + I_2(1 - a_1 + a_1b_1 + 1 - b_1 + 1 + a_1b_1 - a_1 - b_1) = 0$$

$$\Rightarrow \begin{cases} 2a_1 + 2b_1 - 2a_1b_1 = 0 \\ -2a_1 - 2b_1 + 2a_1b_1 = 0 \end{cases}$$

$\Rightarrow 2(a_1 + b_1 - a_1b_1) = 0$ , so  $a_1 + b_1 - a_1b_1$  is a zero divisor, which is possible if and only if  $a_1 + b_1 - a_1b_1 = 0 \Rightarrow a_1 + b_1 = a_1b_1$ .

The possible solutions are  $(a_1 = b_1 = 0), (a_1 = b_1 = 2)$

$$A_3 = 1 + I_2, A_3^{-1} = A_3, A_4 = 1 + 2I_1 - I_2 \text{ and } A_4^{-1} = A_4.$$

Case 3: if  $a_0 = b_0 = 2$  and  $a_0 + a_1 + a_2 = b_0 + b_1 + b_2 = 1$ , then  $a_2 = 2 - a_1$  and  $b_2 = 2 - b_1$ .

The possible units have the form:

$$A = 2a_1I_1 + (2 - a_1)I_2, A^{-1} = 2 + b_1I_1 + (2 - b_1)I_2.$$

$$A.A^{-1} = 1 + 2b_1I_1 + (4 - 2b_1)I_2 + 2a_1I_1 + a_1b_1I_2 + (2a_1 - a_1b_1)I_1 + (4 - 2a_1)I_2 + (2b_1 - a_1b_1)I_2 + (4 - 2b_1 - 2a_1 + a_1b_1)I_2 = 1,$$

$$\Rightarrow I_1(2b_1 + 2a_1 + 2a_1 - a_1b_1 + 2b_1 - a_1b_1) + I_2(4 - 2b_1 + a_1b_1 + 4 - 2a_1 + 4 - 2b_1 - 2a_1 + a_1b_1) = 0,$$

$$\Rightarrow \begin{cases} a_1 + b_1 - 2a_1b_1 = 0 \\ -a_1 - b_1 + 2a_1b_1 = 0 \end{cases} \Rightarrow a_1 + b_1 = 2a_1b_1,$$

The possible solutions are ( $a_1 = b_1 = 0$  or  $a_1 = b_1 = 1$ ).

Thus the corresponding units are:

$$A_5 = 2 + 2I_2, A_6 = 2 + I_1 + I_2 \text{ where } A_5^{-1} = A_5, A_6^{-1} = A_6.$$

Case 4: if  $a_0 = b_0 = 2$  and  $a_0 + a_1 + a_2 = b_0 + b_1 + b_2 = 2$ , we get  $a_1 + a_2 = b_1 + b_2 = 0$ , hence  $a_2 = -a_1, b_2 = -b_1$ .

The possible units are:

$$A = 2 + a_1I_1 - a_1I_2, A^{-1} = 2 + b_1I_1 - b_1I_2.$$

$$A.A^{-1} = 1 \Rightarrow 1 + 2b_1I_1 - 2b_1I_2 + 2a_1I_1 + a_1b_1I_2 - a_1b_1I_1 - 2a_1I_2 - a_1b_1I_1 + a_1b_1I_2 = 1$$

$$\Rightarrow I_1(2b_1 + 2a_1 - 2a_1b_1) + I_2(-2b_1 - 2a_1 + 2a_1b_1) = 0$$

$\Rightarrow \{2b_1 + 2a_1 - 2a_1b_1 = 0$  so that  $2(b_1 + a_1 - a_1b_1) = 0$ , hence  $b_1 + a_1 = a_1b_1$  is a zero divisor, thus

$$b_1 + a_1 - a_1b_1 = 0, \text{ this means that } b_1 + a_1 = a_1b_1.$$

The possible solutions are ( $a_1 = b_1 = 0$  or  $a_1 = b_1 = 2$ ).

Thus the corresponding units are:

$$A_7 = 2, A_8 = 2 + 2I_1 \text{ where } A_8^{-1} = A_8.$$

Since the order of any unit is 2, we get the proof.

### Theorem

The 2-cyclic refined neutrosophic ring of real numbers  $R_2(I) = \{t_0 + t_1I_1 + t_2I_2; t_i \in R\}$  has infinite group of units.

Proof.

Let  $A = a_0 + a_1I_1 + a_2I_2 \in R_2(I)$  is a unit, then  $B = A^{-1} = b_0 + b_1I_1 + b_2I_2; a_0b_0 = 1(I)$ , and

$$(a_0 + a_1 + a_2)(b_0 + b_1 + b_2) = 1(II).$$

From equation (I), we get:

$$b_0 = \frac{1}{a_0}; a_0 \neq 0.$$

From equation (II), we get:

$$b_0 + b_1 + b_2 = \frac{1}{a_0 + a_1 + a_2}, \text{ thus implies that } b_0 + b_1 + b_2 \neq 0, a_0 + a_1 + a_2 \neq 0$$

$$\text{and } \frac{1}{a_0} + b_1 + b_2 = \frac{1}{a_0 + a_1 + a_2}, \text{ thus } b_2 = \frac{1}{a_0 + a_1 + a_2} - \frac{1}{a_0} - b_1 = \frac{-a_1 - a_2}{a_0(a_0 + a_1 + a_2)} - b_1.$$

$$\text{On the other hand, we have } A \cdot A^{-1} = 1, \text{ thus } (a_0 + a_1 I_1 + a_2 I_2) \left( \frac{1}{a_0} + b_1 I_1 + \left[ \frac{-a_1 - a_2}{a_0(a_0 + a_1 + a_2)} - b_1 \right] I_2 \right) = 1$$

$$\Rightarrow 1 + a_0 b_1 I_1 + I_2 \left( \frac{-a_1 - a_2}{a_0(a_0 + a_1 + a_2)} - a_0 b_1 \right) + \frac{a_1}{a_0} I_1 + a_1 b_1 I_2 + I_1 \left( \frac{-a_1^2 - a_1 a_2}{a_0(a_0 + a_1 + a_2)} - a_1 b_1 \right) + \frac{a_2}{a_0} I_2 + a_2 b_1 I_1 + I_2 \left( \frac{-a_1 a_2 - a_2^2}{a_0(a_0 + a_1 + a_2)} - a_2 b_2 \right) = 1,$$

$$\Rightarrow I_1 \left( a_0 b_1 + \frac{a_1}{a_0} - a_1 b_1 \frac{-a_1^2 - a_1 a_2}{a_0(a_0 + a_1 + a_2)} + a_2 b_1 \right) + I_2 \left( \frac{-a_1 - a_2}{a_0(a_0 + a_1 + a_2)} - a_0 b_1 + \frac{a_2}{a_0} - a_2 b_2 + \frac{-a_1 a_2 - a_2^2}{a_0(a_0 + a_1 + a_2)} \right) = 0,$$

We get the following equations:

$$(*) a_0 b_1 + \frac{a_1}{a_0} - a_1 b_1 \frac{-a_1^2 - a_1 a_2}{a_0(a_0 + a_1 + a_2)} + a_2 b_1 = 0$$

$$(**) \frac{-a_1 - a_2}{a_0(a_0 + a_1 + a_2)} - a_0 b_1 + \frac{a_2}{a_0} - a_2 b_2 + \frac{-a_1 a_2 - a_2^2}{a_0(a_0 + a_1 + a_2)} = 0$$

We simplify (\*) as follows:

$$b_1(a_0 - a_1 + a_2) + \frac{a_1(a_0 + a_1 + a_2)}{a_0(a_0 + a_1 + a_2)} + \frac{-a_1^2 - a_1 a_2}{a_0(a_0 + a_1 + a_2)} = 0$$

Hence

$$b_1(a_0 - a_1 + a_2) + \frac{a_0 a_1}{a_0(a_0 + a_1 + a_2)} = 0$$

So

$$b_1 = \frac{-a_1}{(a_0 - a_1 + a_2)(a_0 + a_1 + a_2)}$$

By solving (\*\*), we get the same solution of (\*).

This implies that the units have the form  $A = a_0 + a_1 I_1 + a_2 I_2$ , with the condition  $a_0 + a_1 + a_2 \neq 0, a_0 \neq 0, a_0 - a_1 + a_2 \neq 0$ . The converse of A is  $A^{-1} = \frac{1}{a_0} + I_1 \left[ \frac{-a_1}{(a_0 - a_1 + a_2)(a_0 + a_1 + a_2)} \right] + I_2 \left[ \frac{-a_1 - a_2}{a_0(a_0 + a_1 + a_2)} + \frac{a_1}{(a_0 + a_1 + a_2)(a_0 - a_1 + a_2)} \right]$ .



**Example:**

Consider the following 2-cyclic refined neutrosophic number  $A = 1 + 2I_1 - I_2$ . It is invertible, that is because

$$a_0 + a_1 + a_2 = 2 \neq 0, a_0 = 1 \neq 0, a_0 - a_1 + a_2 = -2 \neq 0.$$

The converse of A is  $A^{-1} = 1 + I_1 \left[ \frac{-2}{(-2)(2)} \right] + I_2 \left[ \frac{-1}{1(2)} + \frac{2}{(2)(-2)} \right] = 1 + \frac{1}{2}I_1 - I_2$ .

**Example:**

Now, we can find the inverse of any square 2-cyclic refined real neutrosophic matrix, as the following example shows.

Consider the following 2-cyclic refined neutrosophic matrix  $M = \begin{pmatrix} 1 & -2I_2 \\ I_1 & 1 - I_2 \end{pmatrix}$ , we have

$\det(M) = 1 + 2I_1 - I_2$ , which is invertible according to the previous example. This implies that M is invertible.

$$M^{-1} = \frac{1}{\det(M)} (\text{adj}(M)) = \left(1 + \frac{1}{2}I_1 - I_2\right) \begin{pmatrix} 1 - I_2 & 2I_2 \\ -I_1 & 1 \end{pmatrix}.$$

**4. Conclusion**

In this paper, we have provided a novel solution of the group of units problem in some 2-cyclic refined neutrosophic rings, such as 2-cyclic refined neutrosophic ring of integers  $Z_2(I)$ , 2-cyclic refined neutrosophic ring of integers modulo 2 and 3  $(Z_2)_2(I)$ ,  $(Z_3)_2(I)$  and 2-cyclic refined neutrosophic ring of real numbers  $R_2(I)$ . Also, we have illustrated many examples especially for the invertibility of any square 2-cyclic refined neutrosophic real matrix.

According to this research, many open questions are coming to light.

**Open problem 1:** If the ring R has no zero divisors, then is the group of units of its corresponding 2-cyclic refined neutrosophic ring  $R_2(I)$  isomorphic to  $U(R) \times U(R) \times U(R)$ . (Remark that the answer is yes if R is the ring of integers, or the ring of integers modulo 3).

**Open problem 2:** Is there a ring homomorphism between the 2-cyclic refined neutrosophic ring  $R_2(I)$  and the direct product  $R \times R \times R$ .

**Open problem 3:** Is the group of units of the 2-cyclic refined neutrosophic ring or real numbers isomorphic to  $R^* \times R^* \times R^*$ .

**Open problem 4:** Determine the structure of the group of units of previous rings in the case of  $n=3$ ,  $n=4$ ,  $n=5$ .

**Open Problem 5:** Answer open problems 1, 2, 3 for the case n is greater than 2.

**References**

- [1]. Edalatpanah. S.A., Systems of Neutrosophic Linear Equations, NSS. 2020.
- [2]. M. Abobala. *n*-Cyclic Refined Neutrosophic Algebraic Systems Of Sub-Indeterminacies, An Application To Rings and Modules, International Journal of Neutrosophic Science, Vol. 12, 2020. pp. 81-95 .
- [3]. E. Adeleke. A. Agboola. , and F. Smarandache. Refined Neutrosophic Rings II. IJNS, 2020.
- [4]. Ibrahim, M., and Abobala, M., An Introduction To Refined Neutrosophic Number Theory, Neutrosophic Sets and Systems, Vol. 45, 2021.
- [5]. E. Adeleke. A. Agboola. , and F. Smarandache. Refined Neutrosophic Rings I. IJNS, 2020.
- [6]. V. Kandasamy and F. Smarandache. Some Neutrosophic Algebraic Structures and Neutrosophic N-Algebraic Structures, Hexis, Phonex, Arizona, 2006.
- [7]. A. Hatip and N.Olgun. On Refined Neutrosophic R-Module. IJNS, 2020.
- [8]. M. Ibrahim. A. Agboola. B.Badmus and S. Akinleye. On refined Neutrosophic Vector Spaces . IJNS, 2020.
- [9]. Abobala, M., On The Characterization of Maximal and Minimal Ideals In Several Neutrosophic Rings, Neutrosophic Sets and Systems, Vol. 45, 2021.
- [10]. Abobala, M., and Hatip, A., An Algebraic Approach To Neutrosophic Euclidean Geometry, Neutrosophic Sets and Systems, Vol. 43, 2021.
- [11]. Bal, M., Ahmad, K., Hajjari, A., Ali, R., The Structure Of Imperfect Triplets In Several Refined Neutrosophic Rings Journal of Neutrosophic and Fuzzy Systems, 2022.
- [12]. Ceven, Y., and Tekin, S., Some Properties of Neutrosophic Integers, Kırklareli University Journal of Engineering and Science, Vol. 6, pp.50-59, 2020.
- [13]. Abobala, M., A Study Of Nil Ideals and Kothe's Conjecture In Neutrosophic Rings, International Journal of Mathematics and Mathematical Sciences, hindawi, 2021

- [14]. Abobala, M., On Some Algebraic Properties of n-Refined Neutrosophic Elements and n-Refined Neutrosophic Linear Equations, *Mathematical Problems in Engineering*, Hindawi, 2021