



NEUTROSOPHIC g^* -CLOSED SETS IN NEUTROSOPHIC TOPOLOGICAL SPACES

Wadei Faris AL-Omeri ¹ *

¹Department of Mathematics, Faculty of Science and Information Technology,
Jadara University, Irbid, Jordan.

Emails: wadeimoon1@hotmail.com

Abstract

A neutrosophic set is a mathematical approach that helps with challenges involving data that is indeterminate, imprecise, or inconsistent. The goal of this manuscript is to present the notion of neutrosophic g^* -closed sets and neutrosophic g^* -open sets. In this situation, we prove various neutrosophic generalized theorems. The findings support previous methodologies in the literature and are backed up by various examples and an application.

Keywords: Neutrosophic g^* continuity, Neutrosophic Set, Neutrosophic g^* closed sets, Neutrosophic Topology, Neutrosophic g^* continuity mapping, Generalized Neutrosophic Set, Neutrosophic continuity mapping.

1 Introduction and Preliminaries

Generalization topological spaces is a classical subject which is a type of classical topological spaces. The concept of fuzzy sets was introduced by Zadeh in his classical paper.³ In 1968, Chang¹⁹ used fuzzy sets to introduce the notion of fuzzy topological spaces. Çoker^{2,18} defined and introduced the intuitionistic fuzzy topological spaces using intuitionistic fuzzy sets Atanassov's²² (IFS for short) on a universe X . Neutrosophic system was defined by Smarandache In various recent papers, has laid the foundation for a whole family of new mathematical theories generalizing both their fuzzy and classical counterparts, and have wide range of real applications for the fields of Medicine, Information Systems, decision making, Applied Mathematic, Computer Science. F. Smarandache and A. Al Shumrani obtained the concept of neutrosophic topology on the non-general and standard interval.^{20,21} Several authors was extended this principle with many applications (see^{10-14,24-26}). In this paper we study and introduce about generalized neutrosophic sets. g^* -closed sets in neutrosophic topological space. We also discussed with details about properties and relationships with other classes of early defined forms. We also introduced application of neutrosophic g^* -closed sets namely neutrosophic $T_{\frac{1}{2}}^*$ space and ${}^*T_{\frac{1}{2}}$ space. Several interesting characterizations and properties are also discussed.

For non-empty fixed set \mathcal{R} and each element $r \in \mathcal{R}$ to the set \tilde{R} . A neutrosophic set¹⁶ (NS for short) \tilde{R} is an object having the form $\tilde{R} = \{\langle r, \mu_{\tilde{R}}(r), \sigma_{\tilde{R}}(r), \gamma_{\tilde{R}}(r) \rangle : r \in \mathcal{R}\}$, where $\mu_{\tilde{R}}(k)$ represent the degree of membership function, $\sigma_{\tilde{R}}(k)$ represent the degree of indeterminacy, and $\gamma_{\tilde{R}}(k)$ represent the degree of non-membership. Neutrosophic sets in \mathcal{S} will be denoted by $\tilde{R}, \mathcal{W}, B, G$, etc., and although subsets of \mathcal{R} will be denoted by \tilde{R}, \tilde{B}, T , etc.

A neutrosophic set $\tilde{R} = \{\langle k, \mu_{\tilde{R}}(k), \sigma_{\tilde{R}}(k), \gamma_{\tilde{R}}(k) \rangle : k \in \mathcal{R}\}$ can be identified to an ordered triple $\langle \mu_{\tilde{R}}(k), \sigma_{\tilde{R}}(k), \gamma_{\tilde{R}}(k) \rangle$ in $]0^-, 1^+]$ on \mathcal{R} .

2 Historical Background

Definition 2.1. ¹⁶ Let \mathcal{Z} be a non-empty set. A neutrosophic set (NS for short) \tilde{R} is an object having the form $\tilde{R} = \{\langle r, \mu_{\tilde{R}}(r), \sigma_{\tilde{R}}(r), \gamma_{\tilde{R}}(r) \rangle : r \in \mathcal{Z}\}$, where $\gamma_{\tilde{R}}(r)$, $\sigma_{\tilde{R}}(r)$, $\mu_{\tilde{R}}(r)$, and the degree of non-membership (namely $\gamma_{\tilde{R}}(r)$), the degree of indeterminacy (namely $\sigma_{\tilde{R}}(r)$), and the degree of membership function (namely $\mu_{\tilde{R}}(r)$), of each element $r \in \mathcal{Z}$ to the set \tilde{R} .

Since our main purpose is to construct the tools for developing neutrosophic set and neutrosophic topology, we must introduce the Neutrosophic sets (NSS) 0_N and 1_N ¹⁶ in \mathcal{Z} are introduced as follows:

$1 - 0_N$ can be defined as four types:

1. $0_N = \{\langle r, 0, 0, 1 \rangle : r \in \mathcal{Z}\}$,
2. $0_N = \{\langle r, 0, 1, 1 \rangle : r \in \mathcal{Z}\}$,
3. $0_N = \{\langle r, 0, 1, 0 \rangle : r \in \mathcal{Z}\}$,
4. $0_N = \{\langle r, 0, 0, 0 \rangle : r \in \mathcal{Z}\}$.

$2- 1_N$ can be defined as four types:

1. $1_N = \{\langle r, 1, 0, 0 \rangle : r \in \mathcal{Z}\}$,
2. $1_N = \{\langle r, 1, 0, 1 \rangle : r \in \mathcal{Z}\}$,
3. $1_N = \{\langle r, 1, 1, 0 \rangle : r \in \mathcal{Z}\}$,
4. $1_N = \{\langle r, 1, 1, 1 \rangle : r \in \mathcal{Z}\}$.

Definition 2.2. ¹⁶ Let X be a non-empty set, and $GNSS$ H and K in the form $H = \{r, \mu_H(r), \sigma_H(r), \gamma_H(r)\}$, $K = \{r, \mu_K(r), \sigma_K(r), \gamma_K(r)\}$, then we may consider two possible definitions for subsets ($H \subseteq K$) ($H \subseteq K$) may be defined as

1. Type 1: $H \subseteq K \Leftrightarrow \mu_H(r) \leq \mu_K(r), \sigma_H(r) \leq \sigma_K(r)$, and $\gamma_H(r) \geq \gamma_K(r)$ or
2. Type 1: $H \subseteq K \Leftrightarrow \mu_H(r) \leq \mu_K(r), \sigma_H(r) \geq \sigma_K(r)$, and $\gamma_H(r) \geq \gamma_K(r)$.

Definition 2.3. ¹⁶ Let $\{A_j : j \in J\}$ be a arbitrary family of NSS in \mathcal{Z} , then

1. $\cap A_j$ may be defined as two types:
 - Type 1: $\cap A_j = \langle r, \bigwedge_{j \in J} \mu_{A_j}(r), \bigwedge_{j \in J} \sigma_{A_j}(r), \bigvee_{j \in J} \gamma_{A_j}(r) \rangle$
 - Type 2: $\cap A_j = \langle r, \bigwedge_{j \in J} \mu_{A_j}(r), \bigvee_{j \in J} \sigma_{A_j}(r), \bigvee_{j \in J} \gamma_{A_j}(r) \rangle$.
2. $\cup A_j$ may be defined as two types:
 - Type 1: $\cup A_j = \langle r, \bigvee_{j \in J} \mu_{A_j}(r), \bigvee_{j \in J} \sigma_{A_j}(r), \bigwedge_{j \in J} \gamma_{A_j}(r) \rangle$
 - Type 2: $\cup A_j = \langle r, \bigvee_{j \in J} \mu_{A_j}(r), \bigwedge_{j \in J} \sigma_{A_j}(r), \bigwedge_{j \in J} \gamma_{A_j}(r) \rangle$

Definition 2.4. Let $\tilde{R} = \langle \mu_{\tilde{R}}(r), \sigma_{\tilde{R}}(r), \gamma_{\tilde{R}}(r) \rangle$ be an NS on \mathcal{Z} .² The complement of the set $\tilde{R}(C(\tilde{R}))$, for short) may be defined as follows:

1. $C(\tilde{R}) = \{\langle r, 1 - \mu_{\tilde{R}}(r), 1 - \gamma_{\tilde{R}}(r) \rangle : r \in \mathcal{Z}\}$,
2. $C(\tilde{R}) = \{\langle r, \gamma_{\tilde{R}}(r), \sigma_{\tilde{R}}(r), \mu_{\tilde{R}}(r) \rangle : r \in \mathcal{Z}\}$,
3. $C(\tilde{R}) = \{\langle r, \gamma_{\tilde{R}}(r), 1 - \sigma_{\tilde{R}}(r), \mu_{\tilde{R}}(r) \rangle : r \in \mathcal{Z}\}$.

Definition 2.5.¹ A neutrosophic topology (NT for short) and a non empty set \mathcal{Z} is a family \mathcal{T} of neutrosophic subsets of \mathcal{Z} satisfying the following axioms

1. $0_N, 1_N \in \mathcal{T}$
2. $H_1 \cap H_2 \in \mathcal{T}$ for any $H_1, H_2 \in \mathcal{T}$
3. $\cup H_i \in \mathcal{T}, \forall \{H_i | j \in J\} \subseteq \mathcal{T}$.

The pair $(\mathcal{Z}, \mathcal{T})$ is called a neutrosophic topological space (NTS for short).

Definition 2.6.¹⁶ Let \tilde{R} be an NS and $(\mathcal{Z}, \mathcal{T})$ an NT where $\tilde{R} = \{r, \mu_{\tilde{R}}(r), \sigma_{\tilde{R}}(r), \gamma_{\tilde{R}}(r)\}$. Then,

1. $NCl(\tilde{R}) = \cap \{H : H \text{ is an NCS in } \mathcal{Z} \text{ and } \tilde{R} \subseteq H\}$
2. $NInt(\tilde{R}) = \cup \{W : W \text{ is an NOS in } \mathcal{Z} \text{ and } W \subseteq \tilde{R}\}$

It can be also shown that $NCl(\tilde{R})$ is an NCS and $NInt(\tilde{R})$ is an NOS in \mathcal{Z} . We have

1. \tilde{R} is in \mathcal{Z} iff $NCl(\tilde{R})$.
2. \tilde{R} is an NCS in \mathcal{Z} iff $NInt(\tilde{R}) = \tilde{R}$.

Definition 2.7.¹⁶ Let $\tilde{R} = \{\mu_{\tilde{R}}(r), \sigma_{\tilde{R}}(r), \gamma_{\tilde{R}}(r)\}$ be a neutrosophic open sets and $B = \{\mu_B(r), \sigma_B(r), \gamma_B(r)\}$ a neutrosophic set on a neutrosophic topological space $(\mathcal{Z}, \mathcal{T})$. Then

1. \tilde{R} is called neutrosophic regular open iff $\tilde{R} = NInt(NCl(\tilde{R}))$.
2. The complement of neutrosophic regular open is neutrosophic regular closed.

Definition 2.8.⁹ Let \tilde{R} be an NS and $(\mathcal{Z}, \mathcal{T})$ an NT . Then

1. Neutrosophic regular-open set ($NROS$) if $\tilde{R} = NInt(NCl(\tilde{R}))$,
2. Neutrosophic regular-closed set ($NROS$) if $\tilde{R} = NCl(NInt(\tilde{R}))$,
3. Neutrosophic semi-open set ($NSOS$) if $\tilde{R} \subseteq NCl(NInt(\tilde{R}))$,
4. Neutrosophic semi-closed set ($NSCS$) if $NInt(NCl(\tilde{R})) \subseteq \tilde{R}$,
5. Neutrosophic pre-open set ($NPOS$) if $\tilde{R} \subseteq NInt(NCl(\tilde{R}))$,
6. Neutrosophic pre-closed set ($NPCS$) if $NCl(NInt(\tilde{R})) \subseteq \tilde{R}$,
7. Neutrosophic α -open set ($N\alpha OS$) if $\tilde{R} \subseteq NInt(NCl(NInt(\tilde{R})))$,
8. Neutrosophic α -closed set ($N\alpha CS$) if $NCl(NInt(NCl(\tilde{R}))) \subseteq \tilde{R}$.

Definition 2.9.⁷

1. Neutrosophic β -closed set ($N\beta CS$) if $NInt(NCl(NInt(\tilde{R}))) \subseteq \tilde{R}$,
2. Neutrosophic β -open set ($N\beta OS$) if $\tilde{R} \subseteq NCl(NInt(NCl(\tilde{R})))$,

Definition 2.10.¹⁷ A (NS) \tilde{R} of an NTs (\mathcal{Z}, Γ) is called generalized neutrosophic closed sets ($GNCs$ in short) if $NCl(\tilde{R}) \subseteq \tilde{B}$ whenever $\tilde{R} \subset \tilde{B}$ and \tilde{B} is neutrosophic closed sets in \mathcal{Z} .


Definition 2.11.⁵ A neutrosophic set \tilde{R} of an NTS (\mathcal{Z}, Γ) is neutrosophic generalized pre-closed set ($NGPCS$ in short) if $NPCl(P) \subseteq G$ whenever $P \subseteq G$ and G is an NOS in \mathcal{Z} .

Definition 2.12. ⁸ Consider a neutrosophic set \tilde{R} in $NTS(\mathcal{Z}, \Gamma)$. Then it is neutrosophic generalized β -closed set ($NG\beta CS$ in short) if $N\beta Cl(P) \subseteq G$ whenever $P \subseteq G$ and G is an NOS .

Definition 2.13. ⁶ A neutrosophic set \tilde{R} of an $NTS(\mathcal{Z}, \Gamma)$ is neutrosophic generalized semi-closed set ($NGSCS$ in short) if $NScl(P) \subseteq G$ whenever $P \subseteq G$ and G is an NOS in \mathcal{Z} .

Definition 2.14. ⁴ Consider a neutrosophic set \tilde{R} in $NTS(\mathcal{Z}, \Gamma)$. Then it is neutrosophic α -generalized closed set ($N\alpha GCS$ in short) if $N\alpha Cl(P) \subseteq G$ whenever $P \subseteq G$ and G is an NOS .

An $NS \tilde{R}$ is called a neutrosophic generalized open set, neutrosophic generalized pre-open set, neutrosophic α -generalized open set, neutrosophic generalized semi-open set, neutrosophic β -generalized open set ($NGOS, NGPOS, N\alpha GOS, NGSOS, NG\beta OS$ respectively), if the complement of \tilde{R}^c is a $NGCS, NGPCS, N\alpha GCS, NGSCS, NG\beta CS$ respectively. The following implications²³ are true:



In this diagram by $A \longrightarrow B$ means A implies B but not conversely and $A \perp B$ means A & B are independent.

Diagram 1

Definition 2.15. ⁹ Let (\mathcal{Z}, Γ) be a neutrosophic topological space and $x_{r,t,s}$ be a neutrosophic point in \mathcal{Z} . A neutrosophic set S of \mathcal{Z} is called a neutrosophic neighbourhood if there exists a neutrosophic open set $x_{r,t,s}$ in \mathcal{Z} such that $p_\epsilon \in x_{r,t,s} \subseteq S$.

Definition 2.16. ²⁷ A neutrosophic point $x_{r,t,s}$ is said to be q -coincident with a neutrosophic set G , denoted by $x_{r,t,s}qG$, if and only if there exists $x_{r,t,s} \not\subseteq G^c$. A neutrosophic set H is said to be neutrosophic quasi-coincident (neutrosophic q -coincident, for short) with G , denoted by HqG if and only if $H \not\subseteq G^c$.

Definition 2.17. ²⁷ A neutrosophic point (NP in short), $x_{r,t,s}$ is said to be a neutrosophic cluster point of a neutrosophic set (Ns in short) G if HqG for each neutrosophic open q -neighborhood H of $x_{r,t,s}$. The union of all neutrosophic cluster points of G is called the neutrosophic closure of G and denoted by $NCl(G)$.

Definition 2.18. ¹⁵ A neutrosophic set (Ns in short), (\mathcal{Z}, Γ) is said to be neutrosophic- $T_{\frac{1}{2}}$ ($NT_{\frac{1}{2}}$ in short) space if every $GNCs$ in \mathcal{Z} is an NCs in \mathcal{Z} .

3 Neutrosophic g^* Closed Sets

In this section we introduce neutrosophic g^* -closed sets in neutrosophic topological spaces and discuss some of their properties.

Definition 3.1. A neutrosophic point (NP in short) $x_{r,t,s}$ is said to be a neutrosophic θ -cluster point of a neutrosophic set (Ns in short) \tilde{R} if and only if $Cl(H)q\tilde{R}$ for each q -neighborhood H . The set of all neutrosophic θ -cluster points of \tilde{R} is said to be neutrosophic θ -closure of \tilde{R} and is denoted by $Cl_\theta(\tilde{R})$. An $Ns \tilde{R}$ will be called a neutrosophic θ -closed ($N\theta CS$ for short) if and only if $\tilde{R} = Cl_\theta(\tilde{R})$. The complement of θCS is said to be neutrosophic θ -open set ($N\theta OS$ for short).

Definition 3.2. A neutrosophic set H in (\mathcal{Z}, Γ) is said to be a neutrosophic g^* -closed set (NG^*CS in short) if $Cl(H) \subseteq W$ whenever $H \subseteq W$ and W is an $NGOS$ in (\mathcal{Z}, Γ) . The family of all G^*CSs of an $NTS(\mathcal{Z}, \Gamma)$ is denoted by $NG^*C(\mathcal{Z})$.

Example 3.3. Let $X = \{u, v\}$ $\Gamma_N = \{0_N, 1_N, F\}$ is a N_Γ on \mathcal{Z} . where $F = \langle x, (\frac{3}{10}, \frac{4}{10}, \frac{4}{10}), (\frac{4}{10}, \frac{3}{10}, \frac{1}{10}) \rangle$. Then the Neutrosophic set $H = \langle x, (\frac{7}{10}, \frac{6}{10}, \frac{6}{10}), (\frac{5}{10}, \frac{5}{10}, \frac{5}{10}) \rangle$ is an NG^*CS in (\mathcal{Z}, Γ) .

Example 3.4. Let $X = \{u, v, w\}$ and $A = \langle (\frac{4}{10}, \frac{5}{10}, \frac{4}{10}), (\frac{5}{10}, \frac{5}{10}, \frac{5}{10}), (\frac{4}{10}, \frac{5}{10}, \frac{4}{10}) \rangle$, $B = \langle (\frac{7}{10}, \frac{6}{10}, \frac{5}{10}), (\frac{3}{10}, \frac{4}{10}, \frac{5}{10}), (\frac{3}{10}, \frac{4}{10}, \frac{4}{10}) \rangle$, and $C = \langle (\frac{5}{10}, \frac{5}{10}, \frac{5}{10}), (\frac{5}{10}, \frac{5}{10}, \frac{5}{10}), (\frac{5}{10}, \frac{5}{10}, \frac{5}{10}) \rangle$. Then $\tau = \{0_N, 1_N, A, B\}$ is a neutrosophic topology on X . Then the Neutrosophic set $C = \langle (\frac{6}{10}, \frac{4}{10}, \frac{5}{10}), (\frac{5}{10}, \frac{6}{10}, \frac{5}{10}), (\frac{5}{10}, \frac{6}{10}, \frac{4}{10}) \rangle$ is an NG^*CS in (\mathcal{Z}, Γ) .

Example 3.5. $X = \{u, v\}$ $\Gamma_N = \{0_N, 1_N, F\}$ is a N_Γ on \mathcal{Z} . where $F = \langle x, (\frac{6}{10}, \frac{5}{10}, \frac{3}{10}), (\frac{8}{10}, \frac{5}{10}, \frac{2}{10}) \rangle$. Then the Neutrosophic set $H = \langle x, (\frac{6}{10}, \frac{5}{10}, \frac{4}{10}), (\frac{3}{10}, \frac{5}{10}, \frac{7}{10}) \rangle$ is not an NG^*CS in (\mathcal{Z}, Γ) .

Theorem 3.6. Every NCS is a NG^*CS but not conversely.

Proof. Let H be a NCS in (\mathcal{Z}, Γ) . Then $Cl(H) = H$. Let $H \subseteq W$ and W is an $NGOS$ in (\mathcal{Z}, Γ) . Therefore $Cl(H) = H \subseteq W$. Thus H is an NG^*CS in (\mathcal{Z}, Γ) . □

Example 3.7. Let $X = \{u, v\}$ $\Gamma_N = \{0_N, 1_N, F\}$ is a N_Γ on \mathcal{Z} , where $F = \langle x, (\frac{4}{10}, \frac{2}{10}, \frac{4}{10}), (\frac{6}{10}, \frac{7}{10}, \frac{5}{10}) \rangle$. Then the Neutrosophic set $H = \langle x, (\frac{6}{10}, \frac{7}{10}, \frac{6}{10}), (\frac{1}{10}, \frac{1}{10}, \frac{1}{10}) \rangle$ is an NG^*CS in (\mathcal{Z}, Γ) but not an NCS in Γ .

Theorem 3.8. Every NG^*CS in neutrosophic topological space (\mathcal{Z}, Γ) is an $NGCS$ but not conversely.

Proof. Let H be an NG^*CSCS in (\mathcal{Z}, Γ) . Let $H \subseteq W$ and W is a NOS in (\mathcal{Z}, Γ) . Since every NOS is $NGOS$ and since H is an NG^*CS in \mathcal{Z} . Therefore $Cl(H) \subseteq W$ whenever $H \subseteq W$, W is an NOS in \mathcal{Z} . Thus H is an $NGCS$ in \mathcal{Z} . □

Example 3.9. Let $X = \{u, v\}$ $\Gamma_N = \{0_N, 1_N, F\}$ is a N_Γ on \mathcal{Z} . where $F = \langle x, (\frac{3}{10}, \frac{7}{10}, \frac{4}{10}), (\frac{6}{10}, \frac{3}{10}, \frac{5}{10}) \rangle$. Then the $NS H = \langle x, (\frac{3}{10}, \frac{8}{10}, \frac{6}{10}), (\frac{7}{10}, \frac{2}{10}, \frac{1}{10}) \rangle$ is an $NGCS$ but not an NG^*CS in Γ .

Theorem 3.10. Every $NG\theta CS$ in neutrosophic topological space (\mathcal{Z}, Γ) is an $N\alpha GCS$ but not conversely.

Proof. It is obvious because every $N\theta CS$ is NCS and by Theorem 3.6 the proof is clear. □

Example 3.11. Let $X = \{u, v\}$ $\Gamma_N = \{0_N, 1_N, F, H, F \cup H\}$ is a N_Γ on \mathcal{Z} . where $H = \langle x, (0, 0, \frac{5}{10}), (1, 1, 0) \rangle$ and $F = \langle x, (1, \frac{5}{10}, 0), (0, \frac{5}{10}, 1) \rangle$. Then the $NS F = \langle x, (1, 1, 0), (0, 0, \frac{5}{10}) \rangle$ is an $NG\theta CS$ but not an $N\alpha GCS$ in Γ .

Theorem 3.12. Every NG^*CS in neutrosophic topological space (\mathcal{Z}, Γ) is an $N\alpha GCS$ but not conversely.

Proof. Let H be an NG^*CS in (\mathcal{Z}, Γ) . By Theorem 3.8 H is an $NGOS$ in (\mathcal{Z}, Γ) . Since $\alpha Cl(H) \subseteq Cl(H)$ and H is a $NGCS$ in \mathcal{Z} . Therefore $\alpha Cl(H) \subseteq Cl(H) \subseteq W$ whenever $H \subseteq W$, W is an NOS in \mathcal{Z} . Thus H is an NG^*CS in \mathcal{Z} . □

Example 3.13. Let $X = \{u, v\}$ $\Gamma_N = \{0_N, 1_N, F\}$ is a N_Γ on \mathcal{Z} . where $F = \langle x, (\frac{3}{10}, \frac{4}{10}, \frac{4}{10}), (\frac{2}{10}, \frac{5}{10}, \frac{5}{10}) \rangle$. Then the $NS H = \langle x, (\frac{1}{10}, \frac{4}{10}, \frac{6}{10}), (\frac{6}{10}, \frac{5}{10}, \frac{2}{10}) \rangle$ is an NG^*CS but not an $N\alpha GCS$ in Γ .

Theorem 3.14. Every $NGRCS$ in neutrosophic topological space (\mathcal{Z}, Γ) is an NG^*CS but not conversely.

Proof. Let H be an $NRCS$ in (\mathcal{Z}, Γ) . Then $H = Cl(Int(H))$. Let $H \subseteq W$, W is an $NGOS$ in \mathcal{Z} . Thus $Cl(H) \subseteq Cl(Int(A))$. This implies $Cl(H) \subseteq H \subset W$. Thus H is an NG^*CS in Γ . H is an NG^*CS in \mathcal{Z} . □

Example 3.15. Let $X = \{u, v\}$ $\Gamma_N = \{0_N, 1_N, F\}$ is a N_Γ on \mathcal{Z} . where $H = \langle x, (\frac{3}{10}, \frac{7}{10}, \frac{2}{10}), (\frac{6}{10}, \frac{3}{10}, \frac{4}{10}) \rangle$. Then the $NS H = \langle x, (\frac{7}{10}, \frac{6}{10}, \frac{3}{10}), (\frac{3}{10}, \frac{4}{10}, \frac{3}{10}) \rangle$ is an NG^*CS but not an $NGRCS$ in Γ .

Theorem 3.16. Every NG^*CS in neutrosophic topological space (\mathcal{Z}, Γ) is an $NG\beta CS$ but not conversely.

Proof. Let H be an NG^*CS in (\mathcal{Z}, Γ) . Let $H \subseteq W$, W is an NOS in \mathcal{Z} . Since every NOS is $NGOS$ and since H is an NG^*CS in (\mathcal{Z}, Γ) . By hypothesis $Cl(H) \subseteq W$. Which implies $Cl(Int(H)) \subseteq W$. Thus $Int(Cl(Int(H))) \subseteq W$. Therefore $\beta Cl(H) \subseteq W$. Thus H is an $NG\beta CS$ in (\mathcal{Z}, Γ) . □

Example 3.17. Let $X = \{u, v\}$ $\Gamma_N = \{0_N, 1_N, F\}$ is a N_Γ on \mathcal{Z} . where $H = \langle x, (\frac{5}{10}, \frac{6}{10}, \frac{4}{10}), (\frac{5}{10}, \frac{4}{10}, \frac{3}{10}) \rangle$. Then the $NS H = \langle x, (\frac{5}{10}, \frac{7}{10}, \frac{3}{10}), (\frac{5}{10}, \frac{3}{10}, \frac{2}{10}) \rangle$ is an $NG * CS$ but not an $NG\beta CS$ in Γ .

Remark 3.18. An $NG * CS$ is independent from $N_\alpha CS$, $NSGCS$, $NPCS$, and $NSCS$ as seen from the following example.

Example 3.19. Let $X = \{u, v\}$ $\Gamma_N = \{0_N, 1_N, F\}$ is a N_Γ on \mathcal{Z} . where $H = \langle x, (\frac{3}{10}, \frac{4}{10}, \frac{4}{10}), (\frac{2}{10}, \frac{5}{10}, \frac{3}{10}) \rangle$. Then the $NS H = \langle x, (\frac{1}{10}, \frac{4}{10}, \frac{5}{10}), (\frac{6}{10}, \frac{5}{10}, \frac{4}{10}) \rangle$ is an $N_\alpha CS$, $NPCS$, $NSCS$ and $NSCS$ but not an $NG * CS$ in (\mathcal{Z}, Γ) .

Example 3.20. Let $X = \{u, v\}$ $\Gamma_N = \{0_N, 1_N, F\}$ is a N_Γ on \mathcal{Z} . where $H = \langle x, (\frac{4}{10}, \frac{3}{10}, \frac{2}{10}), (\frac{6}{10}, \frac{7}{10}, \frac{3}{10}) \rangle$. Then the $NS H = \langle x, (\frac{6}{10}, \frac{7}{10}, \frac{4}{10}), (\frac{1}{10}, \frac{1}{10}, \frac{4}{10}) \rangle$ is an $NG * CS$ but neither $NSCS$ or $N_\alpha CS$ in (\mathcal{Z}, Γ) .

Example 3.21. Let $X = \{u, v\}$ $\Gamma_N = \{0_N, 1_N, F\}$ is a N_Γ on \mathcal{Z} . where $H = \langle x, (0, \frac{9}{10}, \frac{7}{10}), (\frac{5}{10}, \frac{1}{10}, \frac{3}{10}) \rangle$. Then the $NS H = \langle x, (0, \frac{3}{10}, \frac{3}{10}), (\frac{7}{10}, \frac{7}{10}, \frac{4}{10}) \rangle$ is an $NPCS$ but not $NG * CS$ in (\mathcal{Z}, Γ) .

Example 3.22. Let $X = \{u, v\}$ $\Gamma_N = \{0_N, 1_N, H_1, H_2\}$ is a N_Γ on \mathcal{Z} . where $H_1 = \langle x, (\frac{5}{10}, \frac{2}{10}, \frac{2}{10}), (\frac{5}{10}, \frac{7}{10}, \frac{5}{10}) \rangle$ and $H_2 = \langle x, (\frac{8}{10}, \frac{8}{10}, \frac{3}{10}), (\frac{2}{10}, \frac{2}{10}, \frac{4}{10}) \rangle$. Then the $NS H_2 = \langle x, (\frac{5}{10}, \frac{7}{10}, \frac{5}{10}), (\frac{5}{10}, \frac{3}{10}, \frac{4}{10}) \rangle$ is an $NG * CS$ but not an $NPGCS$ in (\mathcal{Z}, Γ) .

Example 3.23. Let $X = \{u, v\}$ $\Gamma_N = \{0_N, 1_N, H_1, H_2\}$ is a N_Γ on \mathcal{Z} . where $H_1 = \langle x, (\frac{5}{10}, \frac{2}{10}, \frac{2}{10}), (\frac{5}{10}, \frac{7}{10}, \frac{5}{10}) \rangle$ and $H_2 = \langle x, (\frac{8}{10}, \frac{8}{10}, \frac{3}{10}), (\frac{2}{10}, \frac{2}{10}, \frac{2}{10}) \rangle$. Then the $NS H_2 = \langle x, (\frac{7}{10}, \frac{7}{10}, \frac{7}{10}), (\frac{3}{10}, \frac{3}{10}, \frac{3}{10}) \rangle$ is an $NG * CS$ but not an $NSGCS$ in (\mathcal{Z}, Γ) .

Remark 3.24. Let (\mathcal{Z}, Γ) be neutrosophic topological space. Then intersection of any two $NG * CS$ s is not an $NG * CS$ in general as seen in the following example.

Example 3.25. Let $X = \{u, v\}$ $\Gamma_N = \{0_N, 1_N, F\}$ is a N_Γ on \mathcal{Z} . where $W = \langle x, (\frac{5}{10}, 0, \frac{1}{10}), (\frac{1}{10}, 1, \frac{9}{10}) \rangle$. Then the $NS H_1 = \langle x, (\frac{2}{10}, 1, \frac{7}{10}), (\frac{7}{10}, 0, \frac{1}{10}) \rangle$, $H_2 = \langle x, (\frac{6}{10}, 0, \frac{1}{10}), (\frac{3}{10}, 1, 1) \rangle$ are $NG * CS$'s in (\mathcal{Z}, Γ) , but $H_1 \cap H_2$ is not an $NG * CS$ in \mathcal{Z} .

Theorem 3.26. Let (\mathcal{Z}, Γ) be neutrosophic topological space. Then the union of any two $NG * CS$ is an $NG * CS$.

Proof. Let H and K be any two $NG * CS$ s in (\mathcal{Z}, Γ) and let $H \cup K \subseteq W$, where W is a $NGOS$ in \mathcal{Z} . Therefore $H \subseteq W$ or $K \subseteq W$ or both subset of W . Since H and K are $NG * CS$, $Cl(H) \subseteq W$ and $Cl(K) \subseteq W$. Therefore $Cl(H \cup K) \subseteq W$. Thus $H \cup K$ is an $NG * CS$. □

Theorem 3.27. Let (\mathcal{Z}, Γ) be neutrosophic topological space. Then H is an $NG * CS$ in (\mathcal{Z}, Γ) iff $\lceil (H_1 q H_2)$ implies $\lceil (Cl(H_1) q H_2)$ for every $NGCS H_2$ of \mathcal{Z} .

Proof. Necessity: Let H_2 be an $NGCS$ of (\mathcal{Z}, Γ) such that $\lceil (H_1 q H_2)$. Then by Definition, $H_1 \subseteq H_2^c$ where H_2^c is an $NGOS$ in \mathcal{Z} . Then $Cl(H_1) \subseteq H_2^c$. Hence $\lceil (Cl(H_1) q H_2)$.

Sufficiency: Let H_2 is an $NGCS$ in \mathcal{Z} . Therefore $\lceil (H_1 q H_2)$, by hypothesis, $\lceil (Cl(H_1) q H_2)$. Then we have $Cl(H_1) \subseteq H_2^c$ whenever $H_1 \subseteq H_2^c$ and H_2^c is an $NGOS$ in \mathcal{Z} . Hence H_1 is an $NG * CS$ in \mathcal{Z} . □

Theorem 3.28. Let (\mathcal{Z}, Γ) be neutrosophic topological space. If H is an $NG * CS$ in (\mathcal{Z}, Γ) , such that $H_1 \subseteq H_2 \subseteq Cl(H_1)$. Then H_2 is also a $NG * CS$ (\mathcal{Z}, Γ) .

Proof. Let W be a $NGOS$ in (\mathcal{Z}, Γ) such that $H_2 \subseteq W$, Since $H_1 \subseteq H_2$, $H_1 \subseteq W$ and W be an $NGOS$. Since H_1 is an $NG * CS$, $Cl(H_1) \subseteq W$. Therefore $H_2 \subseteq Cl(H_1)$, by hypothesis. $Cl(H_2) \subseteq Cl(Cl(H_1)) \subseteq W$. Then we have $Cl(H_2) \subseteq W$. Hence H_2 is an $NG * CS$ in \mathcal{Z} . □

Theorem 3.29. Let (\mathcal{Z}, Γ) be neutrosophic topological space. Then $NGO(\Gamma) = NGC(\Gamma)$ iff every NS in (\mathcal{Z}, Γ) is an $NG * CS$ in \mathcal{Z} .

Proof. Necessity: Suppose that $NGO(\Gamma) = NGC(\Gamma)$. Let $H_1 \subseteq W$ and W is an $NGOS$ in Γ . This implies $Cl(H_1) \subseteq Cl(W)$. Since W is an $NGOS$ in Γ . Since by hypothesis W is an $NGCS$ in Γ , $Cl(W) \subseteq W$. This implies $Cl(H_1) \subseteq W$. Therefore H_1 is an $NG * CS$ in Γ .

Sufficiency: Suppose that every NS in (\mathcal{Z}, Γ) is an $NG * CS$ in Γ . Let $W \in NO(\Gamma)$, then $W \in NGO(\Gamma)$. Since $W \subseteq W$ and W is NOS in Γ , by hypothesis $Cl(W) \subseteq W$. That is $W \in NGC(\Gamma)$. Hence $NGO(\Gamma) \subseteq NGC(\Gamma)$. Let $H \in NGC(\Gamma)$ then H^c is an $NGOS$ in Γ . But $NGO(\Gamma) \subseteq NGC(\Gamma)$. Therefore $H^c \in NGC(\Gamma)$. That is $H \in NGO(\Gamma)$. Hence $NGC(\Gamma) \subseteq NGO(\Gamma)$. Thus $NGO(\Gamma) \subseteq NGC(\Gamma)$. □

Theorem 3.30. Let (Z, Γ) be neutrosophic topological space. If H is NG^*CS and $NGOS$ in (Z, Γ) , then H_1 is an NCS in Z .

Proof. Let H_1 is an $NGOS$ in Z . Since $H_1 \subseteq H_1$, by hypothesis $Cl(H_1) \subseteq H_1$. But from the Definition, $H_1 \subseteq Cl(H_1)$. Therefore $Cl(H_1) = H_1$. Hence H_1 is an NCS of Z . \square

Theorem 3.31. Let (Z, Γ) be neutrosophic topological space. If H is an NG^*CS in (Z, Γ) and $w(u, v)$ be a neutrosophic point in Z such that $w(u, v)qCl(H)$ then $Cl(w(u, v)qH)$.

Proof. Let H be an NG^*CS in Z and let $w(u, v)qCl(H)$. Let $\}(Cl(w(u, v)qH)$, then $H \subseteq [Cl(w(u, v))]^c$, where $[Cl(w(u, v))]^c$ is an $NGOS$ in Z . Since $Cl(H) \subseteq [Cl(w(u, v))]^c \subseteq [w(u, v)]^c$, by hypothesis, we have $\}(w(u, v)qCl(H))$, which is a contradiction to the hypothesis. Therefore $Cl((w(u, v)qH)$. \square


Theorem 3.32. Let (Z, Γ) be neutrosophic topological space. If H is an NOS and an NG^*CS in (Z, Γ) , then

1. H is an $NROS$ in Z ,
2. H is an $NRCS$ in Z .

Proof. (1) Let H be an NOS and an NG^*CS in Z . Hence $Cl(H) \subseteq H$. That is $Int(Cl(H)) \subseteq H$. Since H is an NOS , H is an $NPOS$ in Z . Then $H \subseteq Int(Cl(H))$. Therefore $H = Int(Cl(H))$. Hence H is an $NROS$ in Z .

(2) Let H be an NOS and an NG^*CS in Z . Then $Cl(H) \subseteq H$. That is $Cl(int(H)) \subseteq H$. Since H is an NOS , H is an $NSOS$ in Z . By hypothesis, we have $H \subseteq Cl(Int(H))$. This implies $H = Int(Cl(H))$. Hence H is an $NRCS$ in Z . \square

Remark 3.33. From above the following implication between NG^*CS and the other existed NCS 's and none of these implications is reversible



Theorem 3.34. Let (Z, Γ) be neutrosophic topological space. If H is an NG^*CS in (Z, Γ) , then $Cl(H) \setminus H$ contains non-empty $NGCS$.

Proof. Suppose that H is an NG^*CS of (Z, Γ) and let W be non-empty $NGCS$ of (Z, Γ) , such that $W \subseteq Cl(H) \setminus H$. Therefore $H \subseteq Z \setminus W$. By hypothesis, we have H is NG^*CS and $Z \setminus W$ is $NGOS$, then $Cl(H) \subseteq Z \setminus W$. This implies $W \subseteq Z \setminus Cl(H)$. So $W \subseteq (Z \setminus cl(H)) \cap (cl(H) \setminus H) \subseteq (Z \setminus Cl(H)) \cap cl(H) = \phi$. Hence W is empty set. \square

4 Neutrosophic g^* -Open Sets

In this section, we introduce Neutrosophic g^* -open set in Neutrosophic topological space and study some of their properties.

Definition 4.1. A neutrosophic set H (Ns in short) is said to be Neutrosophic g^* -open set (NG^*OS in short) in (\mathcal{Z}, Γ) if the complement H^c is an NG^*CS in \mathcal{Z} . The family of all NG^*OS 's of an NTS (\mathcal{Z}, Γ) is denoted by $NG^*O(\mathcal{Z})$.

Theorem 4.2. Let (\mathcal{Z}, Γ) be neutrosophic topological space. Then A subset H_1 of \mathcal{Z} is an NG^*OS if and only if $H_2 \subseteq Int(H_1)$ whenever H_2 is an $NGCS$ in \mathcal{Z} and $H_2 \subseteq H_1$.

Proof. Necessity: Let H_1 is an NG^*OS in \mathcal{Z} . Let H_2 be an $NGCS$ in \mathcal{Z} and $H_2 \subseteq H_1$. Then H_2^c is an $NGOS$ in \mathcal{Z} such that $H_1^c \subseteq H_2^c$. Since H_1^c is an NG^*CS , we have $Cl(H_1^c) \subseteq H_2^c$. Hence $(Int(H_1))^c \subseteq H_2^c$. Therefore $H_2 \subseteq Int(H_1)$.

Sufficiency: Let $H_2 \subseteq Int(H_1)$ whenever H_2 is an $NGCS$ in and $H_2 \subseteq H_1$. Then $H_1^c \subseteq H_2^c$ and H_2^c is an $NGOS$. By hypothesis, $(Int(H_1))^c \subseteq H_2^c$, which implies $Cl(H_1^c) \subseteq H_2^c$. Therefore H_1^c is an NG^*CS of \mathcal{Z} . Hence H_1 is an NG^*OS in \mathcal{Z} . \square

Theorem 4.3. Let (\mathcal{Z}, Γ) be neutrosophic topological space. Then for every NOS is an NG^*OS but not conversely.

Proof. Let H be an NOS . Then H^c is an NCS . By Theorem 3.6, every NCS is an NG^*CS . Therefore H^c is an NG^*CS . Hence H is an NG^*OS . \square

Example 4.4. Let $X = \{u, v\} \Gamma_N = \{0_N, 1_N, F\}$ is a N_Γ on \mathcal{Z} . where $H = \langle x, (\frac{5}{10}, \frac{3}{10}, \frac{3}{10}), (\frac{7}{10}, \frac{8}{10}, \frac{8}{10}) \rangle$. Then the $NS H = \langle x, (\frac{2}{10}, \frac{2}{10}, \frac{2}{10}), (\frac{7}{10}, \frac{8}{10}, \frac{8}{10}) \rangle$ is an NG^*OS but not an NCS in (\mathcal{Z}, Γ) .

Theorem 4.5. Let (\mathcal{Z}, Γ) be neutrosophic topological space. Then every $NROS$ is an NG^*OS but not conversely.

Proof. Let H be an $NROS$ in (\mathcal{Z}, Γ) . Then H^c is an $NRCS$. By Theorem 3.14, every $NRCS$ is an NG^*CS . Therefore H^c is an NG^*CS . Hence H is an NG^*OS . \square

Example 4.6. Let $X = \{u, v\} \Gamma_N = \{0_N, 1_N, F\}$ is a N_Γ on \mathcal{Z} . where $H = \langle x, (\frac{3}{10}, \frac{7}{10}, \frac{7}{10}), (\frac{6}{10}, \frac{3}{10}, \frac{3}{10}) \rangle$. Then the $NS H = \langle x, (\frac{3}{10}, \frac{4}{10}, \frac{4}{10}), (\frac{7}{10}, \frac{6}{10}, \frac{6}{10}) \rangle$ is an NG^*OS but not an $NROS$ in (\mathcal{Z}, Γ) .

Theorem 4.7. Let (\mathcal{Z}, Γ) be neutrosophic topological space. Then every NG^*OS is an $NGOS$ but not conversely.

Proof. Let H be an NG^*OS in (\mathcal{Z}, Γ) . Then H^c is an NG^*CS . By Theorem 3.8, every NG^*CS is an $NGCS$. Therefore H^c is an $NGCS$. Hence H is an $NGOS$. \square

Example 4.8. Let $X = \{u, v\} \Gamma_N = \{0_N, 1_N, F\}$ is a N_Γ on \mathcal{Z} . where $H = \langle x, (\frac{5}{10}, \frac{2}{10}, \frac{3}{10}), (\frac{4}{10}, \frac{2}{10}, \frac{2}{10}) \rangle$. Then the $NS H = \langle x, (\frac{4}{10}, \frac{2}{10}, \frac{4}{10}), (\frac{6}{10}, \frac{6}{10}, \frac{6}{10}) \rangle$ is an $NGOS$ but not an NG^*OS in (\mathcal{Z}, Γ) .

Theorem 4.9. Let (\mathcal{Z}, Γ) be neutrosophic topological space. Then every NG^*OS is a $N\alpha GOS$ but not conversely.

Proof. Let H be a NG^*OS in (\mathcal{Z}, Γ) . Then H^c is a NG^*CS . By Theorem 3.12, every NG^*CS is a $N\alpha GCS$. Therefore H^c is a $N\alpha GCS$. Hence H is a $N\alpha GOS$. \square

Example 4.10. Let $X = \{u, v\} \Gamma_N = \{0_N, 1_N, F\}$ is a N_Γ on \mathcal{Z} . where $H = \langle x, (\frac{4}{10}, \frac{2}{10}, \frac{3}{10}), (\frac{6}{10}, \frac{7}{10}, \frac{6}{10}) \rangle$. Then the $NS H = \langle x, (\frac{6}{10}, \frac{5}{10}, \frac{4}{10}), (\frac{4}{10}, \frac{3}{10}, \frac{6}{10}) \rangle$ is an $N\alpha GOS$ but not a NG^*OS in (\mathcal{Z}, Γ) .

Theorem 4.11. Let (\mathcal{Z}, Γ) be neutrosophic topological space. Then The intersection of any two NG^*OS 's is a NG^*OS .

Proof. Let H_1 and H_2 be the any two NG^*OS 's in \mathcal{Z} , H_1^c and H_2^c are NG^*CS sets. By Theorem 3.26, $H_1^c \cup H_2^c$ is an NG^*CS in \mathcal{Z} . Therefore $(H_1 \cap H_2)^c$ is an NG^*CS . Thus $H_1 \cap H_2$ is an NG^*OS in \mathcal{Z} . □

Remark 4.12. Let (\mathcal{Z}, Γ) be NT Space. The union of any two NG^*OS 's is not a NG^*OS in general as seen in the following example.


Example 4.13. Let $X = \{u, v\} \Gamma_N = \{0_N, 1_N, F\}$ is a N_Γ on \mathcal{Z} . where $F = \langle x, (\frac{5}{10}, 0, 0), (\frac{1}{10}, 1, 1) \rangle$. Then the NS 's $H_1 = \langle x, (\frac{7}{10}, 0, 0), (\frac{2}{10}, 1, 1) \rangle$, $H_2 = \langle x, (\frac{3}{10}, 1, 1), (\frac{6}{10}, 0, 0) \rangle$ is are NG^*OS in (\mathcal{Z}, Γ) but $H_1 \cup H_2$ is not an NG^*CS in (\mathcal{Z}, Γ) .

Theorem 4.14. Let (\mathcal{Z}, Γ) be neutrosophic topological space. Then every NG^*OS is an $N\beta GOS$ but not conversely.

Proof. Let H be an NG^*OS in (\mathcal{Z}, Γ) . Then H^c is an NG^*CS . By Theorem 4.3, every NG^*CS is an $N\beta GCS$. Therefore H^c is a $N\beta GCS$. Hence H is a $N\beta GOS$. □

Example 4.15. Let $X = \{u, v\} \Gamma_N = \{0_N, 1_N, F\}$ is a N_Γ on \mathcal{Z} . where $F = \langle x, (\frac{5}{10}, \frac{6}{10}, \frac{6}{10}), (\frac{5}{10}, \frac{4}{10}, \frac{4}{10}) \rangle$. Then the NS $H = \langle x, (\frac{5}{10}, \frac{3}{10}, \frac{4}{10}), (\frac{5}{10}, \frac{7}{10}, \frac{6}{10}) \rangle$ is an $N\beta GOS$ but not an NG^*OS in (\mathcal{Z}, Γ) .

Remark 4.16. From above the following implication between NG^*OS and the other existed NOS 's, $NGOS$'s and none of these implications is reversible



Theorem 4.17. Let (\mathcal{Z}, Γ) be neutrosophic topological space. Then if H is a NS of \mathcal{Z} then have the following equivalent properties:

1. H in $NG^*O(\Gamma)$,
2. $F \subseteq Int(H)$ whenever $F \subseteq H$ and F is a $NGCS$ in \mathcal{Z} ,
3. There exists OS 's Q_1 and Q_2 such that $Q_2 \subseteq F \subseteq Q_1$, where $Q_1 = Int(H)$, $F \subseteq H$ and F is a NCS in \mathcal{Z} .

Proof. (1) \rightarrow (2): Let H in $NG^*O(\mathcal{Z})$. Then H^c is an NG^*CS in \mathcal{Z} . Therefore $Cl(H^c) \subseteq W$ whenever $H^c \subseteq W$ and W is a $NGOS$ in \mathcal{Z} . By taking the complement on both sides $[Cl(H^c)]^c \supseteq W^c$ whenever $[H^c]^c \supseteq W^c$. Therefore $W^c \subseteq Int(H)$ whenever $W^c \subseteq H$ and W^c is a $NGCS \in \mathcal{Z}$. Replacing W^c by F , $F \subseteq Int(H)$ whenever $F \subseteq H$ and F is a $NGCS \in \mathcal{Z}$.

(2) \rightarrow (3): $F \subseteq Int(H)$ whenever $F \subseteq H$ and F is a $NGCS \in \mathcal{Z}$. Hence $Int(F) \subseteq F \subseteq Int(H)$ then there exists NOS 's Q_1 and Q_2 such that $Q_2 \subseteq F \subseteq Q_1$, where $Q_1 = Int(H)$ and $Q_2 = Int(F)$.

(3) \rightarrow (1): Suppose that there exists NOS 's Q_1 and Q_2 such that $Q_2 \subseteq F \subseteq Q_1$. That is $F \subseteq Int(H)$. Then $Cl(H^c) \subseteq F^c$ whenever $H^c \subseteq F^c$ and F^c is a $NGOS \in \mathcal{Z}$. Hence H^c is a $NG^*CS \in \mathcal{Z}$. Therefore H in $NG^*O(\mathcal{Z})$. □

Theorem 4.18. Let (\mathcal{Z}, Γ) be neutrosophic topological space. Then if H is a NS of \mathcal{Z} . Then for every H in $NG^*O(\mathcal{Z})$ and every R in $NS(\mathcal{Z})$, $Int(H) \subseteq R \subseteq H$ implies R in $NG^*O(\mathcal{Z})$.

Proof. Since $Int(H) \subseteq R \subseteq H$. By taking the complement on both sides, we get $H^c \subseteq R^c \subseteq Cl(H^c)$. Let $R^c \subseteq F$ and F is a $NGOS \in \mathcal{Z}$. Since $H^c \subseteq R^c$, $H^c \subseteq F$. Since H^c is a NG^*CS , $Cl(H^c) \subseteq F$. Therefore $Cl(R^c) \subseteq Cl(H^c) \subseteq F$. Hence R^c is an $NG^*CS \in \mathcal{Z}$. Therefore R is a $G^*OS \in \mathcal{Z}$. That is R in $NG^*O(\mathcal{Z})$. \square

5 Separation Axioms of Neutrosophic g^* - Closed Sets

In this section, we study and introduce the concepts of neutrosophic $T_{\frac{1}{2}}^*$ space ($NT_{\frac{1}{2}}^*$ in short), neutrosophic $^*T_{\frac{1}{2}}$ space and some of their basic properties. Also we study some applications of Neutrosophic g^* - Closed Sets.

Definition 5.1. An neutrosophic topological space (\mathcal{Z}, Γ) is said to be an neutrosophic $T_{\frac{1}{2}}^*$ space (in short $NT_{\frac{1}{2}}^*$) if every NG^*CS of (\mathcal{Z}, Γ) is a NCS of (\mathcal{Z}, Γ) .

Theorem 5.2. A neutrosophic space topological (\mathcal{Z}, Γ) is neutrosophic $T_{\frac{1}{2}}^*$ space iff $NG^*OS(\mathcal{Z}) = NOS(\mathcal{Z})$

Proof. Necessity: Let H be an NG^*CS in \mathcal{Z} then H^c is a NG^*CS in \mathcal{Z} . By our assumption H^c is an NCS in \mathcal{Z} , this implies H is a NOS of \mathcal{Z} . Then NG^*OS of $\mathcal{Z} = NOS(\mathcal{Z})$.

Sufficiency: Let H be a NG^*CS in \mathcal{Z} , the H^c is a NG^*OS of \mathcal{Z} . By hypothesis H^c is $NOS \in \mathcal{Z}$, therefore H is a $NCS \in \mathcal{Z}$. Then (\mathcal{Z}, Γ) is neutrosophic $T_{\frac{1}{2}}^*$ space. \square

Theorem 5.3. Let (\mathcal{Z}, Γ) be neutrosophic topological space. Then every neutrosophic $T_{\frac{1}{2}}$ space is neutrosophic $T_{\frac{1}{2}}^*$ space.

Proof. Assume (\mathcal{Z}, Γ) be an $NT_{\frac{1}{2}}$ space and let H be an NG^*CS in (\mathcal{Z}, Γ) . Since every NG^*CS is a $NGCS$. However, H is an NCS . Hence \mathcal{Z} is an $NT_{\frac{1}{2}}^*$ space. \square

Remark 5.4. Every neutrosophic $T_{\frac{1}{2}}^*$ space need not be neutrosophic $T_{\frac{1}{2}}$ space in (\mathcal{Z}, Γ) in general as seen from the following example.

Example 5.5. Let $X = \{u, v\}$ $\Gamma_N = \{0_N, 1_N, F\}$ is a N_Γ on \mathcal{Z} . where $F = \langle x, (\frac{9}{10}, \frac{9}{10}, \frac{9}{10}), (\frac{1}{10}, \frac{1}{10}, \frac{1}{10}) \rangle$. Clearly (\mathcal{Z}, Γ) is an neutrosophic $T_{\frac{1}{2}}^*$ space, but not neutrosophic $T_{\frac{1}{2}}$ space.

Theorem 5.6. For any neutrosophic space topological (\mathcal{Z}, Γ) the following are equivalent:

1. (\mathcal{Z}, Γ) is neutrosophic $T_{\frac{1}{2}}$ space,
2. Every singleton element of \mathcal{Z} is either $NGCS$ or NOS .

Proof. (1) \rightarrow (2): Let $r \in \mathcal{Z}$ and Suppose r is not a $NGCS$ in (\mathcal{Z}, Γ) . Then $\mathcal{Z} \setminus r$ is not $NGOS$, This implies \mathcal{Z} is the only $NGOS$ containing $\mathcal{Z} \setminus r$. Therefore $\mathcal{Z} \setminus r$ is a $NGOS$ of (\mathcal{Z}, Γ) . Since (\mathcal{Z}, Γ) is a neutrosophic $T_{\frac{1}{2}}^*$ space, then $\mathcal{Z} \setminus r$ is NCS or equivalently r is an NOS in (\mathcal{Z}, Γ) .

(2) \rightarrow (1) Let H be an NG^*CS in (\mathcal{Z}, Γ) . Its Obvious that $H \subseteq Cl(H)$. Let $r \in Cl(H)$. By (2) r is either NOS or $NGCS$. Case (1) let r is NOS . Since $r \in Cl(H)$, then $r \cap H \neq \phi$. So $r \in H$. Then in any case $r \in H$. Thus $H = Cl(H)$ or equivalently H is a NCS . Thus every NG^*CS is NCS . Hence (\mathcal{Z}, Γ) is neutrosophic $T_{\frac{1}{2}}^*$ space. Case (2) let r is $NGCS$. If $r \notin H$ then $Cl(H) \setminus H$ contains non-empty $NGCS$ r . But this is not possible according to the theorem 3.34 as H is NG^*CS . Hence $r \in H$. \square

Theorem 5.7. Let (\mathcal{Z}, Γ) be a NTS and \mathcal{Z} is a neutrosophic $T_{\frac{1}{2}}^*$ space. Then the following properties hold:


1. Any intersection of NG^*OS is NG^*OS in \mathcal{Z} ,
2. Any union of NG^*CS is NG^*CS in \mathcal{Z} .

Proof. (1) It is obvious from (1) by taking complement.

(2) Let $H_{j \in I}$ be a collection of NG^*CS 's in a neutrosophic $T_{\frac{1}{2}}^*$ space. Thus, every NG^*CS is an NCS of \mathcal{Z} . However, the union of NCS is a NCS in \mathcal{Z} . Hence $\cup H_{j \in I}$ is an $NCS \in \mathcal{Z}$. Since every NCS is an NG^*CS , $\cup H_{j \in I}$ is an NG^*CS in \mathcal{Z} . Therefore any union of NG^*CS 's is NG^*CS in \mathcal{Z} . \square

Definition 5.8. A neutrosophic topological space (\mathcal{Z}, Γ) is said to be a neutrosophic ${}^*T_{\frac{1}{2}}$ space (in short $N^*T_{\frac{1}{2}}$) if every $NGCS$ of (\mathcal{Z}, Γ) is a NG^*CS of (\mathcal{Z}, Γ) .

Remark 5.9. The following diagram, we have provided the relation between $NT_{\frac{1}{2}}^*$, $N^*T_{\frac{1}{2}}$ and $NT_{\frac{1}{2}}$.



Theorem 5.10. If (\mathcal{Z}, Γ) is an $N^*T_{\frac{1}{2}}$ space then each $r \in \mathcal{Z}$, r is either NCS or NG^*OS .

Proof. Suppose (\mathcal{Z}, Γ) is a $N^*T_{\frac{1}{2}}$ space. Let $r \in \mathcal{Z}$ and let that r is $NGCS$ since \mathcal{Z} is the only open set which contains $\mathcal{Z} \setminus r$. Since (\mathcal{Z}, Γ) is a $N^*T_{\frac{1}{2}}$ then $\mathcal{Z} \setminus r$ is an NG^*CS or equivalently r is NG^*OS . \square

Theorem 5.11. For any neutrosophic topological space (\mathcal{Z}, Γ) , neutrosophic $T_{\frac{1}{2}}$ space is neutrosophic ${}^*T_{\frac{1}{2}}$ space.

Proof. Let \mathcal{Z} be a $N^*T_{\frac{1}{2}}$ space and let H be an $NGCS$ of (\mathcal{Z}, Γ) . By hypothesis H is an NCS . Since every NCS in (\mathcal{Z}, Γ) is a NG^*CS of (\mathcal{Z}, Γ) . Hence \mathcal{Z} is a $N^*T_{\frac{1}{2}}$ space. \square

Theorem 5.12. A neutrosophic topological space (\mathcal{Z}, Γ) is an $NT_{\frac{1}{2}}$ space iff it is both $N^*T_{\frac{1}{2}}$ space and $NT_{\frac{1}{2}}^*$ space.

Proof. Necessity: Follows from Theorem 5.3 and 5.11.

Sufficiency: Suppose (\mathcal{Z}, Γ) is both $N^*T_{\frac{1}{2}}$ space and $NT_{\frac{1}{2}}^*$ space. Let H be a $NGCS$ of (\mathcal{Z}, Γ) . Since (\mathcal{Z}, Γ) is an ${}^*T_{\frac{1}{2}}$ space, then H is NG^*CS of (\mathcal{Z}, Γ) . Since (\mathcal{Z}, Γ) is a $NT_{\frac{1}{2}}^*$ space, then H is NCS of (\mathcal{Z}, Γ) . Thus (\mathcal{Z}, Γ) is a $NT_{\frac{1}{2}}$ space. \square

Theorem 5.13. A neutrosophic topological space (\mathcal{Z}, Γ) is a $N^*T_{\frac{1}{2}}$ space iff $NG^*OS(\mathcal{Z}) = NGOS(\mathcal{Z})$

Proof. Necessity: Assume H be a $NGOS(\mathcal{Z}) \in \mathcal{Z}$, then H^c is a $NGCS \in \mathcal{Z}$. However H^c is a NG^*CS in \mathcal{Z} , and this implies H is an NG^*OS in \mathcal{Z} . Hence $NGOS(\mathcal{Z}) = NG^*OS(\mathcal{Z})$.

Sufficiency: Assume H be a $NGCS$ of \mathcal{Z} , the H^c is an $NGOS$ in \mathcal{Z} . By Hypothesis H^c is a NG^*OS in \mathcal{Z} , which implies H is a NG^*CS in \mathcal{Z} . Hence (\mathcal{Z}, Γ) is a $N^*T_{\frac{1}{2}}$ space. \square

6 Conclusion

In this write up, we established inequalities and introduce the notion of Neutrosophic g^* set, Neutrosophic $*T_{\frac{1}{2}}$ space by using Ng^* set in Neutrosophic topological spaces. Furthermore, we investigate and define NG^*CS , NG^*OS . Then we introduce and study the relationships between NG^*CS , NG^*OS and other generalization closed and open sets respectively, and also present some properties and applications of Neutrosophic g^* set in Neutrosophic topological space. This work can easily be extended in various structures like Neutrosophic Pg^* , Neutrosophic Sg^* , Neutrosophic αg^* sets. The other properties of these types of sets can be found and one can introduce some other relations to these types of sets to develop the skills of learning mathematics.

7 Author's contributions

This article was written in collaboration by all of the contributors. The final manuscript was read and approved by all writers.

8 Conflicts of interest

There are no competing interests declared by the authors.

References

- [1] M. Scheepers, *Combinatorics of open covers i: Ramsey theory*, Topology Appl. vol. 69, pp. 31–62, 1996.
- [2] J. Dontchev, *Survey on preopen sets*, proc. Yatsushiro Topological Conf. pp. 1–8, 1998.
- [3] A. A. El-Aziz, *On generalized forms of compactness*, Masters thesis, Faculty of Science, Tanta University, Egypt.
- [4] D.Jayanthi, α Generalized closed Sets in Neutrosophic Topological Spaces, International Journal of Mathematics Trends and Technology (IJMTT)- Special Issue ICRMIT March. Infinite Study, (2018).
- [5] A.Mary Margaret, and M.Trinita Pricilla., Neutrosophic Vague Generalized Pre-closed Sets in Neutrosophic Vague Topological Spaces, International Journal of Mathematics And its Applications, Vol. 5, Issue 4-E, pp. 747–759, 2017.
- [6] V.K.Shanthi.V.K.,S.Chandrasekar.S, K.SafinaBegam, Neutrosophic Generalized Semi closed Sets in Neutrosophic Topological Spaces, International Journal of Research in Advent Technology, Vol.(ii),6, No.7,July pp. 1739–1743,2018.
- [7] Renu Thomas, and S. Anila, On Neutrosophic Semi-preopen Sets and Semi-preclosed Sets in a Neutrosophic Topological Space, International Journal of Scientific Research in Mathematical and Statistical Sciences, Vol. 5, No. 5, pp. 138–143, 2018.
- [8] Narmatha, S., E. Glory Bebina, and R. Vis.hnu Priyaa. On $\pi g\beta$ -Closed Sets and Mappings in Neutrosophic Topological Spaces. International Journal of Innovative Technology and Exploring Engineering (IJITEE) 8.125, pp. 505–510, 2019.
- [9] W. F. Al-Omeri, F. Smarandache, New Neutrosophic Sets via Neutrosophic Topological Spaces, Brussels (Belgium): Pons, pp. 189–209, 2017.
- [10] W. F. Al-Omeri, Neutrosophic crisp sets via neutrosophic crisp topological spaces, Neutrosophic Sets and Systems.13. pp. 96–104, 2016.

- [11] W. F. Al-Omeri, On Mixed b -Fuzzy Topological Spaces, Journal International Journal of Fuzzy Logic and Intelligent Systems, Volume 20,3. Pp. 242–246,2020.
- [12] W. F. Al-Omeri, S.Jafari and F. Smarandache, Neutrosophic fixed point theorems and cone metric spaces, Neutrosophic Sets and Systems, 2020, 31, 250-265. doi:10.5281/zenodo.3640600.
- [13] W. F. Al-Omeri, F. Smarandache, and S. Jafari, (Φ, Ψ) -Weak Contractions in Neutrosophic Cone Metric Spaces via Fixed Point Theorems,2020, Journal Mathematical Problems in Engineering,8, Pages 1-8.
- [14] A Ghareeb, Wadei F. Al-Omeri, New degrees for functions in (L, M) -fuzzy topological spaces based on (L, M) -fuzzy semiopen and (L, M) -fuzzy preopen operators,Journal of Intelligent and Fuzzy Systems, 36 (1), 787-803.
- [15] W. Al-Omeri, and S. Jafari. On Generalized Closed Sets and Generalized Pre-Closed Sets in Neutrosophic Topological Spaces, Mathematics, pp. 1–12, 2019.
- [16] A. A. Salama, Said Broumi, S. A. Alblowi, Introduction to Neutrosophic Topological Spatial Region, Possible Application to GIS Topological Rules, IJ. Information
- [17] R. Dhavaseelan, and S. Jafari. Generalized Neutrosophic closed sets, New trends in Neutrosophic theory and applications Volume II-261-273,(2018) R. Dhavaseelan, S. Jafari and md. Hanif page, Neutrosophic generalized α -contra-continuity, creat. math. inform Vol. 2334, pp.133–139, 2018.
- [18] D. Coker, An introduction to intuitionistic fuzzy topological space, Fuzzy Sets and Systems, Vol. 88, No. 1, pp. 81–89, 1997.
- [19] C. L. Chang, Fuzzy topological spaces, Journal of Mathematical Analysis and Applications Vol. 24, No. 1 , pp. 182–190, 1968.
- [20] F. Smarandache, Neutrosophic Probability, Set, and Logic, ProQuest Information, Learning, Ann Arbor., Michigan, USA., pp. 105, 1998.
- [21] M. A. A. Shumrani, F. Smarandache, Introduction to non-standard neutrosophic topology, Symmetry Vol. 11 (0), pp. 1-14,2019, basel, Switzerland. doi:10.3390/sym11050000.
- [22] K. Atanassov, Intuitionistic fuzzy sets, VII ITKR's Session, (Deposed in Central Sci.- Techn. Library of Bulg. Acad. of Sci., 1697/84) (in Bulg.), 1984.
- [23] C. Maheswari, and S. Chandrasekar. Neutrosophic gb -closed Sets and Neutrosophic gb -Continuity. Infinite Study, 2020.
- [24] M. Parimala , M. Karthika , Florentin Smarandache , Said Broumi, On $\alpha\omega$ -closed sets and its connectedness in terms of neutrosophic topological spaces, International Journal of Neutrosophic Science, Vol. 2 , No. 2 , (2020) : 82-88 (Doi : <https://doi.org/10.54216/IJNS.020204>)
- [25] A. Vadivel , C. John Sundar, Neutrosophic δ -Open Maps and Neutrosophic δ -Closed Maps, International Journal of Neutrosophic Science, Vol. 13 , No. 2 , pp. 66–74, 2021 (Doi : <https://doi.org/10.54216/IJNS.130203>)
- [26] P. Basker , Broumi Said, $N\Psi_0^\# \alpha$ and $N\Psi_1^\# \alpha$ -spaces in Neutrosophic Topological Spaces, International Journal of Neutrosophic Science, Vol. 16 , No. 1 , pp.09–15, 2021 (Doi : <https://doi.org/10.54216/IJNS.160101>)
- [27] Acikgoz, Ahu, and Ferhat Esenbel. A study on connectedness in neutrosophic opological spaces. AIP Conference Proceedings. Vol. 2334. No. 1. AIP Publishing LLC, 2021.