



Decision Support Modeling For Agriculture Land Selection Based On Sine Trigonometric Single Valued Neutrosophic Information

¹ Shahzaib Ashraf, ² Saleem Abdullah

^{1,2}Department of Mathematics, Abdul Wali Khan University, Mardan 23200, PAKISTAN

shahzaibashraf@awkum.edu.pk¹, saleemabdullah@awkum.edu.pk²

Abstract

A single valued neutrosophic set (SVNS) is a useful tool to portray uncertainty in multi attribute decision-making. In this article, we develop hybrid averaging and hybrid geometric aggregation operator using sine trigonometric function to handle uncertainty in single valued neutrosophic information, which are, sine trigonometric-single valued neutrosophic hybrid weighted averaging (ST-SVNHWA) operator and , sine trigonometric-single valued neutrosophic hybrid weighted geometric (ST-SVNHWG) operator. We investigate properties, namely, idempotancy, monotonicity and boundedness for the proposed operators. Moreover, we give an algorithm to solve multi-criteria decision-making issues which involve SVN information with ST-SVNHWA and ST-SVNHWG operators. Finally, an illustrative example of agricultural land selection is provided to verify the effectiveness. Sensitivity and comparative analyses are also implemented to assess the stability and validity of our method.

Keywords:

Single valued neutrosophic set, Sine trigonometric single valued neutrosophic information, Agriculture land selection, Decision Support.

1 Introduction

Multi-criteria decision-making (MCDM) is performing a vital role in different areas, including social, physical, medical and environmental sciences. MCDM methods use not only to evaluate an appropriate object but also to rank the objects in a given problem. To solve different uncertain problems for decision-making, Atanassov¹³ presented the concept of intuitionistic fuzzy set (IFS) to include both membership and non membership parts, an extension of fuzzy set⁴⁷ in which simply membership part is characterized. After that, various hybrid models of fuzzy sets (FSs) have been presented and investigated such as, Pythagorean fuzzy sets (PyFSs),⁴⁶ Picture fuzzy sets (PFSs),¹⁷ Spherical fuzzy sets (SFSs)^{1,2} and single-valued neutrosophic set (SVNS).^{38,40}

Aggregation operators (AOs) perform an important role in order to combine data into a single form and solve MCDM problems. Aggregation implies the invention of a numeral of things to a cluster or a bunch of objects that have come or been taken together. In the past few years, aggregation operators based on FSs and its various hybrid compositions have made very much attention and become attractive because they can quickly execute functional areas of diverse regions. For example, Yager⁴⁵ introduced weighted aggregation operators (AOs). Khan et al.,²³ presented the probabilistic hesitant based DM technique. Xu⁴¹ proposed some new AOs under IFSs. Khan et al.,²⁵ established the novel decision making (DM) methodology under generalized intuitionistic fuzzy soft information. Khan et al.,²⁴ established the Dombi AOs under PyFSs. Ashraf et al.,⁵ proposed the fuzzy decision support modelling for internet finance soft power evaluation based on sine trigonometric Pythagorean fuzzy information. Batool et al., established the entropy based DM method under probabilistic Pythagorean hesitant fuzzy information. Sajjad et al.,³⁹ established the TOPSIS approach under PyFSs. Ashraf et al.,⁶ presented the AOs based on algebraic norm under PFSs. Khan et al.,²⁶ presented the AOs based on Einstein norm under PFSs. Ashraf et al.,⁷ introduced the DM method under picture cubic fuzzy sets (PCFSs). Qiyas et al.,^{34,35} established the AOs based on the Dombi and algebraic norm using

linguistic picture fuzzy information respectively. Rafiq et al.,³⁷ proposed the cosine trigonometric function based similarity measure under SFSs. Zang et al.,⁴⁸ proposed the DM approach using TOPSIS under spherical fuzzy rough set. Ashraf et al.,⁸ introduced the norm representation under SFSs. Ashraf et al.,⁹ presented the AOs using Dombi norm under SFSs. Jin et al.,²⁰ introduced the AOs for hybrid structure of linguistic and SFSs. Ashraf et al.,¹⁰ presented the AOs for SFSs using the symmetric sum technique and discussed their application in DM. Jin et al.,²¹ established the logarithmic function based AOs under SFSs. Ashraf et al.,¹¹ presented the DM technique using distance measures under SFSs. Barukab et al.,¹⁴ introduced the entropy measure based extended TOPSIS under SFSs. Ashraf et al.,¹² established the AOs using logarithmic function under SVNNSs. Ye⁴² established the correlation coefficient based DM approach under SVNNSs. Liu et al.,²⁸ proposed the Dombi norm based DM methodology under linguistic SVNNSs. Liu et al.,²⁹ established the muirhead mean based power AOs under SVNNSs. Liu et al.,^{30,31} presented the Heronian mean based power AOs using linguistic SVNNSs and cubic neutrosophic information respectively. Ji et al.,²² presented the Bonferroni mean AOs using Frank norm under SVNNSs. Ye⁴³ introduced the list of novel AOs using exponential function under SVNNSs and discussed their application to tackle the uncertainty in decision making problems.

Since single valued neutrosophic model is more general than the fuzzy sets, IFS, PyFS, PFS and SFSs due to the wider range of applicability over different complex problems. The SVN sets can explain uncertainties concurrently more precisely than the other existing methods, like fuzzy set and existing structures.^{16,27,36}

The motivation of developed AOs is summarized as below.

(1) A very difficult MCDM problem is the estimation of the supreme option in a single valued neutrosophic environment due to the involvement of several imprecise factors. Assessment of information in different MCDM techniques is simply depicted through fuzzy and their existing structures which may not consider all the data in a real-world problem.

(2) As a general theory, single valued neutrosophic numbers describes efficient execution in the assessment process about uncertain, imprecise and vague information. Thus, single valued neutrosophic theory provide an excellent approach for the assessment of objects under multinary data.

(3) In view of the fact that sine trigonometric hybrid AOs are simple but provide a pioneering tool for solving MCDM problems when combine with other powerful mathematical tools, this article aims to develop sine trigonometric hybrid AOs in a single valued neutrosophic environment to handle complex problems.

(4) A single valued neutrosophic model is different from the mathematical tools like fuzzy sets and their extensions. Because the fuzzy set and their extensions can only handle one dimensional data, two dimensional data and three dimensional data, respectively, which may prompt a loss in data. Nevertheless, in many daily life problems, we handle the situations having higher dimension to sort out all the attributes.

(5) The sine trigonometric hybrid AOs employed in the construction of SVN sine trigonometric hybrid AOs are more suitable than all other aggregation approaches to tackle the MCDM situations as developed AOs have ability to consider all the information within the aggregation procedure.

(6) sine trigonometric AOs make the optimal outcomes more accurate and definite when utilized in practical MCDM problems under single valued neutrosophic environment. However, the proposed single valued neutrosophic operators handle the drawbacks of AOs present in the literature.

Therefore, some single valued neutrosophic sine trigonometric hybrid AOs are developed to choose the best option in different decision-making situations. The developed operators has some advantages over other approaches which are given as below:

(1) Our proposed methods explain the problems more accurately which involve multiple attributes because they consider single valued neutrosophic numbers.

(2) The developed AOs are more precise and efficient with single attribute.

(3) To solve practical problems by using sine trigonometric hybrid AOs with single valued neutrosophic numbers is very significant.

The rest of this article is structured as follows: Section 2 recalls some fundamental definitions and operations of the SVNNSs. Section 3 presents novel sine hybrid aggregation operators. Section 4 develops a methodology of these AOs to MCDM problems under single valued neutrosophic environment. Section 5 discusses a application of the selection of best agricultural land. Section 6 provides comparative analysis of developed approaches with different aggregating methodologies. Section 7 discusses the conclusions and future directions.

2 Preliminaries

In this section some essential notions of PFS, SFS, and SVNNS are examined.

Definition 2.1. ¹⁷A PFS \mathcal{U} in fixed universe set \mathfrak{J} is defined as

$$\mathcal{U} = \{ \langle b, \tilde{\rho}_\varphi(b), \neg_\varphi(b), \tilde{n}_\varphi(b) \rangle \mid b \in \mathfrak{J} \},$$

where the positive, neutral and negative membership grades, $\tilde{\rho}_\varphi : \mathfrak{J} \rightarrow \Theta$, $\neg_\varphi : \mathfrak{J} \rightarrow \Theta$ and $\tilde{n}_\varphi : \mathfrak{J} \rightarrow \Theta$, respectively and also, $\Theta = [0, 1]$ is the unit interval. Furthermore, $0 \leq \tilde{\rho}_\varphi(b) + \neg_\varphi(b) + \tilde{n}_\varphi(b) \leq 1$, for each $b \in \mathfrak{J}$.

Definition 2.2. ^{1,2}A SFS \mathcal{U} in fixed universe set \mathfrak{J} is defined as

$$\mathcal{U} = \{ \langle b, \tilde{\rho}_\varphi(b), \neg_\varphi(b), \tilde{n}_\varphi(b) \rangle \mid b \in \mathfrak{J} \},$$

where the positive, neutral and negative membership grades, $\tilde{\rho}_\varphi : \mathfrak{J} \rightarrow \Theta$, $\neg_\varphi : \mathfrak{J} \rightarrow \Theta$ and $\tilde{n}_\varphi : \mathfrak{J} \rightarrow \Theta$, respectively and also, $\Theta = [0, 1]$ is the unit interval. Furthermore, $0 \leq \tilde{\rho}_\varphi^2(b) + \neg_\varphi^2(b) + \tilde{n}_\varphi^2(b) \leq 1$, for each $b \in \mathfrak{J}$.

Definition 2.3. ³⁸A neutrosophic set \mathcal{U} in a fixed universe set \mathfrak{J} is defined as

$$\mathcal{U} = \{ \langle b, \tilde{\rho}_\varphi(b), \neg_\varphi(b), \tilde{n}_\varphi(b) \rangle \mid b \in \mathfrak{J} \},$$

where the positive, neutral and negative membership grades, $\tilde{\rho}_\varphi : \mathfrak{J} \rightarrow]0^-, 1^+[$, $\neg_\varphi : \mathfrak{J} \rightarrow]0^-, 1^+[$ and $\tilde{n}_\varphi : \mathfrak{J} \rightarrow]0^-, 1^+[$, respectively. Furthermore, $0^- \leq \tilde{\rho}_\varphi(b) + \neg_\varphi(b) + \tilde{n}_\varphi(b) \leq 3^+$, for each $b \in \mathfrak{J}$.

Definition 2.4. ⁴⁰A SVNS \mathcal{U} in a fixed universe set \mathfrak{J} is defined as

$$\mathcal{U} = \{ \langle b, \tilde{\rho}_\varphi(b), \neg_\varphi(b), \tilde{n}_\varphi(b) \rangle \mid b \in \mathfrak{J} \},$$

where the positive, neutral and negative membership grades, $\tilde{\rho}_\varphi : \mathfrak{J} \rightarrow \Theta$, $\neg_\varphi : \mathfrak{J} \rightarrow \Theta$ and $\tilde{n}_\varphi : \mathfrak{J} \rightarrow \Theta$, respectively and also, $\Theta = [0, 1]$ is the unit interval. Furthermore, $0 \leq \tilde{\rho}_\varphi(b) + \neg_\varphi(b) + \tilde{n}_\varphi(b) \leq 3$, for each $b \in \mathfrak{J}$.

In simplification, throughout the whole work, the triplet $\mathcal{U} = \{ \tilde{\rho}_\varphi, \neg_\varphi, \tilde{n}_\varphi \}$ called single valued neutrosophic number (SVNN) and their collection denoted by $SVNN(\mathfrak{J})$.

Here we highlight the some basic operations for SVNNs, ^{40,44} & ⁵⁰ as follows:

Definition 2.5. ⁵⁰ Let $\mathcal{U}_1 = \{ \tilde{\rho}_{\varphi_1}, \neg_{\varphi_1}, \tilde{n}_{\varphi_1} \}$ and $\mathcal{U}_2 = \{ \tilde{\rho}_{\varphi_2}, \neg_{\varphi_2}, \tilde{n}_{\varphi_2} \} \in SVNN(\mathfrak{J})$. then,

- (1) $\mathcal{U}_1 \subseteq \mathcal{U}_2$ if and only if $\tilde{\rho}_{\varphi_1} \leq \tilde{\rho}_{\varphi_2}$, $\neg_{\varphi_1} \geq \neg_{\varphi_2}$ and $\tilde{n}_{\varphi_1} \geq \tilde{n}_{\varphi_2}$ for each $b \in \mathfrak{J}$.
- (2) $\mathcal{U}_1 = \mathcal{U}_2$ if and only if $\mathcal{U}_1 \subseteq \mathcal{U}_2$ and $\mathcal{U}_2 \subseteq \mathcal{U}_1$.
- (2) $\mathcal{U}_1 \cap \mathcal{U}_2 = \{ \min(\tilde{\rho}_{\varphi_1}, \tilde{\rho}_{\varphi_2}), \max(\neg_{\varphi_1}, \neg_{\varphi_2}), \max(\tilde{n}_{\varphi_1}, \tilde{n}_{\varphi_2}) \}$,
- (3) $\mathcal{U}_1 \cup \mathcal{U}_2 = \{ \max(\tilde{\rho}_{\varphi_1}, \tilde{\rho}_{\varphi_2}), \min(\neg_{\varphi_1}, \neg_{\varphi_2}), \min(\tilde{n}_{\varphi_1}, \tilde{n}_{\varphi_2}) \}$,
- (4) $\mathcal{U}_1^c = \{ \tilde{n}_{\varphi_1}, \neg_{\varphi_1}, \tilde{\rho}_{\varphi_1} \}$.

Definition 2.6. ^{40,44} Let $\mathcal{U}_1 = \{ \tilde{\rho}_{\varphi_1}, \neg_{\varphi_1}, \tilde{n}_{\varphi_1} \}$ and $\mathcal{U}_2 = \{ \tilde{\rho}_{\varphi_2}, \neg_{\varphi_2}, \tilde{n}_{\varphi_2} \} \in SVNN(\mathfrak{J})$ with $\vartheta > 0$. then,

- (1) $\mathcal{U}_1 \otimes \mathcal{U}_2 = \{ \tilde{\rho}_{\varphi_1} \tilde{\rho}_{\varphi_2}, \neg_{\varphi_1} + \neg_{\varphi_2} - \neg_{\varphi_1} \cdot \neg_{\varphi_2}, \tilde{n}_{\varphi_1} + \tilde{n}_{\varphi_2} - \tilde{n}_{\varphi_1} \cdot \tilde{n}_{\varphi_2} \}$;
- (2) $\mathcal{U}_1 \oplus \mathcal{U}_2 = \{ \tilde{\rho}_{\varphi_1} + \tilde{\rho}_{\varphi_2} - \tilde{\rho}_{\varphi_1} \tilde{\rho}_{\varphi_2}, \neg_{\varphi_1} \neg_{\varphi_2}, \tilde{n}_{\varphi_1} \tilde{n}_{\varphi_2} \}$;
- (3) $(\mathcal{U}_1)^\vartheta = \{ (\tilde{\rho}_{\varphi_1})^\vartheta, 1 - (1 - \neg_{\varphi_1})^\vartheta, 1 - (1 - \tilde{n}_{\varphi_1})^\vartheta \}$;
- (4) $\vartheta \cdot \mathcal{U}_1 = \{ 1 - (1 - \tilde{\rho}_{\varphi_1})^\vartheta, (\neg_{\varphi_1})^\vartheta, (\tilde{n}_{\varphi_1})^\vartheta \}$.

Definition 2.7. ⁴ Let $\mathcal{U}_h = \{ \tilde{\rho}_{\varphi_h}, \neg_{\varphi_h}, \tilde{n}_{\varphi_h} \} \in SVNN(\mathfrak{J})$ ($h = 1, 2, 3, \dots, n$). Then, the Algebraic averaging aggregation operator for $SVNN(\mathfrak{J})$ is denoted by $SVNWA$ and defined as follows:

$$\begin{aligned} SVNWA(\mathcal{U}_1, \mathcal{U}_2, \mathcal{U}_3, \dots, \mathcal{U}_n) &= \sum_{h=1}^n \vartheta_h \mathcal{U}_h, \\ &= \{ 1 - \prod_{h=1}^n (1 - \tilde{\rho}_{\varphi_h})^{\vartheta_h}, \prod_{h=1}^n (\neg_{\varphi_h})^{\vartheta_h}, \prod_{h=1}^n (\tilde{n}_{\varphi_h})^{\vartheta_h} \} \end{aligned}$$

where the weights of \mathcal{U}_h represented by ϑ_h ($h = 1, 2, \dots, n$) having $\vartheta_h \geq 0$ and $\sum_{h=1}^n \vartheta_h = 1$.

Definition 2.8. ⁴ Let $\mathcal{U}_h = \{ \tilde{\rho}_{\varphi_h}, \neg_{\varphi_h}, \tilde{n}_{\varphi_h} \} \in SVNN(\mathfrak{J})$ ($h = 1, 2, 3, \dots, n$). Then, the Algebraic geometric aggregation operator for $SVNN(\mathfrak{J})$ is denoted by $SVNWG$ and defined as follows:

$$\begin{aligned} SFWG(\mathcal{U}_1, \mathcal{U}_2, \mathcal{U}_3, \dots, \mathcal{U}_n) &= \prod_{h=1}^n (\mathcal{U}_h)^{\vartheta_h}, \\ &= \{ \prod_{h=1}^n (\tilde{\rho}_{\varphi_h})^{\vartheta_h}, 1 - \prod_{h=1}^n (1 - \neg_{\varphi_h})^{\vartheta_h}, 1 - \prod_{h=1}^n (1 - \tilde{n}_{\varphi_h})^{\vartheta_h} \} \end{aligned}$$

where ϑ_h ($h = 1, 2, \dots, n$) represents the weights of \mathcal{U}_h ($h = 1, 2, 3, \dots, n$) with $\vartheta_h \geq 0$ and $\sum_{h=1}^n \vartheta_h = 1$.

Ashraf et al.,⁴ introduced the sine trigonometric operational laws for single valued neutrosophic environments.

Definition 2.9. ⁴ Let $\mathcal{U} = \{\tilde{\rho}_\varphi, \ulcorner_\varphi, \tilde{n}_\varphi\} \in SVNN(\mathfrak{J})$. If

$$\sin(\mathcal{U}) = \left\{ \left(\ulcorner, \sin\left(\frac{\pi}{2}\tilde{\rho}_\varphi(\mathfrak{J})\right), 1 - \sin\left(\frac{\pi}{2}(1 - \ulcorner_\varphi(\mathfrak{J}))\right), 1 - \sin\left(\frac{\pi}{2}(1 - \tilde{n}_\varphi(\mathfrak{J}))\right) \right) \mid \mathfrak{b} \in \mathfrak{J} \right\}$$

then, $\sin(\mathcal{U})$ is said to be sine trigonometric operator and their value known to be sine trigonometric SVNN (ST-SVNN).

Definition 2.10. ⁴ Let $\sin(\mathcal{U}_1) = \left\{ \left(\begin{matrix} \sin\left(\frac{\pi}{2}\tilde{\rho}_{\varphi_1}\right), \\ 1 - \sin\left(\frac{\pi}{2}(1 - \ulcorner_{\varphi_1})\right), \\ 1 - \sin\left(\frac{\pi}{2}(1 - \tilde{n}_{\varphi_1})\right) \end{matrix} \right) \right\}$ and $\sin(\mathcal{U}_2) = \left\{ \left(\begin{matrix} \sin\left(\frac{\pi}{2}\tilde{\rho}_{\varphi_2}\right), \\ 1 - \sin\left(\frac{\pi}{2}(1 - \ulcorner_{\varphi_2})\right), \\ 1 - \sin\left(\frac{\pi}{2}(1 - \tilde{n}_{\varphi_2})\right) \end{matrix} \right) \right\}$

be two ST-SVNNs. The basic operations for SVNNs are as follows

- (1) $\sin(\mathcal{U}_1) \oplus \sin(\mathcal{U}_2) = \left(\begin{matrix} 1 - (1 - \sin\left(\frac{\pi}{2}\tilde{\rho}_{\varphi_1}\right))(1 - \sin\left(\frac{\pi}{2}\tilde{\rho}_{\varphi_2}\right)), \\ (1 - \sin\left(\frac{\pi}{2}(1 - \ulcorner_{\varphi_1})\right))(1 - \sin\left(\frac{\pi}{2}(1 - \ulcorner_{\varphi_2})\right)), \\ (1 - \sin\left(\frac{\pi}{2}(1 - \tilde{n}_{\varphi_1})\right))(1 - \sin\left(\frac{\pi}{2}(1 - \tilde{n}_{\varphi_2})\right)) \end{matrix} \right),$
- (2) $\psi \cdot \sin(\mathcal{U}_1) = \left(\begin{matrix} 1 - (1 - \sin\left(\frac{\pi}{2}\tilde{\rho}_{\varphi_1}\right))^\psi, (1 - \sin\left(\frac{\pi}{2}(1 - \ulcorner_{\varphi_1})\right))^\psi, \\ (1 - \sin\left(\frac{\pi}{2}(1 - \tilde{n}_{\varphi_1})\right))^\psi \end{matrix} \right),$
- (3) $\sin(\mathcal{U}_1) \otimes \sin(\mathcal{U}_2) = \left(\begin{matrix} \sin\left(\frac{\pi}{2}\tilde{\rho}_{\varphi_1}\right)\sin\left(\frac{\pi}{2}\tilde{\rho}_{\varphi_2}\right), \\ 1 - (\sin\left(\frac{\pi}{2}(1 - \ulcorner_{\varphi_1})\right))(\sin\left(\frac{\pi}{2}(1 - \ulcorner_{\varphi_2})\right)), \\ 1 - (\sin\left(\frac{\pi}{2}(1 - \tilde{n}_{\varphi_1})\right))(\sin\left(\frac{\pi}{2}(1 - \tilde{n}_{\varphi_2})\right)) \end{matrix} \right),$
- (4) $(\sin(\mathcal{U}_1))^\psi = \left(\begin{matrix} (\sin\left(\frac{\pi}{2}\tilde{\rho}_{\varphi_1}\right))^\psi, 1 - (\sin\left(\frac{\pi}{2}(1 - \ulcorner_{\varphi_1})\right))^\psi, \\ 1 - (\sin\left(\frac{\pi}{2}(1 - \tilde{n}_{\varphi_1})\right))^\psi \end{matrix} \right).$

Definition 2.11. Let $\mathcal{U} = \{\tilde{\rho}_\varphi, \ulcorner_\varphi, \tilde{n}_\varphi\} \in SVNN(\mathfrak{J})$. Then, the score and accuracy of \mathcal{U} is denoted and defined as

- (1) $\check{s}\check{c}(\mathcal{U}) = \tilde{\rho}_\varphi - \ulcorner_\varphi - \tilde{n}_\varphi$, and
- (2) $\check{a}\check{c}(\mathcal{U}) = \tilde{\rho}_\varphi + \ulcorner_\varphi + \tilde{n}_\varphi$.

Definition 2.12. Let $\mathcal{U}_1 = \{\tilde{\rho}_{\varphi_1}, \ulcorner_{\varphi_1}, \tilde{n}_{\varphi_1}\}$ and $\mathcal{U}_2 = \{\tilde{\rho}_{\varphi_2}, \ulcorner_{\varphi_2}, \tilde{n}_{\varphi_2}\} \in SVNN(\mathfrak{J})$. Then,

- (1) If $\check{s}\check{c}(\mathcal{U}_1) < \check{s}\check{c}(\mathcal{U}_2)$ then $\mathcal{U}_1 < \mathcal{U}_2$,
- (2) If $\check{s}\check{c}(\mathcal{U}_1) > \check{s}\check{c}(\mathcal{U}_2)$ then $\mathcal{U}_1 > \mathcal{U}_2$,
- (3) If $\check{s}\check{c}(\mathcal{U}_1) = \check{s}\check{c}(\mathcal{U}_2)$ then
 - (a) $\check{a}\check{c}(\mathcal{U}_1) < \check{a}\check{c}(\mathcal{U}_2)$ then $\mathcal{U}_1 < \mathcal{U}_2$,
 - (b) $\check{a}\check{c}(\mathcal{U}_1) > \check{a}\check{c}(\mathcal{U}_2)$ then $\mathcal{U}_1 > \mathcal{U}_2$,
 - (c) $\check{a}\check{c}(\mathcal{U}_1) = \check{a}\check{c}(\mathcal{U}_2)$ then $\mathcal{U}_1 = \mathcal{U}_2$.

3 Novel Sine Trigonometric Hybrid AOs for SVNNs

This section propose the novel sine trigonometric hybrid AOs under SVN information.

Definition 3.1. Let $\mathcal{U}_{\tilde{h}} = \{\tilde{\rho}_{\varphi_{\tilde{h}}}(b), \ulcorner_{\varphi_{\tilde{h}}}(b), \tilde{n}_{\varphi_{\tilde{h}}}(b)\} \in SVNN(\mathfrak{J})$ ($\tilde{h} \in \mathbb{N}$). The sine trigonometric hybrid weighted averaging AO for $SVNN(\mathfrak{J})$ is represented by $ST - SVNHWA$ and discribed as follows:

$$\begin{aligned} ST - SVNHWA(\mathcal{U}_1, \mathcal{U}_2, \dots, \mathcal{U}_n) &= \sigma_1 \sin(\mathcal{U}'_{v(1)}) \oplus \sigma_2 \sin(\mathcal{U}'_{v(2)}) \oplus \dots \oplus \sigma_n \sin(\mathcal{U}'_{v(n)}) \\ &= \sum_{\tilde{h}=1}^n \sigma_{\tilde{h}} \sin(\mathcal{U}'_{v(\tilde{h})}), \end{aligned}$$

where the weights of $\mathcal{U}_{\tilde{h}}$ is represented by $\varrho_{\tilde{h}}$ ($\tilde{h} = 1, 2, \dots, n$) having $\varrho_{\tilde{h}} \geq 0$ and $\sum_{\tilde{h}=1}^n \varrho_{\tilde{h}} = 1$ and gth biggest weighted value is $\mathcal{U}'_{v(\tilde{h})}$ ($\mathcal{U}'_{v(\tilde{h})} = n\varrho_{\tilde{h}}\mathcal{U}_{v(\tilde{h})} \mid \tilde{h} = 1, 2, \dots, n$) consequently by total order ($v(1), v(2), v(3), \dots, v(n)$). Also the associated weight vector $\sigma = \sigma_{\tilde{h}}$ ($\tilde{h} = 1, 2, \dots, n$) with $\sigma_{\tilde{h}} \geq 0$ and $\sum_{\tilde{h}=1}^n \sigma_{\tilde{h}} = 1$.

Theorem 3.2. Let $\mathcal{U}_{\tilde{h}} = \{\tilde{\rho}_{\varphi_{\tilde{h}}}(b), \ulcorner_{\varphi_{\tilde{h}}}(b), \tilde{n}_{\varphi_{\tilde{h}}}(b)\} \in SVNN(\mathfrak{J})$ ($\tilde{h} \in \mathbb{N}$) and the weights of $\mathcal{U}_{\tilde{h}}$ represented by $(\varrho_1, \varrho_2, \dots, \varrho_n)^T$ subject to $\sum_{\tilde{h}=1}^n \varrho_{\tilde{h}} = 1$. The $ST - SVNHWA$ AO is defined by a mapping $G^n \rightarrow G$

with associated weight vector $\sigma_{\bar{h}}$ ($\bar{h} = 1, 2, \dots, n$) having $\sigma_{\bar{h}} \geq 0$ and $\sum_{\bar{h}=1}^n \sigma_{\bar{h}} = 1$:

$$\begin{aligned}
 ST - SVNHWA(\mathcal{U}_1, \mathcal{U}_2, \dots, \mathcal{U}_n) &= \sum_{\bar{h}=1}^n \sigma_{\bar{h}} \sin(\mathcal{U}'_{v(\bar{h})}) \\
 &= \left(\begin{array}{c} 1 - \prod_{\bar{h}=1}^n \left(1 - \sin\left(\frac{\pi}{2} \tilde{\rho}'_{\varphi_{v(\bar{h})}}\right) \right)^{\sigma_{\bar{h}}}, \\ \prod_{\bar{h}=1}^n \left(1 - \sin\left(\frac{\pi}{2} 1 - \mathcal{T}'_{\varphi_{v(\bar{h})}}\right) \right)^{\sigma_{\bar{h}}}, \\ \prod_{\bar{h}=1}^n \left(1 - \sin\left(\frac{\pi}{2} 1 - \tilde{n}'_{\varphi_{v(\bar{h})}}\right) \right)^{\sigma_{\bar{h}}} \end{array} \right) \quad (1)
 \end{aligned}$$

Proof. By using mathematical induction on n we prove Theorem 3.2. Then the mathematical induction steps below were implemented.

Step-1: For $n = 2$, we obtained

$$ST - SVNHWA(\mathcal{U}_1, \mathcal{U}_2) = \sigma_1 \sin(\mathcal{U}'_{v(1)}) \oplus \sigma_2 \sin(\mathcal{U}'_{v(2)}).$$

Since by the Definition 2.9, $\sin(\mathcal{U}_1)$ and $\sin(\mathcal{U}_2)$ are SVNNS and hence $\sigma_1 \sin(\mathcal{U}'_{v(1)}) \oplus \sigma_2 \sin(\mathcal{U}'_{v(2)})$ is also SVNNS. Further, for \mathcal{U}_1 and \mathcal{U}_2 , we have

$$\begin{aligned}
 &ST - SVNHWA(\mathcal{U}_1, \mathcal{U}_2) \\
 &= \sigma_1 \sin(\mathcal{U}'_{v(1)}) \oplus \sigma_2 \sin(\mathcal{U}'_{v(2)}) \\
 &= \left(\begin{array}{c} 1 - \left(1 - \sin\left(\frac{\pi}{2} \tilde{\rho}'_{\varphi_{v(1)}}\right) \right)^{\sigma_1}, \\ \left(1 - \sin\left(\frac{\pi}{2} 1 - \mathcal{T}'_{\varphi_{v(1)}}\right) \right)^{\sigma_1}, \\ \left(1 - \sin\left(\frac{\pi}{2} 1 - \tilde{n}'_{\varphi_{v(1)}}\right) \right)^{\sigma_1} \end{array} \right) \oplus \left(\begin{array}{c} 1 - \left(1 - \sin\left(\frac{\pi}{2} \tilde{\rho}'_{\varphi_{v(2)}}\right) \right)^{\sigma_2}, \\ \left(1 - \sin\left(\frac{\pi}{2} 1 - \mathcal{T}'_{\varphi_{v(2)}}\right) \right)^{\sigma_2}, \\ \left(1 - \sin\left(\frac{\pi}{2} 1 - \tilde{n}'_{\varphi_{v(2)}}\right) \right)^{\sigma_2} \end{array} \right) \\
 &= \left(\begin{array}{c} 1 - \prod_{\bar{h}=1}^2 \left(1 - \sin\left(\frac{\pi}{2} \tilde{\rho}'_{\varphi_{v(\bar{h})}}\right) \right)^{\sigma_{\bar{h}}}, \\ \prod_{\bar{h}=1}^2 \left(1 - \sin\left(\frac{\pi}{2} 1 - \mathcal{T}'_{\varphi_{v(\bar{h})}}\right) \right)^{\sigma_{\bar{h}}}, \\ \prod_{\bar{h}=1}^2 \left(1 - \sin\left(\frac{\pi}{2} 1 - \tilde{n}'_{\varphi_{v(\bar{h})}}\right) \right)^{\sigma_{\bar{h}}} \end{array} \right)
 \end{aligned}$$

Step-2: Suppose that Equation 1 is holds for $n = \kappa$, we have

$$ST - SVNHWA(\mathcal{U}_1, \mathcal{U}_2, \dots, \mathcal{U}_{\kappa}) = \left(\begin{array}{c} 1 - \prod_{\bar{h}=1}^{\kappa} \left(1 - \sin\left(\frac{\pi}{2} \tilde{\rho}'_{\varphi_{v(\bar{h})}}\right) \right)^{\sigma_{\bar{h}}}, \\ \prod_{\bar{h}=1}^{\kappa} \left(1 - \sin\left(\frac{\pi}{2} 1 - \mathcal{T}'_{\varphi_{v(\bar{h})}}\right) \right)^{\sigma_{\bar{h}}}, \\ \prod_{\bar{h}=1}^{\kappa} \left(1 - \sin\left(\frac{\pi}{2} 1 - \tilde{n}'_{\varphi_{v(\bar{h})}}\right) \right)^{\sigma_{\bar{h}}} \end{array} \right)$$

Step-3: Now we have to prove that Equation 1 is holds for $n = \kappa + 1$.

$$\begin{aligned}
 & ST - SVNHWA(\mathcal{U}_1, \mathcal{U}_2, \dots, \mathcal{U}_{\kappa+1}) \\
 &= \sum_{\hbar=1}^{\kappa} \sigma_{\hbar} \sin(\mathcal{U}'_{v(\hbar)}) \oplus \sigma_{\kappa+1} \sin(\mathcal{U}'_{v(\kappa+1)}) \\
 &= \left(\begin{array}{c} 1 - \prod_{\hbar=1}^{\kappa} \left(1 - \sin\left(\frac{\pi}{2} \tilde{\rho}'_{\varphi_{v(\hbar)}}\right) \right)^{\sigma_{\hbar}}, \\ \prod_{\hbar=1}^{\kappa} \left(1 - \sin\left(\frac{\pi}{2} 1 - \mathcal{T}'_{\varphi_{v(\hbar)}}\right) \right)^{\sigma_{\hbar}}, \\ \prod_{\hbar=1}^{\kappa} \left(1 - \sin\left(\frac{\pi}{2} 1 - \tilde{n}'_{\varphi_{v(\hbar)}}\right) \right)^{\sigma_{\hbar}} \end{array} \right) \oplus \left(\begin{array}{c} 1 - \left(1 - \sin\left(\frac{\pi}{2} \tilde{\rho}'_{\varphi_{v(\kappa+1)}}\right) \right)^{\sigma_{\kappa+1}}, \\ \left(1 - \sin\left(\frac{\pi}{2} 1 - \mathcal{T}'_{\varphi_{v(\kappa+1)}}\right) \right)^{\sigma_{\kappa+1}}, \\ \left(1 - \sin\left(\frac{\pi}{2} 1 - \tilde{n}'_{\varphi_{v(\kappa+1)}}\right) \right)^{\sigma_{\kappa+1}} \end{array} \right) \\
 &= \left(\begin{array}{c} 1 - \prod_{\hbar=1}^{\kappa+1} \left(1 - \sin\left(\frac{\pi}{2} \tilde{\rho}'_{\varphi_{v(\hbar)}}\right) \right)^{\sigma_{\hbar}}, \\ \prod_{\hbar=1}^{\kappa+1} \left(1 - \sin\left(\frac{\pi}{2} 1 - \mathcal{T}'_{\varphi_{v(\hbar)}}\right) \right)^{\sigma_{\hbar}}, \\ \prod_{\hbar=1}^{\kappa+1} \left(1 - \sin\left(\frac{\pi}{2} 1 - \tilde{n}'_{\varphi_{v(\hbar)}}\right) \right)^{\sigma_{\hbar}} \end{array} \right)
 \end{aligned}$$

that is, when $n = z + 1$, Equation 1 also holds.

Hence, Equation 1 holds for any n . The proof is completed. □

Next, we give some properties that are apparently carried by the proposed ST-SVNHWA aggregation operator.

(1) Let $\mathcal{U}_{\hbar} = \{\tilde{\rho}_{\varphi_{\hbar}}, \mathcal{T}_{\varphi_{\hbar}}, \tilde{n}_{\varphi_{\hbar}}\} \in SVN(\mathfrak{J})$ ($\hbar = 1, 2, 3, \dots, n$) such that $\mathcal{U}_{\hbar} = \mathcal{U}$. Then

$$ST - SVNHWA(\mathcal{U}_1, \mathcal{U}_2, \dots, \mathcal{U}_n) = \sin(\mathcal{U}).$$

(2) Let $\mathcal{U}_{\hbar} = \{\tilde{\rho}_{\varphi_{\hbar}}, \mathcal{T}_{\varphi_{\hbar}}, \tilde{n}_{\varphi_{\hbar}}\}$, $\mathcal{U}_{\hbar}^- = \{\min(\tilde{\rho}_{\varphi_{\hbar}}), \max(\mathcal{T}_{\varphi_{\hbar}}), \max(\tilde{n}_{\varphi_{\hbar}})\}$ and $\mathcal{U}_{\hbar}^+ = \{\max(\tilde{\rho}_{\varphi_{\hbar}}), \min(\mathcal{T}_{\varphi_{\hbar}}), \min(\tilde{n}_{\varphi_{\hbar}})\} \in SVN(\mathfrak{J})$ ($\hbar = 1, 2, 3, \dots, n$). Then,

$$\sin(\mathcal{U}_{\hbar}^-) \leq ST - SVNHWA(\mathcal{U}_1, \mathcal{U}_2, \dots, \mathcal{U}_n) \leq \sin(\mathcal{U}_{\hbar}^+).$$

(3) Let $\mathcal{U}_{\hbar} = \{\tilde{\rho}_{\varphi_{\hbar}}, \mathcal{T}_{\varphi_{\hbar}}, \tilde{n}_{\varphi_{\hbar}}\}$, $\mathcal{U}_{\hbar}^* = \{\tilde{\rho}_{\varphi_{\hbar}}^*, \mathcal{T}_{\varphi_{\hbar}}^*, \tilde{n}_{\varphi_{\hbar}}^*\} \in SVN(\mathfrak{J})$ ($\hbar = 1, 2, 3, \dots, n$). If $\tilde{\rho}_{\varphi_{\hbar}} \leq \tilde{\rho}_{\varphi_{\hbar}}^*$, $\mathcal{T}_{\varphi_{\hbar}} \leq \mathcal{T}_{\varphi_{\hbar}}^*$ and $\tilde{n}_{\varphi_{\hbar}} \leq \tilde{n}_{\varphi_{\hbar}}^*$, then

$$ST - SVNHWA(\mathcal{U}_1, \mathcal{U}_2, \dots, \mathcal{U}_n) \leq ST - SVNHWA(\mathcal{U}_1^*, \mathcal{U}_2^*, \dots, \mathcal{U}_n^*).$$

Definition 3.3. Let $\mathcal{U}_{\hbar} = \{\tilde{\rho}_{\varphi_{\hbar}}(b), \mathcal{T}_{\varphi_{\hbar}}(b), \tilde{n}_{\varphi_{\hbar}}(b)\} \in SVN(\mathfrak{J})$ ($\hbar \in \mathbb{N}$). The sine trigonometric hybrid weighted geometric AO for $SVNN(\mathfrak{J})$ is represented by $ST - SVNHWG$ and discribed as follows:

$$\begin{aligned}
 ST - SVNHWG(\mathcal{U}_1, \mathcal{U}_2, \dots, \mathcal{U}_n) &= \left(\sin(\mathcal{U}'_{v(1)}) \right)^{\sigma_1} \otimes \left(\sin(\mathcal{U}'_{v(2)}) \right)^{\sigma_2} \otimes \dots \otimes \left(\sin(\mathcal{U}'_{v(n)}) \right)^{\sigma_n} \\
 &= \prod_{\hbar=1}^n \left(\sin(\mathcal{U}'_{v(\hbar)}) \right)^{\sigma_{\hbar}}
 \end{aligned}$$

where the weights of \mathcal{U}_{\hbar} is represented by ϱ_{\hbar} ($\hbar = 1, 2, \dots, n$) having $\varrho_{\hbar} \geq 0$ and $\sum_{\hbar=1}^n \varrho_{\hbar} = 1$ and g th biggest weighted value is $\mathcal{U}'_{v(\hbar)}$ ($\mathcal{U}'_{v(\hbar)} = n\varrho_{\hbar}\mathcal{U}_{v(\hbar)} | \hbar = 1, 2, \dots, n$) consequently by total order $(v(1), v(2), v(3), \dots, v(n))$. Also the associated weight vector $\sigma = \sigma_{\hbar}$ ($\hbar = 1, 2, \dots, n$) with $\sigma_{\hbar} \geq 0$ and $\sum_{\hbar=1}^n \sigma_{\hbar} = 1$.

Theorem 3.4. Let $\mathcal{U}_{\hbar} = \{\tilde{\rho}_{\varphi_{\hbar}}(b), \mathcal{T}_{\varphi_{\hbar}}(b), \tilde{n}_{\varphi_{\hbar}}(b)\} \in SVN(\mathfrak{J})$ ($\hbar \in \mathbb{N}$) and the weights of \mathcal{U}_{\hbar} represented by $(\varrho_1, \varrho_2, \dots, \varrho_n)^T$ subject to $\sum_{\hbar=1}^n \varrho_{\hbar} = 1$. The $ST - SVNHWG$ AO is defined by a mapping $G^n \rightarrow G$ with associated weight vector σ_{\hbar} ($\hbar = 1, 2, \dots, n$) having $\sigma_{\hbar} \geq 0$ and $\sum_{\hbar=1}^n \sigma_{\hbar} = 1$:

$$\begin{aligned}
 ST - SVNHWG(\mathcal{U}_1, \mathcal{U}_2, \dots, \mathcal{U}_n) &= \prod_{\hbar=1}^n \left(\sin(\mathcal{U}'_{v(\hbar)}) \right)^{\sigma_{\hbar}} \\
 &= \left(\begin{array}{c} \prod_{\hbar=1}^n \left(\sin\left(\frac{\pi}{2} \tilde{\rho}'_{\varphi_{v(\hbar)}}\right) \right)^{\sigma_{\hbar}}, \\ 1 - \prod_{\hbar=1}^n \left(\sin\left(\frac{\pi}{2} 1 - \mathcal{T}'_{\varphi_{v(\hbar)}}\right) \right)^{\sigma_{\hbar}}, \\ 1 - \prod_{\hbar=1}^n \left(\sin\left(\frac{\pi}{2} 1 - \tilde{n}'_{\varphi_{v(\hbar)}}\right) \right)^{\sigma_{\hbar}} \end{array} \right) \tag{2}
 \end{aligned}$$

Proof. By using mathematical induction on n we prove Theorem 3.4. Then the mathematical induction steps below were implemented.

Step-1: For $n = 2$, we get

$$ST - SVNHWG(\mathcal{U}_1, \mathcal{U}_2) = \left(\sin(\mathcal{U}'_{v(1)})\right)^{\sigma_1} \otimes \left(\sin(\mathcal{U}'_{v(2)})\right)^{\sigma_2}.$$

Since by the Definition 2.9, $\sin(\mathcal{U}_1)$ and $\sin(\mathcal{U}_2)$ are SFNs and hence $\left(\sin(\mathcal{U}'_{v(1)})\right)^{\sigma_1} \otimes \left(\sin(\mathcal{U}'_{v(2)})\right)^{\sigma_2}$ is also SVN. Further, for \mathcal{U}_1 and \mathcal{U}_2 , we have

$$\begin{aligned} & ST - SVNHWG(\mathcal{U}_1, \mathcal{U}_2) \\ &= \left(\sin(\mathcal{U}'_{v(1)})\right)^{\sigma_1} \otimes \left(\sin(\mathcal{U}'_{v(2)})\right)^{\sigma_2} \\ &= \left(\begin{array}{c} \left(\sin\left(\frac{\pi}{2}\tilde{\rho}'_{\varphi_{v(1)}}\right)\right)^{\sigma_1}, \\ 1 - \left(\sin\left(\frac{\pi}{2}1 - \mathcal{T}'_{\varphi_{v(1)}}\right)\right)^{\sigma_1}, \\ 1 - \left(\sin\left(\frac{\pi}{2}1 - \tilde{n}'_{\varphi_{v(1)}}\right)\right)^{\sigma_1} \end{array} \right) \otimes \left(\begin{array}{c} \left(\sin\left(\frac{\pi}{2}\tilde{\rho}'_{\varphi_{v(2)}}\right)\right)^{\sigma_1}, \\ 1 - \left(\sin\left(\frac{\pi}{2}1 - \mathcal{T}'_{\varphi_{v(2)}}\right)\right)^{\sigma_1}, \\ 1 - \left(\sin\left(\frac{\pi}{2}1 - \tilde{n}'_{\varphi_{v(2)}}\right)\right)^{\sigma_1} \end{array} \right) \\ &= \left(\begin{array}{c} \prod_{\hbar=1}^2 \left(\sin\left(\frac{\pi}{2}\tilde{\rho}'_{\varphi_{v(\hbar)}}\right)\right)^{\sigma_{\hbar}}, \\ 1 - \prod_{\hbar=1}^2 \left(\sin\left(\frac{\pi}{2}1 - \mathcal{T}'_{\varphi_{v(\hbar)}}\right)\right)^{\sigma_{\hbar}}, \\ 1 - \prod_{\hbar=1}^2 \left(\sin\left(\frac{\pi}{2}1 - \tilde{n}'_{\varphi_{v(\hbar)}}\right)\right)^{\sigma_{\hbar}} \end{array} \right) \end{aligned}$$

Step-2: Suppose that Equation 2 is holds for $n = \kappa$, we have

$$ST - SVNHWG(\mathcal{U}_1, \mathcal{U}_2, \dots, \mathcal{U}_{\kappa}) = \left(\begin{array}{c} \prod_{\hbar=1}^{\kappa} \left(\sin\left(\frac{\pi}{2}\tilde{\rho}'_{\varphi_{v(\hbar)}}\right)\right)^{\sigma_{\hbar}}, \\ 1 - \prod_{\hbar=1}^{\kappa} \left(\sin\left(\frac{\pi}{2}1 - \mathcal{T}'_{\varphi_{v(\hbar)}}\right)\right)^{\sigma_{\hbar}}, \\ 1 - \prod_{\hbar=1}^{\kappa} \left(\sin\left(\frac{\pi}{2}1 - \tilde{n}'_{\varphi_{v(\hbar)}}\right)\right)^{\sigma_{\hbar}} \end{array} \right)$$

Step-3: Now we have to prove that Equation 2 is holds for $n = \kappa + 1$.

$$\begin{aligned} & ST - SVNHWG(\mathcal{U}_1, \mathcal{U}_2, \dots, \mathcal{U}_{\kappa+1}) \\ &= \prod_{\hbar=1}^{\kappa} \left(\sin(\mathcal{U}'_{v(\hbar)})\right)^{\sigma_{\hbar}} \otimes \left(\sin(\mathcal{U}'_{v(\kappa+1)})\right)^{\sigma_{\kappa+1}} \\ &= \left(\begin{array}{c} \prod_{\hbar=1}^{\kappa} \left(\sin\left(\frac{\pi}{2}\tilde{\rho}'_{\varphi_{v(\hbar)}}\right)\right)^{\sigma_{\hbar}}, \\ 1 - \prod_{\hbar=1}^{\kappa} \left(\sin\left(\frac{\pi}{2}1 - \mathcal{T}'_{\varphi_{v(\hbar)}}\right)\right)^{\sigma_{\hbar}}, \\ 1 - \prod_{\hbar=1}^{\kappa} \left(\sin\left(\frac{\pi}{2}1 - \tilde{n}'_{\varphi_{v(\hbar)}}\right)\right)^{\sigma_{\hbar}} \end{array} \right) \otimes \left(\begin{array}{c} \left(\sin\left(\frac{\pi}{2}\tilde{\rho}'_{\varphi_{v(\kappa+1)}}\right)\right)^{\sigma_{\kappa+1}}, \\ 1 - \left(\sin\left(\frac{\pi}{2}1 - \mathcal{T}'_{\varphi_{v(\kappa+1)}}\right)\right)^{\sigma_{\kappa+1}}, \\ 1 - \left(\sin\left(\frac{\pi}{2}1 - \tilde{n}'_{\varphi_{v(\kappa+1)}}\right)\right)^{\sigma_{\kappa+1}} \end{array} \right) \\ &= \left(\begin{array}{c} \prod_{\hbar=1}^{\kappa+1} \left(\sin\left(\frac{\pi}{2}\tilde{\rho}'_{\varphi_{v(\hbar)}}\right)\right)^{\sigma_{\hbar}}, \\ 1 - \prod_{\hbar=1}^{\kappa+1} \left(\sin\left(\frac{\pi}{2}1 - \mathcal{T}'_{\varphi_{v(\hbar)}}\right)\right)^{\sigma_{\hbar}}, \\ 1 - \prod_{\hbar=1}^{\kappa+1} \left(\sin\left(\frac{\pi}{2}1 - \tilde{n}'_{\varphi_{v(\hbar)}}\right)\right)^{\sigma_{\hbar}} \end{array} \right) \end{aligned}$$

that is, when $n = z + 1$, Equation 2 also holds.

Hence, Equation 2 holds for any n . The proof is completed. □

Next, we give some properties that are apparently carried by the proposed ST-SVNHWG aggregation operator.

(1) Let $\mathcal{U}_{\hbar} = \{\tilde{\rho}_{\varphi_{\hbar}}, \mathcal{T}_{\varphi_{\hbar}}, \tilde{n}_{\varphi_{\hbar}}\} \in SVN(\mathfrak{I})$ ($\hbar = 1, 2, 3, \dots, n$) such that $\mathcal{U}_{\hbar} = \mathcal{U}$. Then

$$ST - SVNHWG(\mathcal{U}_1, \mathcal{U}_2, \dots, \mathcal{U}_n) = \sin(\mathcal{U}).$$

(2) Let $\mathcal{U}_{\tilde{h}} = \{\tilde{\rho}_{\varphi_{\tilde{h}}}, \top_{\varphi_{\tilde{h}}}, \tilde{n}_{\varphi_{\tilde{h}}}\}$, $\mathcal{U}_{\tilde{h}}^- = \{\min(\tilde{\rho}_{\varphi_{\tilde{h}}}), \max(\top_{\varphi_{\tilde{h}}}), \max(\tilde{n}_{\varphi_{\tilde{h}}})\}$ and $\mathcal{U}_{\tilde{h}}^+ = \{\max(\tilde{\rho}_{\varphi_{\tilde{h}}}), \min(\top_{\varphi_{\tilde{h}}}), \min(\tilde{n}_{\varphi_{\tilde{h}}})\}$ $\in SVN(\mathfrak{I})$ ($\tilde{h} = 1, 2, 3, \dots, n$). Then,

$$\sin(\mathcal{U}_{\tilde{h}}^-) \leq ST - SVNHWG(\mathcal{U}_1, \mathcal{U}_2, \dots, \mathcal{U}_n) \leq \sin(\mathcal{U}_{\tilde{h}}^+).$$

(3) Let $\mathcal{U}_{\tilde{h}} = \{\tilde{\rho}_{\varphi_{\tilde{h}}}, \top_{\varphi_{\tilde{h}}}, \tilde{n}_{\varphi_{\tilde{h}}}\}$, $\mathcal{U}_{\tilde{h}}^* = \{\tilde{\rho}_{\varphi_{\tilde{h}}}^*, \top_{\varphi_{\tilde{h}}}^*, \tilde{n}_{\varphi_{\tilde{h}}}^*\} \in SVN(\mathfrak{I})$ ($\tilde{h} = 1, 2, 3, \dots, n$). If $\tilde{\rho}_{\varphi_{\tilde{h}}} \leq \tilde{\rho}_{\varphi_{\tilde{h}}}^*$, $\top_{\varphi_{\tilde{h}}} \leq \top_{\varphi_{\tilde{h}}}^*$ and $\tilde{n}_{\varphi_{\tilde{h}}} \leq \tilde{n}_{\varphi_{\tilde{h}}}^*$, then

$$ST - SVNHWG(\mathcal{U}_1, \mathcal{U}_2, \dots, \mathcal{U}_n) \leq ST - SVNHWG(\mathcal{U}_1^*, \mathcal{U}_2^*, \dots, \mathcal{U}_n^*).$$

4 Decision Making Algorithm

This section describes a decision-making algorithm to address the uncertainty of decision-making problems (DMPs) in the SVN environment, supported by an illustrative example. The following notions are utilized to tackle the MCDM situations having SVN information. Suppose that $\{\mathfrak{I}_1, \mathfrak{I}_2, \dots, \mathfrak{I}_k\}$ is a universal set and $\{r_1, r_2, r_3, \dots, r_n\}$ is the universe of attributes. Assume $\vartheta = \{\vartheta_1, \vartheta_2, \dots, \vartheta_n\}$ is a weigh-vector with $\vartheta_{\tilde{h}} \in [0, 1]$ such that $\sum_{\tilde{h}=1}^n \vartheta_{\tilde{h}} = 1$. Consider $D^{(k)} = (\mathcal{U}_{ij}^{(k)})_{k \times n}$ is a single valued neutrosophic decision-matrix, which represents the membership values evaluated by the experts.

We construct an algorithmic method to handle MCDM problems by proposed aggregation operators. The algorithm includes steps below:

Step-1 Summarize the values of each alternative in term of decision matrix $D^{(k)} = (\mathcal{U}_{ij}^{(k)})_{n \times m}$ with SVNns.

Step-2 Construct the normalized decision matrix $P = (p_{ij})$ from $D = (\mathcal{U}_{ij})$, where p_{ij} is calculated as

$$p_{ij} = \begin{cases} (\tilde{\rho}_{ij}, \top_{ij}, \tilde{n}_{ij}) & \text{If criteria are benefit type} \\ (\tilde{n}_{ij}, \top_{ij}, \tilde{\rho}_{ij}) & \text{If criteria are cost type} \end{cases} \tag{3}$$

Step-3 If the attribute weights are known as a prior then utilize them. Otherwise, we compute them by utilizing the concept of the entropy measure. For it, the information entropy of criteria t_j is computed as

$$En_j = \frac{1 + \frac{1}{n} \sum_{i=1}^n (\tilde{\rho}_{\varphi_{ij}} \log(\tilde{\rho}_{\varphi_{ij}}) + \top_{\varphi_{ij}} \log(\top_{\varphi_{ij}}) + \tilde{n}_{\varphi_{ij}} \log(\tilde{n}_{\varphi_{ij}}))}{\sum_{j=1}^m \left(1 + \frac{1}{n} \sum_{i=1}^n \tilde{\rho}_{\varphi_{ij}} \log(\tilde{\rho}_{\varphi_{ij}}) + \top_{\varphi_{ij}} \log(\top_{\varphi_{ij}}) + \tilde{n}_{\varphi_{ij}} \log(\tilde{n}_{\varphi_{ij}}) \right)}$$

Step-4 Using proposed aggregation operators defined in Theorem 3.2 & 3.4 and attributes weight vector, the aggregated single valued neutrosophic numbers of the each alternative $\{\mathcal{U}_1, \mathcal{U}_2, \mathcal{U}_3, \dots, \mathcal{U}_n\}$ are obtained.

Step-6 Evaluate the scores values $\check{s}\check{c}(\mathcal{U}_{\tilde{h}})$ using the Definition 2.11 of collective spherical fuzzy numbers $\mathcal{U}_{\tilde{h}}$ ($\tilde{h} = 1, 2, \dots, n$) and rank using the Definition 2.12 according the maximum score values.

Step-6 Select the optimal alternative according the maximum score value or accuracy degree.

5 Application Decision Making Algorithm

In this segment, the numerical implementation of agricultural land selection is used to demonstrate the MCDM methodology developed.

5.1 Practical case study

Agriculture is an important component of the Economic System of Pakistan. This area directly supports the population of the country and accounts for 26% of gross domestic product (GDP). The major agricultural crops are sugarcane, wheat, rice, cotton, vegetables and fruit. A businessman wants to invest in the agriculture sector and to look for appropriate land. The options in his brain are $\mathcal{Y}_1, \mathcal{Y}_2, \mathcal{Y}_3, \mathcal{Y}_4$ and \mathcal{Y}_5 . He consults to an expert to get his suggestion about the alternatives based on the following desired parameters:

- (1) Location (r_1),
- (2) Climate (r_2),
- (3) Fertility (r_3),
- (4) Price (r_4) and
- (5) Water Availability (r_5).

The expert was asked to use SVN data in this assessment and weights of the attributes are $(0.15, 0.28, 0.20, 0.22, 0.15)^T$. The expert's findings are summarized in Table-1:

Table-2.: Expert Information

$$D^1 = \begin{matrix} & r_1 & r_2 & r_3 & r_4 & r_5 \\ \begin{matrix} \mathcal{Y}_1 \\ \mathcal{Y}_2 \\ \mathcal{Y}_3 \\ \mathcal{Y}_4 \\ \mathcal{Y}_5 \end{matrix} & \left(\begin{matrix} (0.5, 0.3, 0.4) & (0.3, 0.2, 0.5) & (0.2, 0.2, 0.6) & (0.4, 0.2, 0.3) & (0.3, 0.3, 0.4) \\ (0.7, 0.1, 0.3) & (0.3, 0.2, 0.7) & (0.6, 0.3, 0.2) & (0.2, 0.4, 0.6) & (0.7, 0.1, 0.2) \\ (0.5, 0.3, 0.4) & (0.4, 0.2, 0.6) & (0.6, 0.1, 0.2) & (0.3, 0.1, 0.5) & (0.6, 0.4, 0.3) \\ (0.7, 0.3, 0.2) & (0.2, 0.2, 0.7) & (0.4, 0.5, 0.2) & (0.2, 0.2, 0.5) & (0.4, 0.5, 0.4) \\ (0.4, 0.1, 0.3) & (0.2, 0.1, 0.5) & (0.4, 0.1, 0.5) & (0.6, 0.3, 0.4) & (0.3, 0.2, 0.4) \end{matrix} \right) \end{matrix}$$

Step-2 The r_1, r_3 & r_5 are benefits type and r_2 & r_4 are cost type criteria. According to defined formula 3, the normalized decision matrix is summarized in Table-2:

Table-2.: Normalized Information

$$P = \begin{matrix} & r_1 & r_2 & r_3 & r_4 & r_5 \\ \begin{matrix} \mathcal{Y}_1 \\ \mathcal{Y}_2 \\ \mathcal{Y}_3 \\ \mathcal{Y}_4 \\ \mathcal{Y}_5 \end{matrix} & \left(\begin{matrix} (0.5, 0.3, 0.4) & (0.5, 0.2, 0.3) & (0.2, 0.2, 0.6) & (0.3, 0.2, 0.4) & (0.3, 0.3, 0.4) \\ (0.7, 0.1, 0.3) & (0.7, 0.2, 0.3) & (0.6, 0.3, 0.2) & (0.6, 0.4, 0.2) & (0.7, 0.1, 0.2) \\ (0.5, 0.3, 0.4) & (0.6, 0.2, 0.4) & (0.6, 0.1, 0.2) & (0.5, 0.1, 0.3) & (0.6, 0.4, 0.3) \\ (0.7, 0.3, 0.2) & (0.7, 0.2, 0.2) & (0.4, 0.5, 0.2) & (0.5, 0.2, 0.2) & (0.4, 0.5, 0.4) \\ (0.4, 0.1, 0.3) & (0.5, 0.1, 0.2) & (0.4, 0.1, 0.5) & (0.4, 0.3, 0.6) & (0.3, 0.2, 0.4) \end{matrix} \right) \end{matrix}$$

Step-3 The Expert provides the following parameters weights

$$\kappa = \{\kappa_1 = 0.15, \kappa_2 = 0.28, \kappa_3 = 0.20, \kappa_4 = 0.22, \kappa_5 = 0.15\}$$

Step-4 In this step, we used proposed AOp namely, ST-SVHWA and ST-SVHWG to aggregate the single valued neutrosophic information as follows:

Firstly, we find out the weighted matrix shown in Table-3:

Table-3.: $(n \mathcal{D}_{\bar{h}} \mathcal{U}_{\nu(\bar{h})} | \bar{h} = 1, 2, \dots, n)$

$$\left(\begin{matrix} (0.60, 0.09, 0.14) & (0.8, 0.004, 0.01) & (0.30, 0.02, 0.17) & (0.48, 0.01, 0.06) & (0.36, 0.09, 0.14) \\ (0.81, 0.01, 0.09) & (0.9, 0.004, 0.01) & (0.80, 0.04, 0.02) & (0.83, 0.06, 0.01) & (0.81, 0.01, 0.05) \\ (0.60, 0.09, 0.14) & (0.9, 0.004, 0.02) & (0.8, 0.004, 0.02) & (0.7, 0.002, 0.03) & (0.71, 0.14, 0.09) \\ (0.81, 0.09, 0.05) & (0.9, 0.004, 0.004) & (0.58, 0.12, 0.02) & (0.74, 0.01, 0.01) & (0.48, 0.20, 0.14) \\ (0.48, 0.01, 0.09) & (0.8, 0.005, 0.04) & (0.5, 0.004, 0.12) & (0.62, 0.03, 0.14) & (0.36, 0.05, 0.14) \end{matrix} \right)$$

Table-4.: Corresponding Score Values

$$\left(\begin{matrix} (0.3557) & (0.8037) & (0.1144) & (0.4113) & (0.1187) \\ (0.6943) & (0.9380) & (0.7444) & (0.7634) & (0.7384) \\ (0.3557) & (0.8687) & (0.7840) & (0.7052) & (0.4650) \\ (0.6601) & (0.9467) & (0.4454) & (0.7140) & (0.1296) \\ (0.3696) & (0.8160) & (0.4603) & (0.4433) & (0.1628) \end{matrix} \right)$$

Now, according to the score values, the $(U'_{v(\bar{h})} : \bar{h} = 1, 2, \dots, n)$ values are shows as follows:

Table-5.: $(U'_{v(\bar{h})} : \bar{h} = 1, 2, \dots, n)$

$U'_{v(11)} = U_{12} = (0.5, 0.2, 0.3)$	$U'_{v(12)} = U_{14} = (0.3, 0.2, 0.4)$	$U'_{v(13)} = U_{11} = (0.5, 0.3, 0.4)$
$U'_{v(14)} = U_{15} = (0.3, 0.3, 0.4)$	$U'_{v(15)} = U_{13} = (0.2, 0.2, 0.6)$	
$U'_{v(21)} = U_{22} = (0.7, 0.2, 0.3)$	$U'_{v(22)} = U_{24} = (0.6, 0.4, 0.2)$	$U'_{v(23)} = U_{23} = (0.6, 0.3, 0.2)$
$U'_{v(24)} = U_{25} = (0.7, 0.1, 0.2)$	$U'_{v(25)} = U_{21} = (0.7, 0.1, 0.3)$	
$U'_{v(31)} = U_{32} = (0.6, 0.2, 0.4)$	$U'_{v(32)} = U_{33} = (0.6, 0.1, 0.2)$	$U'_{v(33)} = U_{34} = (0.5, 0.1, 0.3)$
$U'_{v(34)} = U_{35} = (0.6, 0.4, 0.3)$	$U'_{v(35)} = U_{31} = (0.5, 0.3, 0.4)$	
$U'_{v(41)} = U_{42} = (0.7, 0.2, 0.2)$	$U'_{v(42)} = U_{44} = (0.5, 0.2, 0.2)$	$U'_{v(43)} = U_{41} = (0.7, 0.3, 0.2)$
$U'_{v(44)} = U_{43} = (0.4, 0.5, 0.2)$	$U'_{v(45)} = U_{45} = (0.4, 0.5, 0.2)$	
$U'_{v(51)} = U_{52} = (0.5, 0.1, 0.2)$	$U'_{v(52)} = U_{53} = (0.4, 0.1, 0.5)$	$U'_{v(53)} = U_{54} = (0.4, 0.3, 0.6)$
$U'_{v(54)} = U_{51} = (0.4, 0.1, 0.3)$	$U'_{v(55)} = U_{55} = (0.3, 0.2, 0.4)$	

Aggregated information of the alternatives are calculated in Table-6 as follows:

Table-6.: Aggregated Information

	<i>ST – SVNHWA</i>	<i>ST – SVNHWG</i>
\mathcal{Y}_1	(0.54515, 0.02797, 0.08164)	(0.50042, 0.03039, 0.08899)
\mathcal{Y}_2	(0.85733, 0.02063, 0.02539)	(0.85066, 0.03635, 0.02741)
\mathcal{Y}_3	(0.77818, 0.01567, 0.04227)	(0.77177, 0.02991, 0.04829)
\mathcal{Y}_4	(0.76483, 0.04584, 0.02450)	(0.71602, 0.06397, 0.02901)
\mathcal{Y}_5	(0.59151, 0.00952, 0.07508)	(0.58134, 0.01529, 0.09549)

Step-6 Score value of the aggregated alternatives are calculated in Table-7:

Table-7.: Score Values

	$\check{s}\check{c}(\mathcal{Y}_1)$	$\check{s}\check{c}(\mathcal{Y}_2)$	$\check{s}\check{c}(\mathcal{Y}_3)$	$\check{s}\check{c}(\mathcal{Y}_4)$	$\check{s}\check{c}(\mathcal{Y}_5)$
<i>ST – SVNHWA</i>	0.43553	0.81130	0.72023	0.69448	0.50690
<i>ST – SVNHWG</i>	0.38103	0.78688	0.69357	0.62302	0.47055

Step-7 Ranking of the alternatives are shown in Table-8:

Table-8.: Ranking

	Score Ranking	Best Alternative
<i>ST – SVNHWA</i>	$\check{s}\check{c}(\mathcal{Y}_2) > \check{s}\check{c}(\mathcal{Y}_3) > \check{s}\check{c}(\mathcal{Y}_4) > \check{s}\check{c}(\mathcal{Y}_5) > \check{s}\check{c}(\mathcal{Y}_1)$	\mathcal{Y}_2
<i>ST – SVNHWG</i>	$\check{s}\check{c}(\mathcal{Y}_2) > \check{s}\check{c}(\mathcal{Y}_3) > \check{s}\check{c}(\mathcal{Y}_4) > \check{s}\check{c}(\mathcal{Y}_5) > \check{s}\check{c}(\mathcal{Y}_1)$	\mathcal{Y}_2

From this above computational process, we can conclude that the alternative \mathcal{Y}_2 is the best among the others and hence it is highly recommendable to select for the required task/plan.

5.2 Comparison Study

In the section, we include some appropriate examples to demonstrate the feasibility and efficacy of the established decision-making approach and make a comparison with existing studies. Here, we presented the different existing aggregated information of the SVNNs namely, SVNWA,³³ SVNOWA,³³ NWA,⁴⁴ SVNFWA,¹⁸ SVNHWA,³² L-SVNWA,¹⁹ L-SVNOWA,¹⁹ ST-SVNWA⁴ and ST-SVNWG⁴ in Table-9, 10 and 11:

Table-9: Aggregated Information

	SVNWA ³³	SVNOWA ³³	NWA ⁴⁴	SVNFWA ¹⁸
\mathcal{Y}_1	(0.37, 0.22, 0.40)	(0.38, 0.24, 0.40)	(0.37, 0.23, 0.42)	(0.37, 0.22, 0.40)
\mathcal{Y}_2	(0.66, 0.20, 0.23)	(0.66, 0.18, 0.24)	(0.66, 0.24, 0.24)	(0.66, 0.20, 0.23)
\mathcal{Y}_3	(0.56, 0.17, 0.31)	(0.55, 0.18, 0.31)	(0.56, 0.21, 0.32)	(0.56, 0.17, 0.31)
\mathcal{Y}_4	(0.57, 0.29, 0.22)	(0.57, 0.31, 0.22)	(0.57, 0.33, 0.23)	(0.56, 0.29, 0.22)
\mathcal{Y}_5	(0.41, 0.14, 0.36)	(0.39, 0.13, 0.36)	(0.41, 0.16, 0.41)	(0.41, 0.14, 0.36)

Table-10: Aggregated Information

	SVNHWA ³² $\gamma = 2$	SVNHWA ³² $\gamma = 3$	L-SVNWA ¹⁹	L-SVNOWA ¹⁹
\mathcal{Y}_1	(0.37, 0.22, 0.40)	(0.36, 0.22, 0.40)	(0.31, 0.17, 0.35)	(0.32, 0.19, 0.36)
\mathcal{Y}_2	(0.66, 0.20, 0.23)	(0.66, 0.20, 0.23)	(0.64, 0.19, 0.23)	(0.65, 0.17, 0.23)
\mathcal{Y}_3	(0.56, 0.17, 0.31)	(0.56, 0.18, 0.31)	(0.49, 0.17, 0.33)	(0.48, 0.18, 0.33)
\mathcal{Y}_4	(0.56, 0.29, 0.22)	(0.56, 0.30, 0.22)	(0.55, 0.27, 0.19)	(0.55, 0.29, 0.19)
\mathcal{Y}_5	(0.41, 0.14, 0.36)	(0.41, 0.14, 0.37)	(0.28, 0.12, 0.37)	(0.24, 0.12, 0.38)

Table-11: Aggregated Information

	ST-SVNWA ⁴	ST-SVNOWA ⁴
\mathcal{Y}_1	(0.56, 0.02, 0.07)	(0.50, 0.02, 0.08)
\mathcal{Y}_2	(0.86, 0.02, 0.02)	(0.85, 0.03, 0.03)
\mathcal{Y}_3	(0.77, 0.01, 0.04)	(0.76, 0.02, 0.05)
\mathcal{Y}_4	(0.78, 0.04, 0.02)	(0.73, 0.06, 0.02)
\mathcal{Y}_5	(0.60, 0.009, 0.06)	(0.59, 0.01, 0.08)

Also, their corresponding ranking are calculated in Table-12:

Table-12: Ranking

Existing Operators	Ranking	Best Alternative
NWA ⁴³	$\mathcal{Y}_2 > \mathcal{Y}_3 > \mathcal{Y}_4 > \mathcal{Y}_5 > \mathcal{Y}_1$	\mathcal{Y}_2
SVNWA ³³	$\mathcal{Y}_2 > \mathcal{Y}_3 > \mathcal{Y}_4 > \mathcal{Y}_5 > \mathcal{Y}_1$	\mathcal{Y}_2
SVNOWA ³³	$\mathcal{Y}_2 > \mathcal{Y}_3 > \mathcal{Y}_4 > \mathcal{Y}_5 > \mathcal{Y}_1$	\mathcal{Y}_2
SVN WG ³³	$\mathcal{Y}_2 > \mathcal{Y}_3 > \mathcal{Y}_4 > \mathcal{Y}_5 > \mathcal{Y}_1$	\mathcal{Y}_2
SVNOWG ³³	$\mathcal{Y}_2 > \mathcal{Y}_3 > \mathcal{Y}_4 > \mathcal{Y}_5 > \mathcal{Y}_1$	\mathcal{Y}_2
SVNFWA ¹⁸	$\mathcal{Y}_2 > \mathcal{Y}_3 > \mathcal{Y}_4 > \mathcal{Y}_5 > \mathcal{Y}_1$	\mathcal{Y}_2
SVNHWA ³² $\gamma = 2$	$\mathcal{Y}_2 > \mathcal{Y}_3 > \mathcal{Y}_4 > \mathcal{Y}_5 > \mathcal{Y}_1$	\mathcal{Y}_2
SVNHWA ³² $\gamma = 3$	$\mathcal{Y}_2 > \mathcal{Y}_3 > \mathcal{Y}_4 > \mathcal{Y}_5 > \mathcal{Y}_1$	\mathcal{Y}_2
NWG ⁴³	$\mathcal{Y}_2 > \mathcal{Y}_3 > \mathcal{Y}_4 > \mathcal{Y}_5 > \mathcal{Y}_1$	\mathcal{Y}_2
SVNFWG ¹⁸	$\mathcal{Y}_2 > \mathcal{Y}_3 > \mathcal{Y}_4 > \mathcal{Y}_5 > \mathcal{Y}_1$	\mathcal{Y}_2
SVNHWG ³² $\gamma = 2$	$\mathcal{Y}_2 > \mathcal{Y}_3 > \mathcal{Y}_4 > \mathcal{Y}_5 > \mathcal{Y}_1$	\mathcal{Y}_2
SVNHWG ³² $\gamma = 3$	$\mathcal{Y}_2 > \mathcal{Y}_3 > \mathcal{Y}_4 > \mathcal{Y}_5 > \mathcal{Y}_1$	\mathcal{Y}_2
SNWEA ⁴⁴	$\mathcal{Y}_2 > \mathcal{Y}_3 > \mathcal{Y}_5 > \mathcal{Y}_4 > \mathcal{Y}_1$	\mathcal{Y}_2
L-SVNWA ¹⁹	$\mathcal{Y}_2 > \mathcal{Y}_4 > \mathcal{Y}_3 > \mathcal{Y}_5 > \mathcal{Y}_1$	\mathcal{Y}_2
L-SVNOWA ¹⁹	$\mathcal{Y}_2 > \mathcal{Y}_4 > \mathcal{Y}_3 > \mathcal{Y}_5 > \mathcal{Y}_1$	\mathcal{Y}_2
L-SVN WG ¹⁹	$\mathcal{Y}_2 > \mathcal{Y}_4 > \mathcal{Y}_3 > \mathcal{Y}_1 > \mathcal{Y}_5$	\mathcal{Y}_2
L-SVNOWG ¹⁹	$\mathcal{Y}_2 > \mathcal{Y}_3 > \mathcal{Y}_4 > \mathcal{Y}_5 > \mathcal{Y}_1$	\mathcal{Y}_2
L-SVNWA ⁴	$\mathcal{Y}_2 > \mathcal{Y}_4 > \mathcal{Y}_3 > \mathcal{Y}_5 > \mathcal{Y}_1$	\mathcal{Y}_2
L-SVN WG ⁴	$\mathcal{Y}_2 > \mathcal{Y}_3 > \mathcal{Y}_4 > \mathcal{Y}_5 > \mathcal{Y}_1$	\mathcal{Y}_2
L-SVNOWA ⁴	$\mathcal{Y}_2 > \mathcal{Y}_4 > \mathcal{Y}_3 > \mathcal{Y}_5 > \mathcal{Y}_1$	\mathcal{Y}_2
L-SVNOWG ⁴	$\mathcal{Y}_2 > \mathcal{Y}_3 > \mathcal{Y}_4 > \mathcal{Y}_5 > \mathcal{Y}_1$	\mathcal{Y}_2

Table-9.: Overall ranking of the alternatives

Proposed Operators	Ranking	Best Alternative
L-SVNHWA	$\mathcal{Y}_2 > \mathcal{Y}_3 > \mathcal{Y}_4 > \mathcal{Y}_5 > \mathcal{Y}_1$	\mathcal{Y}_2
L-SVNHWG	$\mathcal{Y}_2 > \mathcal{Y}_3 > \mathcal{Y}_4 > \mathcal{Y}_5 > \mathcal{Y}_1$	\mathcal{Y}_2

It is evident that the above conversation confirms the effectiveness and applicability of the proposed decision-making methodology based on new sine trigonometric aggregation operators under SVN environments.

6 Conclusion

Due to the existence of multiple attributes/criteria in many real-world problems, classical MCDM methods are not useful to tackle complicated decision-making situations. To overcome the difficulties of existing models, we have combined SVN Ss with sine trigonometric AOs.

In this article, we have discussed MCDM issues based on single valued neutrosophic information. Motivated by sine trigonometric function based operational laws, we have proposed different AOs, namely, ST-SVNHWA and ST-SVNHWG aggregation operators. We have investigated different features of these operators. We have employed these AOs to enlarged the applicability scope of MCDM. We have given real-life application for the selection of best agricultural land for best investment. At the end, we have provided a comparison of developed AOs with existing aggregation techniques in the literatures and authenticate the proposed strategy by effectiveness comparison study. In the future, we are extending our work in interval valued single valued neutrosophic environment.

Acknowledgments: In this section you can acknowledge any support given which is not covered by the author contribution or funding sections. This may include administrative and technical support, or donations in kind (e.g., materials used for experiments).

Conflicts of Interest: The authors declare no conflict of interest.

References

- [1] Ashraf, S., Abdullah, S. and Mahmood, T., "GRA method based on spherical linguistic fuzzy Choquet integral environment and its application in multi-attribute decision-making problems", *Mathematical Sciences*, 12(4), pp.263-275, 2018.
- [2] Ashraf, S. and Abdullah, S., "Spherical aggregation operators and their application in multiattribute group decision-making", *International Journal of Intelligent Systems*, 34(3), pp.493-523, 2019.
- [3] Ashraf, S., Abdullah, S., Mahmood, T., Ghani, F. and Mahmood, T., "Spherical fuzzy sets and their applications in multi-attribute decision making problems", *Journal of Intelligent & Fuzzy Systems*, 36(3), pp.2829-2844, 2019.
- [4] Ashraf, S., Abdullah, S., Zeng, S., Jin, H. and Ghani, F., "Fuzzy Decision Support Modeling for Hydrogen Power Plant Selection Based on Single Valued Neutrosophic Sine Trigonometric Aggregation Operators", *Symmetry*, 12(2), p.298, 2020.
- [5] Ashraf, S., Abdullah, S., Khan, S., "Fuzzy Decision support modeling for internet finance soft power evaluation Based on sine trigonometric Pythagorean fuzzy information", *Journal of Ambient Intelligence and Humanized Computing*, 2020. (Pre print)
- [6] Ashraf, S., Mahmood, T., Abdullah, S. and Khan, Q., "Different approaches to multi-criteria group decision making problems for picture fuzzy environment", *Bulletin of the Brazilian Mathematical Society, New Series*, 50(2), pp.373-397, 2019.
- [7] Ashraf, S., Abdullah, S., Mahmood, T. and Aslam, M., "Cleaner production evaluation in gold mines using novel distance measure method with cubic picture fuzzy numbers", *International Journal of Fuzzy Systems*, 21(8), pp.2448-2461, 2019.
- [8] Ashraf, S., Abdullah, S., Aslam, M., Qiyas, M. and Kutbi, M.A., "Spherical fuzzy sets and its representation of spherical fuzzy t-norms and t-conorms", *Journal of Intelligent & Fuzzy Systems*, 36(6), pp.6089-6102, 2019.
- [9] Ashraf, S., Abdullah, S. and Mahmood, T., "Spherical fuzzy Dombi aggregation operators and their application in group decision making problems", *Journal of Ambient Intelligence and Humanized Computing*, 11, pp.2731-2749, 2020.
- [10] Ashraf, S., Abdullah, S. and Aslam, M., "Symmetric sum based aggregation operators for spherical fuzzy information: Application in multi-attribute group decision making problem", *Journal of Intelligent & Fuzzy Systems*, 38(4), pp.5241-5255, 2020.
- [11] Ashraf, S., Abdullah, S. and Abdullah, L., "Child development influence environmental factors determined using spherical fuzzy distance measures", *Mathematics*, 7(8), p.661, 2019.
- [12] Ashraf, S., Abdullah, S. and Smarandache, F., "Logarithmic hybrid aggregation operators based on single valued neutrosophic sets and their applications in decision support systems", *Symmetry*, 11(3), p.364, 2019.
- [13] Atanassov, K.T., "Intuitionistic fuzzy sets", *Fuzzy Set and systems*, 20, pp.87-96, 1986.

- [14] Barukab, O., Abdullah, S., Ashraf, S., Arif, M. and Khan, S.A., "A New Approach to Fuzzy TOPSIS Method Based on Entropy Measure under Spherical Fuzzy Information", *Entropy*, 21(12), p.1231, 2019.
- [15] Batool, B., Ahmad, M., Abdullah, S., Ashraf, S. and Chinram, R., "Entropy Based Pythagorean Probabilistic Hesitant Fuzzy Decision Making Technique and Its Application for Fog-Haze Factor Assessment Problem", *Entropy*, 22(3), p.318, 2020.
- [16] Berti, L.A.C., "Application of the neutrosophic system to tax havens with a criminal approach", *International Journal of Neutrosophic Science*, 5(2), pp.91-106, 2020.
- [17] Cuong, B.C. and Kreinovich, V., "Picture Fuzzy Sets-a new concept for computational intelligence problems", In 2013 Third World Congress on Information and Communication Technologies (WICT 2013) (pp. 1-6), IEEE, 2013.
- [18] Garg, H., "Novel single-valued neutrosophic aggregated operators under Frank norm operation and its application to decision-making process", *International Journal for Uncertainty Quantification*, 6(4), pp.361-375, 2016.
- [19] Garg, H., "New logarithmic operational laws and their applications to multiattribute decision-making for single-valued neutrosophic numbers", *Cogn. Syst. Res.*, 52, pp.931-946, 2018.
- [20] Jin, H., Ashraf, S., Abdullah, S., Qiyas, M., Bano, M. and Zeng, S., "Linguistic spherical fuzzy aggregation operators and their applications in multi-attribute decision making problems", *Mathematics*, 7(5), p.413, 2019.
- [21] Jin, Y., Ashraf, S. and Abdullah, S., "Spherical fuzzy logarithmic aggregation operators based on entropy and their application in decision support systems", *Entropy*, 21(7), p.628, 2019.
- [22] Ji, P., Wang, J.Q. and Zhang, H.Y., "Frank prioritized Bonferroni mean operator with single-valued neutrosophic sets and its application in selecting third-party logistics providers", *Neural Computing and Applications*, 30(3), pp.799-823, 2018.
- [23] Khan, M.A., Ashraf, S., Abdullah, S. and Ghani, F., "Applications of probabilistic hesitant fuzzy rough set in decision support system", *Soft Computing*, 2020. <https://doi.org/10.1007/s00500-020-04971-z>
- [24] Khan, A.A., Ashraf, S., Abdullah, S., Qiyas, M., Luo, J. and Khan, S.U., "Pythagorean fuzzy Dombi aggregation operators and their application in decision support system", *Symmetry*, 11(3), p.383, 2019.
- [25] Khan, M.J., Kumam, P., Liu, P., Kumam, W. and Ashraf, S., "A novel approach to generalized intuitionistic fuzzy soft sets and its application in decision support system", *Mathematics*, 7(8), p.742, 2019.
- [26] Khan, S., Abdullah, S. and Ashraf, S., "Picture fuzzy aggregation information based on Einstein operations and their application in decision making", *Mathematical Sciences*, 13(3), pp.213-229, 2019.
- [27] Khan, M., Beg, I. and Gulistan, M., "Exponential Laws and Aggregation Operators on Neutrosophic Cubic Sets", *International Journal of Neutrosophic Science*, 4(1), pp.47-71, 2020.
- [28] Liu, P., Khan, Q., Mahmood, T., Smarandache, F. and Li, Y., "Multiple attribute group decision making based on 2-tuple linguistic neutrosophic Dombi power Heronian mean operators", *IEEE Access*, 7, pp.100205-100230, 2019.
- [29] Liu, P., Khan, Q. and Mahmood, T., "Some single-valued neutrosophic power muirhead mean operators and their application to group decision making", *Journal of Intelligent & Fuzzy Systems*, 37(2), pp.2515-2537, 2019.
- [30] Liu, P., Mahmood, T. and Khan, Q., "Group decision making based on power Heronian aggregation operators under linguistic neutrosophic environment", *International Journal of Fuzzy Systems*, 20(3), pp.970-985, 2018.
- [31] Liu, P., Khan, Q. and Mahmood, T., "Group decision making based on power Heronian aggregation operators under neutrosophic cubic environment", *Soft Computing*, 24(3), pp.1971-1997, 2020.
- [32] Liu, P., Chu, Y., Li, Y. and Chen, Y., "Some generalized neutrosophic number Hamacher aggregation operators and their application to group decision making", *International Journal of fuzzy systems*, 16(2), pp.242-255, 2014.

- [33] Peng, J.J., Wang, J.Q., Wang, J., Zhang, H.Y. and Chen, X.H., "Simplified neutrosophic sets and their applications in multi-criteria group decision-making problems", *International journal of systems science*, 47(10), pp.2342-2358, 2016.
- [34] Qiyas, M., Abdullah, S., Ashraf, S. and Abdullah, L., "Linguistic Picture Fuzzy Dombi Aggregation Operators and Their Application in Multiple Attribute Group Decision Making Problem", *Mathematics*, 7(8), p.764, 2019.
- [35] Qiyas, M., Abdullah, S., Ashraf, S. and Aslam, M., "Utilizing linguistic picture fuzzy aggregation operators for multiple-attribute decision-making problems", *International Journal of Fuzzy Systems*, 22(1), pp.310-320, 2020.
- [36] Rana, S., Saeed, M., Qayyum, M. and Smarandache, F., "Plithogenic Subjective Hyper-Super-Soft Matrices with New Definitions & Local, Global, Universal Subjective Ranking Model", *International Journal of Neutrosophic Science*, 6(2), pp.56-79, 2020.
- [37] Rafiq, M., Ashraf, S., Abdullah, S., Mahmood, T. and Muhammad, S., "The cosine similarity measures of spherical fuzzy sets and their applications in decision making", *Journal of Intelligent & Fuzzy Systems*, 36(6), pp.6059-6073, 2019.
- [38] Smarandache, F., "Neutrosophy Neutrosophic Probability, Set and Logic", American Research Press, (1998) Rehoboth, USA.
- [39] Sajjad Ali Khan, M., Abdullah, S., Yousaf Ali, M., Hussain, I. and Farooq, M., "Extension of TOPSIS method base on Choquet integral under interval-valued Pythagorean fuzzy environment", *Journal of Intelligent & Fuzzy Systems*, 34(1), pp.267-282, 2018.
- [40] Wang, H., Smarandache, F., Zhang, Y.Q. and Sunderraman, R., "Single valued neutrosophic sets", *Multispace and Multistructure*, 4, pp.410-413, 2010.
- [41] Xu, Z., "Intuitionistic fuzzy aggregation operators", *IEEE Transactions on fuzzy systems*, 15(6), pp.1179-1187, 2007.
- [42] Ye, J., "Multicriteria decision-making method using the correlation coefficient under single-valued neutrosophic environment", *International Journal of General Systems*, 42(4), pp.386-394, 2013.
- [43] Ye, J., "Exponential operations and aggregation operators of interval neutrosophic sets and their decision-making methods", *SpringerPlus* 2016, 5, 1488.
- [44] Ye, J., "Subtraction and division operations of simplified neutrosophic sets", *Information*, 8(51), 2017.
- [45] Yager, R.R., "On ordered weighted averaging aggregation operators in multi criteria decision making", *IEEE Transactions on systems, Man, and Cybernetics*, 18(1), pp.183-190, 1988.
- [46] Yager, R.R., "Pythagorean membership grades in multicriteria decision making", *IEEE Transactions on Fuzzy Systems*, 22(4), pp.958-965, 2013.
- [47] Zadeh, L.A., "Fuzzy sets", *Information and control*, 8(3), pp.338-353, 1965.
- [48] Zeng, S., Hussain, A., Mahmood, T., Irfan Ali, M., Ashraf, S. and Munir, M., "Covering-based spherical fuzzy rough set model hybrid with TOPSIS for multi-attribute decision-making", *Symmetry*, 11(4), p.547, 2019.
- [49] Zeng, S., Chen, S.M. and Kuo, L.W., "Multiattribute decision making based on novel score function of intuitionistic fuzzy values and modified VIKOR method", *Information Sciences*, 488, pp.76-92, 2019.
- [50] Zhang, X.; Bo, C.; Smarandache, F.; Dai, J., "New inclusion relation of neutrosophic sets with applications and related lattice structure", *Int. J. Mach. Learn. Cybern.*, 9, pp.1753-1763, 2018.